



## **Seismic Structure of Southern African Cratons**

### **A study based on teleseismic receiver functions and finite-frequency tomography**

Soliman, Mohammad Youssouf Ahmad; Artemieva, Irina; Levander, Alan; Thybo, Hans

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2014

*Document version*  
Early version, also known as pre-print

*Citation for published version (APA):*  
Soliman, M. Y. A., Artemieva, I., Levander, A., & Thybo, H. (2014). Seismic Structure of Southern African Cratons: A study based on teleseismic receiver functions and finite-frequency tomography. *Geophysical Research Abstracts*, 16, [10884].



## **Seismic Structure of Southern African Cratons: A study based on teleseismic receiver functions and finite-frequency tomography**

Mohammad Youssof (1,3), Irina Artemieva (1), Alan Levander (2), and Hans Thybo (1)

(1) Geology Section, IGN, University of Copenhagen, Copenhagen DK-1350, Denmark., (2) Department of Earth Science, Rice University, Houston, TX 77005-1892, USA., (3) Currently at Institute of Geophysics and Planetary Physics, Scripps Institution of Oceanography, University of California, San Diego, CA 92093-0225, USA.

Cratons are extremely stable continental crustal areas above thick depleted lithosphere. These regions have remained largely unchanged for more than 2.5 Ga. This study presents a new seismic model of the seismic structure of the crust and lithospheric mantle constrained by seismic receiver functions and finite-frequency tomography based on data from the South Africa Seismic Experiment (SASE). Combining the two methods provides high vertical and lateral resolution.

The main results obtained are (1) the presence of a highly heterogeneous crustal structure, in terms of thickness, composition (as shown by estimated  $V_p/V_s$ ), and sharpness of the discontinuities, (2) observation of an unexpectedly strong crustal azimuthal anisotropy, and (3) a fast lithospheric keel of the Kaapvaal Craton which reaches depths of 300-350 km and relatively slow anomalies beneath both the paleo-orogenic Limpopo Belt and modified Bushveld Complex down to depth of  $\sim 250$  km and  $\sim 150$  km, respectively.