



Københavns Universitet

The gut microbiota influence behavior in the subchronic PCP induced animal model of schizophrenia

Jørgensen, Bettina Merete Pyndt; Redrobe, Paul; Brønnum Pedersen, Tina ; Selfjord, Ellika Marie; Nielsen, Dennis Sandris; Plath, Niels; Hansen, Axel Kornerup; Sørensen, Dorte Bratbo

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Jørgensen, B. M. P., Redrobe, P., Brønnum Pedersen, T., Selfjord, E. M., Nielsen, D. S., Plath, N., ... Sørensen, D. B. (2013). *The gut microbiota influence behavior in the subchronic PCP induced animal model of schizophrenia*. Poster session presented at Neurodag 2013, Copenhagen, Denmark.

The Gut Microbiota Influence Behavior in the Subchronic PCP-induced Model of Schizophrenia

Pyndt B.^a, Redrobe J.P.^b, Pedersen T.B.^c, Selfjord E.^a, Nielsen D.S.^d, Plath N.^e, Hansen A.K.^a, Sørensen D.B.^a

^aSection of Experimental Animal Models, Department of Veterinary Disease Biology, The Faculty of Health and Medical Sciences, University of Copenhagen, Denmark,

^bMedical Writing, H. Lundbeck A/S, Valby, Denmark,

^cNonclinical Safety Research, H. Lundbeck A/S, Valby, Denmark.

^dDepartment of Food Science, The Faculty of Science, University of Copenhagen, Denmark

^eSynaptic transmission 1, H. Lundbeck A/S, Valby, Denmark.

Introduction: Evidence is accumulating that the gut microbiota (GM) impact on the individual to a degree, which were not previously believed. We investigated the impact of the GM on behavior in the subchronic phencyclidine (PCP) induced model of Schizophrenia.

Materials and methods: Three batches of 24 male Lister Hooded rats, were injected with either PCP (5 mg/kg) or saline twice a day for seven days, followed by seven days wash out. Hereafter the rats were tested in the novel object recognition test and the locomotor activity test, with a timespan of three and six weeks in between for batch two and three, respectively. Fecal samples were collected at relevant time points. The GM was analyzed by denaturation gradient gel electrophoresis, followed by a principal component analysis, and T-tests or regression analysis.

Results: PCP treatment changed the GM up till three weeks after wash out ($p < 0.05$) and decreased visual cognition in the same period. ($p < 0.05$). Locomotor activity was increased for at least six weeks after wash out. The composition of the GM were strongly correlated to cognitive capabilities in both groups ($p < 0.001$ for both). In both groups the GM also correlated to basal activity of the rats, ($p < 0.05$ for both), but after an acute injection of PCP in the locomotor activity test, the GM only correlated to behavior in the rats which had previously been exposed to PCP ($p < 0.05$).

Conclusion: The study demonstrates an influence of the GM in rats on especially cognitive capabilities, but also on activity. This relationship was evident in the vehicle treated group, but also in the PCP treated group, where the GM composition was changed and correlated to the degree of schizophrenia-like symptoms expressed. The results indicate the relevance of including the variation of the gut microbiota in the analysis of studies using this model.