



## **Arabidopsis thaliana PEN1 (AtSYP121) mediates triggering of innate immunity by bacterial lipo-oligosaccharides (LOS)**

Erbs, Gitte; Dow, John Maxwell; Molinaro, Antonio; Newman, Mari-Anne

*Publication date:*  
2009

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Erbs, G., Dow, J. M., Molinaro, A., & Newman, M-A. (2009). *Arabidopsis thaliana PEN1 (AtSYP121) mediates triggering of innate immunity by bacterial lipo-oligosaccharides (LOS)*. Poster session presented at IS-MPMI 2009 XIV International Congress on Molecular Plant-Microbe Interactions, Quebec, Canada.



XIV  
International Congress

On Molecular  
Plant-Microbe Interactions

July 19-23, 2009, Quebec City, Canada

**IS-MPMI 2009 XIV International Congress  
on Molecular Plant-Microbe Interactions  
Abstracts of Poster Presentations**



# *Arabidopsis thaliana* PEN1 (AtSYP121) mediates triggering of innate immunity by bacterial lipo-oligosaccharides (LOS)






Gitte ERBS<sup>1</sup>, John Maxwell DOW<sup>2</sup>, Antonio MOLINARO<sup>3</sup> and Mari-Anne NEWMAN<sup>1</sup>

<sup>1</sup>Faculty of Life Sciences, Dept of Plant Biology & Biotechnology, University of Copenhagen, Denmark. <sup>2</sup>BIOMERIT Research Centre, Dept. of Microbiology, National University of Ireland, Cork, Ireland. <sup>3</sup>Dipartimento di Chimica Organica e Biochimica, Università di Napoli Federico II, Italy.



## Abstract

In eukaryotes, proteins of the soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) family are believed to be required for docking and fusion of intracellular transport vesicles with acceptor/target membranes. The *Arabidopsis* syntaxin PEN1 (AtSYP121) is a SNARE protein that has been shown to play a role in pathogen resistance in *Arabidopsis* towards fungi (Kwon *et al.*, 2008). We present data to show that *Arabidopsis* PEN1 is also involved in signal transduction leading to the induction of the innate immune responses by particular microbe-associated molecular patterns (MAMPs) of bacterial origin. Specifically we show that PEN1 is required for induction of *PR1* gene induction, callose deposition and generation of reactive oxygen species (ROS) by LOS but not by flagellin.



## Summary

Taken together our results suggest that PEN1 has a role in triggering immune responses in *Arabidopsis* in response to LOS but not in response to flg22. One explanation could be that PEN1 is required for the correct localisation at the plant cell plasma-membrane of the putative receptor(s) for LOS, but not for the flagellin receptor FLS2. Alternatively, PEN1 may be required for endocytosis of an LOS complex, which may allow signalling to cytoplasmically located proteins to trigger defence responses. Although we favour a model in which PEN1 is involved in exocytosis required for *Xcc* LOS triggered immunity in *Arabidopsis*, we cannot discount an alternate role in endocytosis. Importantly our findings indicate that PEN1 may have roles in plant disease resistance that have not been appreciated thus far.