



**MicroRNA 101b Is Downregulated in the Prefrontal Cortex of a Genetic Model of Depression and Targets the Glutamate Transporter SLC1A1 (EAAT3) in Vitro**

Bin Wei, Ya; Melas, Philippe A.; Villaescusa, J. Carlos; Liu, Jia Jia; Xu, Ning; Christiansen, Soren Hofman; Elbrønd-Bek, Heidi; Woldbye, David Paul Drucker; Wegener, Gregers; Mathe, Aleksander A.; Lavebratt, Catharina

*Published in:*  
International Journal of Neuropsychopharmacology

*DOI:*  
[10.1093/ijnp/pyw069](https://doi.org/10.1093/ijnp/pyw069)

*Publication date:*  
2016

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/)

*Citation for published version (APA):*  
Bin Wei, Y., Melas, P. A., Villaescusa, J. C., Liu, J. J., Xu, N., Christiansen, S. H., ... Lavebratt, C. (2016). MicroRNA 101b Is Downregulated in the Prefrontal Cortex of a Genetic Model of Depression and Targets the Glutamate Transporter SLC1A1 (EAAT3) in Vitro. *International Journal of Neuropsychopharmacology*, 19(12), [pyw069]. <https://doi.org/10.1093/ijnp/pyw069>

## REGULAR RESEARCH ARTICLE

# MicroRNA 101b Is Downregulated in the Prefrontal Cortex of a Genetic Model of Depression and Targets the Glutamate Transporter SLC1A1 (EAAT3) *in Vitro*

Ya Bin Wei, PhD; Philippe A. Melas, PhD; J. Carlos Villaescusa, PhD; Jia Jia Liu, PhD; Ning Xu, PhD; Søren Hofman Christiansen, PhD; Heidi Elbrønd-Bek, PhD; David Paul Drucker Woldbye, PhD; Gregers Wegener, MD, PhD; Aleksander A. Mathé, MD, PhD; Catharina Lavebratt, PhD

Department of Molecular Medicine and Surgery, Neurogenetics Unit (Drs Wei, Melas, Liu, and Lavebratt), Center for Molecular Medicine (Drs Wei, Melas, Villaescusa, Liu, Xu, and Lavebratt), Department of Clinical Neuroscience (Drs Melas and Mathé), Department of Molecular Biochemistry and Biophysics, Neurogenetics Unit (Dr Villaescusa), and Department of Medicine (Dr Xu), Karolinska Institutet, Stockholm, Sweden; Department of Neuroscience and Pharmacology, Laboratory for Neural Plasticity, University of Copenhagen, Denmark (Drs Christiansen, Elbrønd-Bek, and Woldbye); Translational Neuropsychiatry Unit, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark (Dr Wegener); Centre of Excellence for Pharmaceutical Sciences, North-West University, Potchefstroom, South Africa (Dr Wegener).

Correspondence: Ya Bin Wei, PhD, Neurogenetics Unit, CMM L8:00, Karolinska University Hospital, 171 76 Stockholm, Sweden ([yabin.wei@ki.se](mailto:yabin.wei@ki.se)).

## Abstract

**Background:** MicroRNAs (miRNAs) are small regulatory molecules that cause translational repression by base pairing with target mRNAs. Cumulative evidence suggests that changes in miRNA expression may in part underlie the pathophysiology and treatment of neuropsychiatric disorders, including major depressive disorder (MDD).

**Methods:** A miRNA expression assay that can simultaneously detect 423 rat miRNAs (miRBase v.17) was used to profile the prefrontal cortex (PFC) of a genetic rat model of MDD (the Flinders Sensitive Line [FSL]) and the controls, the Flinders Resistant Line (FRL). Gene expression data from the PFC of FSL/FRL animals (GEO accession no. GSE20388) were used to guide mRNA target selection. Luciferase reporter assays were used to verify miRNA targets *in vitro*.

**Results:** We identified 23 miRNAs that were downregulated in the PFC of the FSL model compared with controls. Interestingly, one of the identified miRNAs (miR-101b) is highly conserved between rat and human and was recently found to be downregulated in the PFC of depressed suicide subjects. Using a combination of *in silico* and *in vitro* analyses, we found that miR-101b targets the neuronal glutamate transporter SLC1A1 (also known as EAAC1 or EAAT3). Accordingly, both mRNA and protein levels of SLC1A1 were found to be upregulated in the PFC of the FSL model.

**Conclusions:** Besides providing a list of novel miRNAs associated with depression-like states, this preclinical study replicated the human association of miR-101 with depression. In addition, since one of the targets of miR-101b appears to be a glutamate transporter, our preclinical data support the hypothesis of a glutamatergic dysregulation being implicated in the etiology of depression.

**Keywords:** epigenetics, miRNA, depression, EAAC1, DCBXA

Received: February 21, 2016; Revised: July 12, 2016; Accepted: August 2, 2016

© The Author 2016. Published by Oxford University Press on behalf of CINP.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

## Significance Statement

MicroRNAs (miRNAs) are small non-coding RNA molecules that cause messenger RNA silencing. In the present study, we found miRNA changes in an established rodent model of major depressive disorder (MDD). Interestingly, one of the dysregulated miRNA (miR-101) was recently also identified as being affected in postmortem brain samples of human MDD subjects. To further our understanding on miR-101b's function, we used *in vitro* assays and found that it targets the neuronal glutamate transporter SLC1A1 (also known as EAAC1 or EAAT3). These preclinical findings suggest a glutamatergic dysregulation in MDD and support the literature showing efficacy of novel antidepressant compounds that act as glutamate receptor modulators.

## Introduction

MicroRNAs (miRNAs) are a family of small, on average 22 nucleotides long, non-coding RNAs that regulate gene expression at the posttranscriptional level (Winter et al., 2009). Most known mammalian miRNAs are expressed in a tissue-specific manner, cause translational repression, and can individually target hundreds of genes (He and Hannon, 2004). A known mechanism, through which miRNAs exert their repressive function, involves their incorporation into the RNA-induced silencing complex (RISC), which uses the miRNA's seed region (nucleotides 2–8 at the 5' end of the miRNA) as a template for recognizing complementary sites in the 3' untranslated region (UTR) of target mRNAs (Winter et al., 2009). The latter mRNA targeting by miRNAs leads to mRNA degradation, destabilization, or translational inhibition (Winter et al., 2009).

The human genome codes for thousands of miRNAs (Londin et al., 2015), and over 60% of protein-coding genes seem to have been under selective pressure to maintain miRNA pairings (Friedman et al., 2009). Not unexpectedly, miRNAs have been found to play an important role in human diseases, including cancer, cardiovascular disease, and other pathological conditions involving stress responsive pathways (Mendell and Olson, 2012). Within the nervous system, miRNAs are thought to contribute to brain development, neural function, and synaptic plasticity (Forero et al., 2010). Dysregulation of specific miRNAs has been observed in both patients and animal models of neuropsychiatric disorders, including addiction, anxiety, autism, and schizophrenia (Forero et al., 2010; Miller and Wahlestedt, 2010; Moreau et al., 2011; Chan and Kocerha, 2012). A critical component necessary for miRNA biogenesis, Dicer1, has also been implicated in posttraumatic stress and anxiolytic responses (Dias et al., 2014; Wingo et al., 2015).

With regard to major depressive disorder (MDD), the evidence supporting a miRNA involvement in the pathophysiology and the treatment of the disorder is increasing (Dwivedi, 2016). More specifically, genetic polymorphisms in different miRNAs (e.g., miR-30e and miR-182) have been associated with MDD (Saus et al., 2010; Xu et al., 2010). In addition, certain miRNAs (e.g., miR-16, miR-135, miR-335, and miR-1202) have been found to contribute to the therapeutic action of antidepressants, including that of selective serotonin re-uptake inhibitors, tricyclics, and ketamine (Baudry et al., 2010; Launay et al., 2011; O'Connor et al., 2013; Issler et al., 2014; Lopez et al., 2014; Li et al., 2015). Separate studies also found that the expression levels of a number of miRNAs were changed in blood samples (Bocchio-Chiavetto et al., 2013; Fan et al., 2014) and fibroblasts (Garbett et al., 2015) of patients with MDD. Finally, a miRNA expression study that profiled the prefrontal cortex (PFC) of antidepressant-free depressed suicide subjects found 21 downregulated miRNAs, including miR-101 (Smalheiser et al., 2012), and one study showed increased miR-511 levels in basolateral amygdala from depressed subjects (Yang et al., 2014).

In the present study, we used a well-established genetic rat model of MDD (the Flinders Sensitive Line [FSL]) and its controls (the Flinders Resistant Line [FRL]) to provide further insights into a possible miRNA dysregulation in depression-like states. FSL rats exhibit some key characteristics of MDD, including anhedonia, emotional memory impairment, and psychomotor retardation, all of which are reversed by antidepressant treatment (Overstreet et al., 2005; Eriksson et al., 2012). At the molecular level, FSL animals have reductions in levels of serotonin receptors (Eriksson et al., 2012) and other key molecules associated with MDD, including BDNF, NPY, and P11 (Melas et al., 2012a, 2012b, 2013; Wei et al., 2015). The FSL also exhibits a dysfunctional regulation of glutamate transmission and has a reduction of both neuronal and glial glutamate receptors and transporters (Eriksson et al., 2012; Gomez-Galan et al., 2013). This is of relevance, given the accumulating evidence that suggests an aberrant glutamatergic signaling in humans with MDD (Sanacora et al., 2012) in combination with the promising role of novel antidepressant compounds that act as glutamate receptor modulators (Caddy et al., 2015).

## Methods

### Animals

Adult, 3-month-old, male FSL and FRL rats ( $n = 6\text{--}7/\text{group}$ ) were used for this study. The number of animals used for each experiment is denoted in the respective methodological section or figure legend. PFC regions and hippocampi of all animals were dissected according to Glowinski and Iversen (1966) and immediately stored at  $-80^{\circ}\text{C}$  until subsequent experimental analyses. All experiments were approved by the Danish National Committee for Ethics in Animal Experimentation and the Ethical Committee for protection of animals at the Karolinska Institutet.

### RNA Extraction and Reverse Transcription

Total RNA from the PFC and one hippocampus of each FSL/FRL animal were extracted using the miRNA Universal Kit (Qiagen) followed by treatment with DNase I (Qiagen) to digest contaminating DNA. Concentrations were determined using the NanoDrop ND-1000 (NanoDrop Technologies Inc.). For miRNA analyses, complementary DNA (cDNA) was synthesized using the Universal cDNA Synthesis Kit II for RT-PCR (Exiqon) according to the manufacturer's protocol. UniSp6 RNA was spiked in for monitoring conversion efficiency. For mRNA analyses, cDNA was synthesized using the SuperScript III First-Strand Synthesis System for RT-PCR (Invitrogen; Life Technologies) according to the manufacturer's protocol. cDNA and RNA samples were stored at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$ , respectively, until further processing.

## miRNA Profiling

The nCounter Rat miRNA Expression Assay (NanoString Technologies, Inc.), which includes 423 rat miRNAs derived from miRBase (version 17), was used to profile miRNA expression in the PFC of FSL/FRL animals. In brief, 100 ng of total RNA in 5  $\mu$ L ( $n = 6$  FSL,  $n = 6$  FRL) was used for hybridizations of each miRNA assay according to the manufacturer's instructions. The miRNA raw data were normalized for lane-to-lane variation using a dilution series of 6 spike-in positive controls. The normalization factor for each lane was calculated as follows: the sum of the 6 positive controls for a given lane, divided by the average sum across lanes. The normalization factor was then multiplied by the raw counts of each lane to produce a normalized value. The background level of each lane was defined as 2 SD above the signal of the no-target sequence (negative) controls. We included a miRNA probe in the subsequent analyses only if at least 5 of 6 samples, of either FSL or FRL, had levels above the background value.

## Gene Expression Microarray Data and Pathway Analysis

In a previous study, Blaveri et al. (2010) used 2 independent cohorts of male FSL and FRL animals to conduct a genome-wide mRNA expression profiling of the hippocampus and PFC. For the purposes of their study, they used the Affymetrix Rat Genome 230 2.0 GeneChips and deposited the microarray data in the GEO microarray database (accession no. GSE20388). To guide our selection of putative miRNA targets, we retrieved the microarray PFC data sets and reprocessed them using a mixed model ANOVA according to the initial publication (Blaveri et al., 2010). Since we found a main downregulation of miRNAs in the PFC of FSL animals, we further filtered the microarray probe-set to include only the mRNAs that showed upregulation in the FSL in both analyzed cohorts with a combined  $P$  value of  $< .05$ . The filtered probe-sets were then imported into the Ingenuity Pathway Analysis (IPA; Qiagen) for miRNA target predictions and also for analyzing pathways, diseases, and biofunctions, including molecular and cellular functions and physiological system development and functions.

## Coexpression Analysis

A miRNA coexpression analysis was performed based on the method in Smalheiser et al. (2011) with a minor modification. MiRNA pairs were filtered based on 3 criteria: (1) they showed significant pairwise expression level correlations ( $r > 0.8$ ) across the individual rats in the FSL group; (2) the corresponding pairwise correlations in the FRL group were significantly less than those in the FSL group ( $r_{\text{FSL}} - r_{\text{FRL}} > 0.8$ ); and (3) the miRNA pairs were not significantly negatively correlated in the FRL group ( $r_{\text{FRL}} > -0.8$ ).

## Validation of miRNAs and mRNA Expression Levels

Validation experiments of miRNA hits and measurements of putative mRNA targets, in the PFC of FSL/FRL animals, were performed using quantitative real-time PCR (qRT-PCR). All qRT-PCR amplifications, as well as that in hippocampal cDNA, were performed in triplicates using Power SYBR Green (Applied Biosystems) on an ABI PRISM 7900 HT Sequence Detection System (Applied Biosystems), with the following conditions: 95°C for 10 minutes, followed by 40 repeats of 95°C for 15 seconds (10 seconds for miRNA amplification), 60°C for 1 minute, and a final dissociation stage to monitor

amplification specificity. For miRNA expression analyses, LNA PCR primers (Exiqon) were used to amplify miRNAs of interest, and the data were normalized to *Rnu5g*. For mRNA expression analyses, conventional primers were used and data were normalized to 2 reference genes (*Mff*, mitochondrial fission factor; and *Cdip1*, CDP-diacylglycerol--inositol 3-phosphatidyltransferase). Relative gene expression quantifications were calculated using the qBase software (version 1.3.4) (Hellemans et al., 2007). The tested genes and corresponding primer pair sequences are listed in [supplementary Table 1](#).

## Protein Expression Levels

Protein expression levels of SLC1A1 (also known as EAAC1, EAAT3, or DCBXA), in the PFC of FSL/FRL animals, were quantified using Western blotting as previously described (Wei et al., 2015). Primary antibody incubations were performed overnight at 4°C with a mouse monoclonal anti-EAAT3 antibody (1:1000 dilution, ab78395; Abcam) and with a mouse monoclonal anti- $\beta$ -actin antibody (1:10,000 dilution, A5316; Sigma-Aldrich) as loading control. Membranes were exposed to the Amersham ECL Plus Western Blotting Detection reagent (GE Healthcare), and immunoreactive bands were detected using the Amersham Hyperfilm ECL (GE Healthcare). Optical densities were quantified using the NIH ImageJ software (1.47 version). SLC1A1 protein expression levels were normalized to the expression levels of  $\beta$ -actin and the data were presented as relative quantifications.

## In Vitro Verification of miRNA Targets

To examine whether SLC1A1 is a direct target of miR-101b, we used LightSwitch 3'UTR luciferase reporter constructs (SwitchGear Genomics). HEK 293 cells were co-transfected with 100 ng of a human SLC1A1 GoClone 3'UTR reporter (S810654, SwitchGear Genomics) and either 50 nM of a miR-101b mimic or 50 nM of a miRNA non-targeting control (Switchgear Genomics) using DharmaFECT Duo transfection reagent (GE Dharmacon). An empty vector (no 3'UTR; SwitchGear Genomics) was used as a control. Luciferase activity was measured on a plate luminometer 24 hours after transfection by adding 100  $\mu$ L of LightSwitch Assay Reagent (SwitchGear Genomics). In total, the experiment was replicated 3 times with at least 3 biological replicates each time.

## Statistical Analyses

Data in the bar graphs are presented as mean values  $\pm 1$  SEM. The normal distribution of the data and the homogeneity of the variances were tested using the Shapiro-Wilk and Levene's tests, respectively. The miRNA expression levels of FSL vs FRL, generated by the NanoString nCounter Rat miRNA assay, were compared using the Mann-Whitney U-test. The difference of miRNA expression levels, generated by RT-PCR, between FSL and FRL was assessed using the 2-tailed Student's  $t$  test if the variances in FRL and FSL were equal or was otherwise assessed using the Mann-Whitney U test. The difference in miRNA expression variation between FSL vs. FRL was assessed by comparing the coefficients of variance of individual miRNAs using Mann-Whitney U-test. The threshold for statistical significance was set at  $P < .05$ . All analyses were performed using IBM SPSS Statistics version 22 (IBM Corporation). Extreme values are defined as potential outliers using box-plot function from SPSS software that are denoted in the figure legends and were excluded from the analyses.


## Results

### Downregulation of miRNAs in the PFC of the FSL Animals

To detect putative changes in miRNA expression in the PFC of FSL (vs control FRL) animals, we used a miRNA expression assay that enables multiplex detection of >400 rat miRNAs. After data normalization, only 121 miRNAs were above the background level in both rat strains, in agreement with the fact that most miRNAs are expressed in a tissue-specific manner (He and Hannon, 2004). We found a global miRNA expression decrease in the FSL animals that is characterized by depression-like behavior (supplementary Table 2). By plotting the relationship between the standard deviations and the mean expression values across all expressed miRNAs, we also observed that the miRNA expression in the FSL rats was significantly more variable compared with control FRL ( $P < .001$ ) (Figure 1). Using individual tests of statistical significance, we identified 23 miRNAs with lower (at least 30%) expression levels in the FSL rats ( $P < .05$ , Benjamini-Hochberg False Discovery Rate  $< 0.22$ ; for the full list of miRNAs, see supplementary Table 2). To validate these observations, we selected 9 miRNAs that showed differential expression and performed individual qRT-PCR experiments. The latter analyses confirmed that all selected miRNAs had decreased expression levels in the FSL, with 8 of the 9 miRNAs reaching statistical significance (Figure 2). We also performed a coexpression analysis and found that in total, 244 miRNA pairs had correlated expression levels only in the FSL group. Of those, 51 miRNAs were significantly coexpressed with at least 3 other miRNAs in this set (supplementary Table 3), including 7 of the 23 miRNAs that were significantly decreased in the FSL. Several of these 51 miRNAs belong to the let-7 miRNA family (let-7a, 7b, 7c, 7d, 7f, and 7i). Some miRNAs (miR-125a-3p and miR-125a-5p, miR-125b-3p and miR-125b-5p, mo-miR-3563-3p and miR-3563-5p) derive from the same precursor miRNA.

### Pathway Analyses and miRNA Target Predictions

Since miRNAs have been shown to correlate negatively with levels of target mRNAs (Winter et al., 2009), the miRNA



**Figure 1.** Plot of  $\log_{10}$  Mean vs  $\log_{10}$  Standard deviation based on the microRNA (miRNA) assay expression levels in the prefrontal cortex (PFC) of the Flinders Sensitive Line (FSL) vs control-Flinders Resistant Line (FRL) animals. MiRNA expression in the FSL depression model showed significantly more variation compared with controls. Polynomial (order = 2) fitting line was added to each rat strain, with the FSL represented by the solid line and the control-FRL by the dashed line.


downregulation observed in the FSL model may lead to an upregulation of their mRNA targets. Among the 23 differentially expressed miRNAs, 16 had information about targets in IPA. In a previous study, Blaveri et al. (2010) used PFC regions of the FSL/FRL model to conduct a genome-wide mRNA profiling study. After retrieving the latter microarray dataset, we reanalyzed the data by focusing on the probe sets that showed a significant upregulation ( $P < .05$ ) in the FSL. Next, we tested for an overlap between the latter mRNA candidates and the targets of the 16 differentially expressed miRNAs using IPA. The analysis revealed 1392 potential mRNA targets that were either predicted *in silico* or had been experimentally observed (supplementary Table 4). The core analysis in IPA suggested that the target genes are involved in canonical pathways involved in protein ubiquitination, Ephrin receptor signaling, and Dopamine-DARPP32 signaling (Table 1). These genes were also associated with neurological diseases, psychiatric and psychological disorders, and monogenic disorders (Table 1).

### Gene Expression Measurements of miR-101b Targets

The human homolog of one of the miRNAs that was confirmed to be downregulated in the FSL, miR-101b, was found to be downregulated in the PFC of depressed suicide subjects (Smalheiser et al., 2012). Therefore, miR-101b was considered to warrant further examination. First, we tested if miR-101b showed different expression levels in the hippocampus, another brain region implicated in MDD, of the FSL/FRL model. We found that hippocampal miR-101b expression levels were not different between FSL and FRL ( $P = .86$ ) (supplemental Figure 1). Further, there is high conservation between the mature rat miR-101b and the mature human miR-101, which share the same seed sequence (Figure 3). A total of 248 genes were predicted to be miR-101b targets (supplementary Table 5), among which 103 genes were also part of the IPA pathway genes. From this list, we identified 8 genes that showed high conservation between human and rat in their 3'UTR and were also implicated in depression-related pathways in IPA (including glutamate, GR or GABA receptor signaling, and NFR2-mediated oxidative stress response). The mRNA levels of these 8 genes (*Slc1a1*, *Gabrb2*, *Adcy5*, *Gclc*, *Pbrm1*, *Prkaa1*, *Rac1*, and *Smarcdca4*) were therefore measured in the PFC of the FSL/FRL animals. We found that 2 genes, *Slc1a1* and *Rac1*, showed significantly increased mRNA levels in the FSL (Figure 4a), in line with the decreased levels of miR-101b.

### MiR-101b Targets SLC1A1 *In Vitro* and Correlates Negatively with SLC1A1 Protein Levels *In Vivo*

Next, we chose to focus on SLC1A1 (also known as EAAC1 or EAAT3), since it is a neuronal glutamate transporter that is highly enriched in the cortex (Rothstein et al., 1994). In addition, the FSL model is known to exhibit a dysfunctional regulation of glutamate transmission (Gomez-Galan et al., 2013). It was therefore of interest to verify that the elevated mRNA levels of *Slc1a1* in the FSL rats were due to a specific miR-101b targeting of the *Slc1a1* 3'UTR. For this purpose, HEK 293 cells were transfected with a luciferase reporter construct containing the human SLC1A1 3'UTR, which contains the same sequence used for miRNA seed complementarity as the rat *Slc1a1* 3'UTR (Figure 3). There was a significant reduction in luciferase activity when the SLC1A1-3'UTR luciferase construct was cotransfected with the miR-101b mimic compared with the SLC1A1-3'UTR luciferase construct alone ( $P = .004$ ) (Figure 4b) or when the SLC1A1-3'UTR luciferase construct was cotransfected with a nontargeting



**Figure 2.** The microRNA (miRNA) assay identified 23 miRNAs with lower expression levels in the prefrontal cortex (PFC) of the Flinders Sensitive Line (FSL) depression model (supplementary Table 2), and we selected 9 of them for individual validation experiments. Eight of the 9 miRNAs reached statistical significance (miR-10a,  $P = .046$ ; miR-101b,  $P = .011$ ; miR-107,  $P = .02$ ; miR-124,  $P = .03$ ; miR-125a,  $P = .048$ ; miR-125b,  $P = .2$ ; miR-133a,  $P = .03$ ; miR-181a,  $P = .032$ , and miR-199a,  $P = .029$ ). Data are presented as relative quantifications (R.Q.)  $\pm$  SEM. Levels of *Rnu5g* were used as reference for normalization of mature miRNA expression levels ( $n = 6-7$  animals/group;  $n = 1$  outlier/group), \* $P < .05$ .

**Table 1.** Core Analysis Result from IPA

Top Pathways, Diseases, and Biofunction	P-Value/P-Value Range
<i>Top Canonical Pathways</i>	
Protein ubiquitination pathway	4.48E-13
Ephrin receptor signaling	8.56E-07
Dopamine-DARPP32 feedback in cAMP signaling	1.64E-06
Ephrin B signaling	3.26E-06
Axonal guidance signaling	3.42E-06
<i>Top Diseases</i>	
Neurological disease	3.19E-03 - 1.41E-16
Psychological disorders	2.72E-03 - 1.41E-16
Skeletal and muscular disorders	2.72E-03 - 1.99E-16
Hereditary disorder	2.97E-03 - 5.43E-16
<i>Molecular and Cellular Functions</i>	
Cellular assembly and organization	3.26E-03 - 6.36E-24
Cellular function and maintenance	3.26E-03 - 6.36E-24
Molecular transport	2.72E-03 - 8.20E-18
Protein trafficking	5.89E-05 - 1.11E-16
Posttranslational modification	3.31E-03 - 7.54E-16
<i>Physiological System Development and Function</i>	
Organismal survival	9.34E-04 - 2.52E-14
Nervous system development and function	3.22E-03 - 1.78E-13
Tissue development	3.28E-03 - 1.78E-13
Behavior	3.28E-03 - 4.57E-08
<i>Top Networks</i>	
Embryonic development, organismal survival, cell death and survival	
neurological disease, psychological disorders, organismal injury and abnormalities	
free radical scavenging, small molecule biochemistry, hereditary disorder	
cellular assembly and organization, cellular movement, nervous system development and function	


Abbreviation: IPA, Ingenuity Pathway Analysis.

control ( $P = .034$ ) (Figure 4b). The latter results supported the notion of miR-101b directly targeting *Slc1a1*, in line with the miR-101b downregulation and the elevated *Slc1a1* mRNA levels observed in the PFC of the FSL. Finally, since mRNA levels do not always correlate with protein levels (Vogel and Marcotte, 2012), we verified that *SLC1A1* protein levels were also upregulated in the PFC of the FSL rats compared with the FRL ( $P = .004$ ) (Figure 4c).


## Discussion

There is evidence supporting the miRNA involvement in the pathophysiology and the treatment of MDD (Dwivedi, 2016). In this study, we used the FSL rat model of depression to provide further insights into a possible miRNA dysregulation in depression-like states. By using a miRNA expression-profiling assay, which covers >400 known miRNAs, we first observed that the overall miRNA expression in the PFC of the FSL model was downregulated. This finding coincides with a recent human study that showed a 17% global downregulation of miRNAs in the PFC of depressed suicide subjects (Smalheiser et al., 2012). However, in another rat model of depression, both up- and downregulation of miRNA expression was observed, suggesting an exposure protocol and strain-specific miRNA expression difference (Dwivedi et al., 2015). In contrast, an increase in global miRNA expression was observed in postmortem cortical regions of subjects with schizophrenia, which was found to correlate with an increase in the RNase III Droscha that is involved in the processing of miRNAs (Beveridge et al., 2010). Droscha has been shown to account for widespread downregulation of miRNAs in cancer (Thomson et al., 2006), and our data (Wei et al., 2016) show that the global downregulation of miRNAs observed in the FSL model is also associated with a basal reduction in Droscha levels.

Using individual tests of statistical significance, we identified 23 specific miRNAs that were decreased in the PFC of the FSL model compared with controls. In general, miRNAs are known to exert a translational repressive function that involves their



**Figure 3.** The sequences of the mature miR-101 and the 3' untranslated region (UTR) of the *SLC1A1* gene in human, rat, and mouse is shown. The mature miR-101 sequence is highly conserved between species, with only one nucleotide difference between human and rat/mouse (the nucleotide difference is underlined in the human sequence). The target site in the 3'-UTR of the *SLC1A1* gene was predicted by TargetScan and shows that both the miR-101 seed region (5'-UACAGUAC-3') and the mRNA target site are conserved between human and rat/mouse.



**Figure 4.** (a) From a list of >200 genes that were predicted to be miR-101b targets (supplementary Table 5), we identified 8 genes that showed high conservation between human and rat in their 3' untranslated region (UTR) and were also implicated in depression-related pathways. The mRNA expression of these 8 genes was quantified in the prefrontal cortex (PFC) of the Flinders Sensitive Line (FSL) and control-Flinders Resistant Line (FRL) animals. Only *Slc1a1* and *Rac1* mRNA levels were significantly increased in the FSL compared with controls (*Slc1a1*:  $P = .001$ ; *Rac1*:  $P = .023$ ;  $P$ -values of the other 6 genes >.2). (b) Luciferase activity was measured in HEK293 cells transfected with either an empty vector (first bar; positive control), an *SLC1A1*-3'UTR luciferase construct (second bar), an *SLC1A1*-3'UTR luciferase construct cotransfected with a nontargeting control (third bar; negative control), or an *SLC1A1*-3'UTR luciferase construct cotransfected with a miR-101b mimic (fourth bar). MiR-101b significantly inhibited the luciferase activity of the reporter construct containing the 3'UTR (fourth bar) compared with both *SLC1A1* alone (second bar) and *SLC1A1*-3'UTR cotransfected with a nontargeting control (third bar). (c) In line with the increased levels of *Slc1a1* mRNA, the FSL also showed increased protein levels of *SLC1A1*. Representative immunoblotting images of *SLC1A1* and  $\beta$ -actin (loading control) are shown below the graph. Data are presented as relative quantifications (R.Q.)  $\pm$  SEM. Two reference genes (*Cd1pt* and *Mff*) were used for normalization in 4a ( $n = 5-7$  animals/group). \* $P < .05$ , \*\* $P < .01$ .

incorporation into the RISC, which uses the miRNA's seed region as a template for recognizing complementary sites in the 3' UTR of target mRNAs (Winter et al., 2009). When we tested for predicted mRNA targets of the significant miRNAs and analyzed the data using pathway analysis software, we found that the putative target genes are involved in protein ubiquitination, Ephrin receptor signaling, and Dopamine-DARPP32 signaling. Protein ubiquitination is known to play an important role in neuronal function (Tai and Schuman, 2008), Ephrin ligands and their receptors regulate synapse formation and neuronal plasticity (Klein, 2009), and Dopamine-DARPP32 is extensively involved in intracellular signaling pathways and integrating dopaminergic and glutamatergic signaling (Girault and Greengard, 2004). Accordingly, the same genes were associated with both neurological and psychiatric disorders, for example MDD, in the pathway analysis. However, besides the overall analysis of the miRNA list, one specific miRNA (miR-101) was of particular interest, since it was found to be downregulated in the PFC of depressed suicide subjects (Smalheiser et al., 2012). Using a combination of *in silico* and *in vitro* analyses, we found that miR-101 targets the 3'UTR of the neuronal glutamate transporter SLC1A1 (also known as EAAC1 or EAAT3). In agreement with this result, both mRNA and protein levels of SLC1A1 were upregulated in the PFC of the FSL model. Whereas decreased levels of SLC1A1 have been previously reported in stress-induced offspring depression (Zhang et al., 2013), this is to our knowledge the first report to show elevated SLC1A1 levels in a depression-like state. Several approaches have been successfully reported to manipulate specific miRNA expression levels that provide possibilities for future *in vivo* examination of miR-101b function in the FSL/FRL model (Issler and Chen, 2015; Li et al., 2016).

In addition to having an important role in transporting glutamate across neuronal membranes, SLC1A1 is also essential for rapidly binding released glutamate to shape synaptic transmission (Tzingounis and Wadiche, 2007). SLC1A1 has also been found to interact with NMDA receptor subunits NR1, NR2A, and NR2B (Waxman et al., 2007). Interestingly, in the FSL model, NR2A and NR2B subunits are decreased in both the PFC and the hippocampus (Eriksson et al., 2012), which may suggest a novel coregulatory mechanism. Whereas the FSL animals display reductions in serotonin receptors (Eriksson et al., 2012), an independent group recently also confirmed an aberrant glutamatergic transmission in this model of depression (Gomez-Galan et al., 2013). In the latter study, the authors examined the hippocampal region of the FSL and found a dysfunctional astrocytic glutamate regulation that was accompanied by a downregulation of the glial glutamate transporter GLAST (also known as EAAT1). Obviously, our finding of increased SLC1A1 (EAAT3) in the PFC may represent just one of many depression-related glutamatergic and/or serotonergic genes that are being targeted by our validated miRNAs. This is supported by the fact that besides SLC1A1, one other gene that was confirmed to be upregulated in the PFC of the FSL was *Rac1*, which can be activated by serotonin through 5-HT<sub>2A</sub> receptor stimulation (Dai et al., 2008) and is a confirmed miR-101 target (Lin et al., 2014). RAC1 is a small Rho family GTPase that acts as a key regulator of actin cytoskeletal dynamics. It plays an important role in dendritic development and spinogenesis. Dysregulation of RAC1 has been associated with cognitive impairment, memory disturbance, and neurodevelopmental disorders (Govek et al., 2005; Martinez and Tejada-Simon, 2011; Stankiewicz and Linseman, 2014), which may support a role of RAC1 in the pathophysiological process of depression. Taken together, these preclinical data support the shift from a solely monoamine-centered hypothesis

in the pathophysiology of depression to one that includes the involvement of the glutamatergic system (Sanacora et al., 2012).

The traditional approach to identify compounds targeting single molecules (usually receptors or their ligands) has been a hinder for the development of effective psychiatric drugs, and an approach targeting the defective components of an entire biological pathway is likely more efficient. Therefore, compared with single-target drugs, miRNAs possess an advantage that is attributed to their pleiotropic nature, which allows them to individually regulate hundreds of genes.

## Supplementary Material

For supplementary material accompanying this paper, visit <http://www.ijnp.oxfordjournals.org/>

## Acknowledgments

We thank Dr. Andor Pivarcsi for helping with the experimental design and granting agencies and funding sources reported in the previous section for supporting and making this work possible. These funding sources had no involvement in the study design, analyses, and interpretation of the data, or in the writing of the manuscript and the decision to submit the report for publication.

This work was supported by the Karolinska Institutet's Faculty Funds, the Swedish Research Council (grant nos. 2010–3631 CL, 10414 AAM), the Fredrik and Ingrid Thuring's Foundation, the regional agreement on medical training and clinical research (ALF) between the Stockholm County Council and Karolinska Institutet, the Danish Medical Research Council (grant 11-107897), and the Lundbeck foundation (grant R118-A11400), and Aarhus University Research Foundation (eMOOD initiative).

## Statement of Interest

None.

## References

- Baudry A, Mouillet-Richard S, Schneider B, Launay JM, Kellermann O (2010) miR-16 targets the serotonin transporter: a new facet for adaptive responses to antidepressants. *Science* 329:1537–1541.
- Beveridge NJ, Gardiner E, Carroll AP, Tooney PA, Cairns MJ (2010) Schizophrenia is associated with an increase in cortical microRNA biogenesis. *Mol Psychiatry* 15:1176–1189.
- Blaveri E, Kelly F, Mallei A, Harris K, Taylor A, Reid J, Razzoli M, Carboni L, Piubelli C, Musazzi L, Racagni G, Mathé A, Popoli M, Domenici E, Bates S (2010) Expression profiling of a genetic animal model of depression reveals novel molecular pathways underlying depressive-like behaviours. *PLoS One* 5:e12596.
- Bocchio-Chiavetto L, Maffioletti E, Bettinsoli P, Giovannini C, Bignotti S, Tardito D, Corrada D, Milanese L, Gennarelli M (2013) Blood microRNA changes in depressed patients during antidepressant treatment. *Eur Neuropsychopharmacol* 23:602–611.
- Caddy C, Amit BH, McCloud TL, Rendell JM, Furukawa TA, McShane R, Hawton K, Cipriani A (2015) Ketamine and other glutamate receptor modulators for depression in adults. *Cochrane Database Syst Rev* 9:CD011612.
- Chan AW, Kocerha J (2012) The Path to microRNA Therapeutics in Psychiatric and Neurodegenerative Disorders. *Front Genet* 3:82.

- Dai Y, Dudek NL, Patel TB, Muma NA (2008) Transglutaminase-catalyzed transamidation: a novel mechanism for Rac1 activation by 5-hydroxytryptamine<sub>2A</sub> receptor stimulation. *J Pharmacol Exp Ther* 326:153–162.
- Dias C et al. (2014) beta-catenin mediates stress resilience through Dicer1/microRNA regulation. *Nature* 516:51–55.
- Dwivedi Y (2016) Pathogenetic and therapeutic applications of microRNAs in major depressive disorder. *Progress in neuro-psychopharmacology & biological psychiatry* 64:341–348.
- Dwivedi Y, Roy B, Lugli G, Rizavi H, Zhang H, Smalheiser NR (2015) Chronic corticosterone-mediated dysregulation of microRNA network in prefrontal cortex of rats: relevance to depression pathophysiology. *Translational psychiatry* 5:e682.
- Eriksson TM, Delagrèze P, Spedding M, Popoli M, Mathé AA, Ogren SO, Svenningsson P (2012) Emotional memory impairments in a genetic rat model of depression: involvement of 5-HT/MEK/Arc signaling in restoration. *Mol Psychiatry* 17:173–184.
- Fan HM, Sun XY, Guo W, Zhong AF, Niu W, Zhao L, Dai YH, Guo ZM, Zhang LY, Lu J (2014) Differential expression of microRNA in peripheral blood mononuclear cells as specific biomarker for major depressive disorder patients. *J Psychiatr Res* 59:45–52.
- Forero DA, van der Ven K, Callaerts P, Del-Favero J (2010) miRNA genes and the brain: implications for psychiatric disorders. *Hum Mutat* 31:1195–1204.
- Friedman RC, Farh KK, Burge CB, Bartel DP (2009) Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 19:92–105.
- Garbett KA, Vereczkei A, Kalman S, Brown JA, Taylor WD, Faludi G, Korade Z, Shelton RC, Mirmics K (2015) Coordinated messenger RNA/microRNA changes in fibroblasts of patients with major depression. *Biol Psychiatry* 77:256–265.
- Girault JA, Greengard P (2004) The neurobiology of dopamine signaling. *Arch Neurol* 61:641–644.
- Glowinski J, Iversen LL (1966) Regional studies of catecholamines in the rat brain. I. The disposition of [3H]norepinephrine, [3H]dopamine and [3H]dopa in various regions of the brain. *J Neurochem* 13:655–669.
- Gomez-Galan M, De Bundel D, Van Eeckhaut A, Smolders I, Lindskog M (2013) Dysfunctional astrocytic regulation of glutamate transmission in a rat model of depression. *Mol Psychiatry* 18:582–594.
- Govek EE, Newey SE, Van Aelst L (2005) The role of the Rho GTPases in neuronal development. *Genes & development* 19:1–49.
- He L, Hannon GJ (2004) MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 5:522–531.
- Hellems J, Mortier G, De Paepe A, Speleman F, Vandesompele J (2007) qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biol* 8:R19.
- Issler O, Chen A (2015) Determining the role of microRNAs in psychiatric disorders. *Nature reviews Neuroscience* 16:201–212.
- Issler O, Haramati S, Paul ED, Maeno H, Navon I, Zwang R, Gil S, Mayberg HS, Dunlop BW, Menke A, Awatramani R, Binder EB, Deneris ES, Lowry CA, Chen A (2014) MicroRNA 135 is essential for chronic stress resiliency, antidepressant efficacy, and intact serotonergic activity. *Neuron* 83:344–360.
- Klein R (2009) Bidirectional modulation of synaptic functions by Eph/ephrin signaling. *Nat Neurosci* 12:15–20.
- Launay JM, Mouillet-Richard S, Baudry A, Pietri M, Kellermann O (2011) Raphe-mediated signals control the hippocampal response to SRI antidepressants via miR-16. *Translational psychiatry* 1:e56.
- Li J, Meng H, Cao W, Qiu T (2015) MiR-335 is involved in major depression disorder and antidepressant treatment through targeting GRM4. *Neuroscience letters* 606:167–172.
- Li Y, Li S, Yan J, Wang D, Yin R, Zhao L, Zhu Y, Zhu X (2016) miR-182 (microRNA-182) suppression in the hippocampus evokes antidepressant-like effects in rats. *Progress in neuro-psychopharmacology & biological psychiatry* 65:96–103.
- Lin X, Guan H, Li H, Liu L, Liu J, Wei G, Huang Z, Liao Z, Li Y (2014) miR-101 inhibits cell proliferation by targeting Rac1 in papillary thyroid carcinoma. *Biomed Rep* 2:122–126.
- Londin E, Loher P, Telonis AG, Quann K, Clark P, Jing Y, Hatzimichael E, Kirino Y, Honda S, Lally M, Ramratnam B, Comstock CE, Knudsen KE, Gomella L, Spaeth GL, Hark L, Katz LJ, Witkiewicz A, Rostami A, Jimenez SA, et al. (2015) Analysis of 13 cell types reveals evidence for the expression of numerous novel primate- and tissue-specific microRNAs. *Proc Natl Acad Sci U S A* 112:E1106–1115.
- Lopez JP, Lim R, Cruceanu C, Crapper L, Fasano C, Labonte B, Mausion G, Yang JP, Yerko V, Vigneault E, El Mestikawy S, Mechawar N, Pavlidis P, Turecki G (2014) miR-1202 is a primate-specific and brain-enriched microRNA involved in major depression and antidepressant treatment. *Nat Med* 20:764–768.
- Major Depressive Disorder Working Group of the Psychiatric GC et al. (2013) A mega-analysis of genome-wide association studies for major depressive disorder. *Mol Psychiatry* 18:497–511.
- Martinez LA, Tejada-Simon MV (2011) Pharmacological inactivation of the small GTPase Rac1 impairs long-term plasticity in the mouse hippocampus. *Neuropharmacology* 61:305–312.
- Melas PA, Mannervik M, Mathé AA, Lavebratt C (2012a) Neuropeptide Y: identification of a novel rat mRNA splice-variant that is downregulated in the hippocampus and the prefrontal cortex of a depression-like model. *Peptides* 35:49–55.
- Melas PA, Rogdaki M, Lennartsson A, Bjork K, Qi H, Witas P, Werme M, Wegener G, Mathé AA, Svenningsson P, Lavebratt C (2012b) Antidepressant treatment is associated with epigenetic alterations in the promoter of P11 in a genetic model of depression. *The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum* 15:669–679.
- Melas PA, Lennartsson A, Vakifahmetoglu-Norberg H, Wei Y, Aberg E, Werme M, Rogdaki M, Mannervik M, Wegener G, Brené S, Mathé AA, Lavebratt C (2013) Allele-specific programming of Npy and epigenetic effects of physical activity in a genetic model of depression. *Translational psychiatry* 3:e255.
- Mendell JT, Olson EN (2012) MicroRNAs in stress signaling and human disease. *Cell* 148:1172–1187.
- Miller BH, Wahlestedt C (2010) MicroRNA dysregulation in psychiatric disease. *Brain Res* 1338:89–99.
- Moreau MP, Bruse SE, David-Rus R, Buyske S, Brzustowicz LM (2011) Altered microRNA expression profiles in postmortem brain samples from individuals with schizophrenia and bipolar disorder. *Biol Psychiatry* 69:188–193.
- O'Connor RM, Grenham S, Dinan TG, Cryan JF (2013) microRNAs as novel antidepressant targets: converging effects of ketamine and electroconvulsive shock therapy in the rat hippocampus. *The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum* 16:1885–1892.
- Overstreet DH, Friedman E, Mathé AA, Yalid G (2005) The Flinders Sensitive Line rat: a selectively bred putative animal model of depression. *Neurosci Biobehav Rev* 29:739–759.
- Rothstein JD, Martin L, Levey AI, Dykes-Hoberg M, Jin L, Wu D, Nash N, Kuncl RW (1994) Localization of neuronal and glial glutamate transporters. *Neuron* 13:713–725.

- Sanacora G, Treccani G, Popoli M (2012) Towards a glutamate hypothesis of depression: an emerging frontier of neuropsychopharmacology for mood disorders. *Neuropharmacology* 62:63–77.
- Saus E, Soria V, Escaramis G, Vivarelli F, Crespo JM, Kagerbauer B, Menchon JM, Urretavizcaya M, Gratacos M, Estivill X (2010) Genetic variants and abnormal processing of pre-miR-182, a circadian clock modulator, in major depression patients with late insomnia. *Hum Mol Genet* 19:4017–4025.
- Smalheiser NR, Lugli G, Rizavi HS, Zhang H, Torvik VI, Pandey GN, Davis JM, Dwivedi Y (2011) MicroRNA expression in rat brain exposed to repeated inescapable shock: differential alterations in learned helplessness vs. non-learned helplessness. *The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum* 14:1315–1325.
- Smalheiser NR, Lugli G, Rizavi HS, Torvik VI, Turecki G, Dwivedi Y (2012) MicroRNA expression is down-regulated and reorganized in prefrontal cortex of depressed suicide subjects. *PLoS One* 7:e33201.
- Stankiewicz TR, Linseman DA (2014) Rho family GTPases: key players in neuronal development, neuronal survival, and neurodegeneration. *Frontiers in cellular neuroscience* 8:314.
- Tai HC, Schuman EM (2008) Ubiquitin, the proteasome and protein degradation in neuronal function and dysfunction. *Nature reviews Neuroscience* 9:826–838.
- Thomson JM, Newman M, Parker JS, Morin-Kensicki EM, Wright T, Hammond SM (2006) Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes & development* 20:2202–2207.
- Tzingounis AV, Wadiche JI (2007) Glutamate transporters: confining runaway excitation by shaping synaptic transmission. *Nature reviews Neuroscience* 8:935–947.
- Vogel C, Marcotte EM (2012) Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nat Rev Genet* 13:227–232.
- Waxman EA, Bacongus I, Lynch DR, Robinson MB (2007) N-methyl-D-aspartate receptor-dependent regulation of the glutamate transporter excitatory amino acid carrier 1. *J Biol Chem* 282:17594–17607.
- Wei YB, Liu JJ, Villaescusa JC, Åberg E, Brené S, Wegener G, Mathé AA, Lavebratt C (2016) Elevation of IL6 is associated with disturbed let-7 biogenesis in a genetic model of depression. *Transl Psychiatry* 6:e869.
- Wei Y, Melas PA, Wegener G, Mathé AA, Lavebratt C (2015) Anti-depressant-like effect of sodium butyrate is associated with an increase in TET1 and in 5-hydroxymethylation levels in the *Bdnf* gene. *The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum* 18.
- Wingo AP, Almlı LM, Stevens JJ, Klengel T, Uddin M, Li Y, Bus-tamante AC, Lori A, Koen N, Stein DJ, Smith AK, Aiello AE, Koenen KC, Wildman DE, Galea S, Bradley B, Binder EB, Jin P, Gibson G, Ressler KJ (2015) DICER1 and microRNA regulation in post-traumatic stress disorder with comorbid depression. *Nat Commun* 6:10106.
- Winter J, Jung S, Keller S, Gregory RI, Diederichs S (2009) Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat Cell Biol* 11:228–234.
- Xu Y, Liu H, Li F, Sun N, Ren Y, Liu Z, Cao X, Wang Y, Liu P, Zhang K (2010) A polymorphism in the microRNA-30e precursor associated with major depressive disorder risk and P300 waveform. *J Affect Disord* 127:332–336.
- Yang X, Yang Q, Wang X, Luo C, Wan Y, Li J, Liu K, Zhou M, Zhang C (2014) MicroRNA expression profile and functional analysis reveal that miR-206 is a critical novel gene for the expression of BDNF induced by ketamine. *Neuromolecular medicine* 16:594–605.
- Zhang XH, Jia N, Zhao XY, Tang GK, Guan LX, Wang D, Sun HL, Li H, Zhu ZL (2013) Involvement of pGluR1, EAAT2 and EAAT3 in offspring depression induced by prenatal stress. *Neuroscience* 250:333–341.