



Københavns Universitet

## EDC IMPACT

Rehfeld, Anders; Egeberg, Dorte; Almstrup, Kristian; Holm Petersen, Jørgen; Dissing, Steen; Skakkebæk, Niels Erik

*Published in:*  
Endocrine Connections

*DOI:*  
[10.1530/EC-17-0156](https://doi.org/10.1530/EC-17-0156)

*Publication date:*  
2018

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY-NC-ND](#)

*Citation for published version (APA):*  
Rehfeld, A., Egeberg, D., Almstrup, K., Holm Petersen, J., Dissing, S., & Skakkebæk, N. E. (2018). EDC IMPACT: Chemical UV filters can affect human sperm function in a progesterone-like manner. *Endocrine Connections*, 7(1), 16-25. <https://doi.org/10.1530/EC-17-0156>

## RESEARCH

# EDC IMPACT: Chemical UV filters can affect human sperm function in a progesterone-like manner

A Rehfeld<sup>1,2,3</sup>, D L Egeberg<sup>1,3</sup>, K Almstrup<sup>1,3</sup>, J H Petersen<sup>1,3,4</sup>, S Dissing<sup>2</sup> and N E Skakkebaek<sup>1,3</sup>

<sup>1</sup>Department of Growth and Reproduction, Copenhagen University Hospital, Rigshospitalet, Denmark

<sup>2</sup>Department of Cellular and Molecular Medicine, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark

<sup>3</sup>International Center for Research and Research Training in Endocrine Disruption of Male Reproduction and Child Health (EDMaRC), University of Copenhagen, Rigshospitalet, Denmark

<sup>4</sup>Department of Biostatistics, University of Copenhagen, Copenhagen, Denmark

Correspondence should be addressed to N E Skakkebaek: [nes@rh.dk](mailto:nes@rh.dk)

This paper forms part of a special series on the effect of endocrine disrupting chemicals (EDCs) on development and male reproduction. This paper is based on work presented at the 9th Copenhagen Workshop on Endocrine Disruptors, 2–5 May 2017, Copenhagen, Denmark

## Abstract

Human sperm cell function must be precisely regulated to achieve natural fertilization. Progesterone released by the cumulus cells surrounding the egg induces a Ca<sup>2+</sup> influx into human sperm cells via the CatSper Ca<sup>2+</sup>-channel and thereby controls sperm function. Multiple chemical UV filters have been shown to induce a Ca<sup>2+</sup> influx through CatSper, thus mimicking the effect of progesterone on Ca<sup>2+</sup> signaling. We hypothesized that these UV filters could also mimic the effect of progesterone on sperm function. We examined 29 UV filters allowed in sunscreens in the US and/or EU for their ability to affect acrosome reaction, penetration, hyperactivation and viability in human sperm cells. We found that, similar to progesterone, the UV filters 4-MBC, 3-BC, Meradimate, Octisalate, BCSA, HMS and OD-PABA induced acrosome reaction and 3-BC increased sperm penetration into a viscous medium. The capacity of the UV filters to induce acrosome reaction and increase sperm penetration was positively associated with the ability of the UV filters to induce a Ca<sup>2+</sup> influx. None of the UV filters induced significant changes in the proportion of hyperactivated cells. In conclusion, chemical UV filters that mimic the effect of progesterone on Ca<sup>2+</sup> signaling in human sperm cells can similarly mimic the effect of progesterone on acrosome reaction and sperm penetration. Human exposure to these chemical UV filters may impair fertility by interfering with sperm function, e.g. through induction of premature acrosome reaction. Further studies are needed to confirm the results *in vivo*.

## Key Words

- ▶ endocrine disrupting chemicals
- ▶ UV filters
- ▶ fertility
- ▶ CatSper
- ▶ progesterone
- ▶ human sperm

*Endocrine Connections*  
(2018) 7, 16–25

## Introduction

Human male infertility is a common problem worldwide (1). The causes are in many cases unknown, but exposure to endocrine disrupting chemicals (EDCs) has been suspected to be involved (2, 3). Sperm cell dysfunction is a common cause of infertility (4) and intra-cytoplasmic sperm injection (ICSI), a method

developed to treat male infertility due to sperm dysfunction, is increasingly used in both the United States (5) and in Europe (6). The reasons for the increasing use of ICSI are unknown, but it has been hypothesized that environmental factors may play a role (1).

Sperm function must be precisely controlled, during the journey of the sperm cells through the female reproductive tract, for natural fertilization to occur (7, 8). Many sperm functions are controlled via the intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ), including sperm motility, chemotaxis and acrosome reaction (7). To be able to fertilize the egg, these individual  $[\text{Ca}^{2+}]_i$ -controlled sperm functions must be triggered at the correct time and in the correct order (7). CatSper (cationic channel of sperm) channels, located in the plasma membrane of the human sperm cell flagellum, are the principal facilitators of channel-mediated  $\text{Ca}^{2+}$  influx (9). CatSper is activated by the natural ligands progesterone and prostaglandins (10, 11), which lead to a rapid  $\text{Ca}^{2+}$  influx into the sperm cell. The cumulus cells surrounding the egg release progesterone and the progesterone-induced  $\text{Ca}^{2+}$  influx has been shown to mediate chemotaxis toward the egg (8, 12), to control sperm motility (13, 14) and to induce acrosome reaction (15).

CatSper can be promiscuously activated by various ligands (16), including multiple EDCs (17, 18, 19, 20, 21). Our recent study examined 29 of the 31 chemical UV filters allowed in sunscreens in the EU and/or US for their ability to induce a rise in  $[\text{Ca}^{2+}]_i$  in human sperm cells and showed that 13 chemical UV filters induced a rise in  $[\text{Ca}^{2+}]_i$  in human sperm cells (21). Nine of these seemed to induce a  $\text{Ca}^{2+}$  influx through interaction with CatSper, thereby mimicking the effect of progesterone. As the progesterone-induced  $\text{Ca}^{2+}$  influx controls important sperm cell functions, including sperm motility and acrosome reaction, we here examined the chemical UV filters for their ability to interfere with the human sperm cell functions acrosome reaction, sperm penetration into a viscous medium and hyperactivation, as well as with sperm viability.

## Materials and methods

### Reagents and chemical UV filters

We were able to obtain 30, out of the 31 chemical UV filters allowed in sunscreens in the EU and/or US (Table 1) from various chemical providers and to dissolve 29 of these in DMSO or ethanol as previously described (21). Progesterone, ionomycin, fluorescein isothiocyanate-conjugated *Pisum sativum* agglutinin (FITC-PSA) and 4000 cP methylcellulose were obtained from Sigma-Aldrich. Human serum albumin (HSA) was obtained from Irvine Scientific (CA, USA). Propidium iodide (PI),

Hoechst-33342 (Hoechst) and S100 were obtained from ChemoMetec A/S (Allerød, Denmark).

### Semen samples

All semen samples were produced by masturbation and ejaculated into clean, wide-mouthed plastic containers, on the same day as the experiment. After ejaculation, the samples were allowed to liquefy for 15–30 min at 37°C.

### Purification of motile sperm cells via swim-up

Motile spermatozoa were recovered from raw ejaculates by swim-up separation in human tubular fluid (HTF<sup>+</sup>) medium containing: 97.8 mM NaCl, 4.69 mM KCl, 0.2 mM  $\text{MgSO}_4$ , 0.37 mM  $\text{KH}_2\text{PO}_4$ , 2.04 mM  $\text{CaCl}_2$ , 0.33 mM Na-pyruvate, 21.4 mM Na-lactate, 2.78 mM glucose, 21 mM HEPES, and 4 mM  $\text{NaHCO}_3$ , adjusted to pH 7.3–7.4 with NaOH as described elsewhere (17). After 1 h at 37°C, the swim-up fraction was removed carefully and sperm concentration was determined by image cytometry as described in (22, 23). After two washes, the sperm samples were adjusted to  $10 \times 10^6/\text{mL}$  in HTF<sup>+</sup> with HSA (3 mg/mL) and the sperm cells were incubated for at least 1 h at 37°C.

For the experiments with capacitated sperm cells, the semen samples were instead adjusted to  $10 \times 10^6/\text{mL}$  (for acrosome reaction experiments) or  $20 \times 10^6/\text{mL}$  (for hyperactivation experiments) in a capacitating medium containing: 72.8 mM NaCl, 4.69 mM KCl, 0.2 mM  $\text{MgSO}_4$ , 0.37 mM  $\text{KH}_2\text{PO}_4$ , 2.04 mM  $\text{CaCl}_2$ , 0.33 mM Na-pyruvate, 21.4 mM Na-lactate, 2.78 mM glucose, 21 mM HEPES, and 25 mM  $\text{NaHCO}_3$ , adjusted to pH 7.3–7.4 with NaOH. 3 mg/mL (for acrosome reaction experiments) or 10 mg/mL (for hyperactivation experiments) HSA was added to the capacitating medium and the sperm cells were incubated for at least 3 h at 37°C in a 10%  $\text{CO}_2$  atmosphere as previously described (21). The higher HSA concentration for the hyperactivation experiments was used to minimize the sperm cells 'sticking-to-glass' phenomenon (24).

### Assessment of acrosome reaction

FITC-PSA can be used to stain the acrosome of sperm cells undergoing acrosome reaction (25, 26). Zoppino *et al.* have used FITC-PSA in combination with PI to identify viable acrosome-reacted sperm cells using flow cytometry (26). Here, we employ a similar approach using an image cytometer. A suspension of capacitated sperm cells with a sperm cell concentration of  $10 \times 10^6/\text{mL}$  was divided into equal aliquots and mixed with a staining solution

**Table 1** Chemical UV filters investigated. UV filters ranked according to their ability to induce Ca<sup>2+</sup> signals 10 μM (21).

Group	Rank	INCI name	CAS #	Abbreviation	Allowance in sunscreens	
					EU (%)	US (%)
UV filters that induce Ca <sup>2+</sup> signals at 10 μM	1	4-Methylbenzylidene camphor	36861-47-9/38102-62-4	4-MBC	4	
	2	3-Benzylidene camphor	15087-24-8	3-BC	2	
	3	Menthyl anthranilate	134-09-8	Meradimate		5
	4	Isoamyl P-methoxycinnamate	71617-10-2	Amiloxate	10	
	5	Ethylhexyl salicylate	118-60-5	Octisalate	5	5
	6	Benzylidene camphor sulfonic acid	56039-58-8	BCSA	6	
	7	Homosalate	118-56-9	HMS	10	15
	8	Ethylhexyl dimethyl PABA	21245-02-3	OD-PABA	8	8
	9	Benzophenone-3	131-57-7	BP-3	10	6
	10	Ethylhexyl methoxycinnamate	5466-77-3	Octinoxate	10	7.5
	11	Octocrylene	6197-30-4	Octocrylene	10	10
	12	Butyl methoxydibenzoylmethane	70356-09-1	Avobenzone	5	3
	UV filters that do not induce Ca <sup>2+</sup> signals at 10 μM	13	Diethylamino hydroxybenzoyl hexyl benzoate	302776-68-7	DHHB	10
14		Benzophenone-8	131-53-3	Dioxybenzone		3
15		Camphor benzalkonium methosulfate	52793-97-2	CBM	6	
16		Polysilicone-15	207574-74-1	Polysilicone-15	10	
17		Drometrizole trisiloxane	155633-54-8	Drometrizole trisolane	15	
18		Benzophenone-4	4065-45-6	BP-4	5	10
19		Diethylhexyl butamido triazone	154702-15-5	Iscotrizinol	10	
20		Ethylhexyl triazone	88122-99-0	Ethylhexyl triazone	5	
21		Cinoxate	104-28-9	Cinoxate		3
22		PEG-25 PABA	116242-27-4	PEG-25 PABA	10	
23		Bis-ethylhexyloxyphenol methoxyphenyl triazine	187393-00-6	Bemotrizinol	10	
24		Tea-salicylate	2174-16-5	TEA salicylate		12
25		Phenylbenzimidazole sulfonic acid	27503-81-7	Ensulizole	8	4
26		PABA	150-13-0	PABA		15
27		Disodium phenyl dibenzimidazole tetrasulfonate	180898-37-7	Bisdisulizole	10	
28		Benzophenone-5	6628-37-1	BP-5	5	
29		Terephthalylidene dicamphor sulfonic acid	92761-26-7/90457-82-2	Ecamsule	10	

Based on their ability to induce Ca<sup>2+</sup> signals, the UV filters are categorized into 'UV filters that induce Ca<sup>2+</sup> signals at 10 μM' and 'UV filters that do not induce Ca<sup>2+</sup> signals at 10 μM'. INCI name, CAS #, abbreviation and allowance in sunscreens in the EU and US are also listed in the table.

containing 5 μg/mL FITC-PSA and 0.5 μg/mL PI in HTF<sup>+</sup> as in (26). 10 μg/mL Hoechst was also added to the staining solution (see explanation below). Chemical UV filters (10 μM) were added to the aliquots of stained capacitated sperm cells. As positive controls, ionomycin (2 μM) and progesterone (10 μM) were added to separate aliquots. As a negative control, 0.2% DMSO was used, as this matched the DMSO concentration of ionomycin, which had the highest DMSO concentration of the treatments. After addition of chemical UV filters and controls, the samples were mixed and placed on a gentle mixing heating plate at 37°C. After 30 min of incubation, the aliquots were

thoroughly mixed by pipetting and a 50 μL sample was drawn and mixed with 100 μL of an immobilizing solution containing 0.6 M NaHCO<sub>3</sub> and 0.37% (v/v) formaldehyde in distilled water. This solution was mixed by pipetting and immediately loaded in an A2 slide (ChemoMetec A/S, Allerød, Denmark) and assessed in a NC-3000 image cytometer (ChemoMetec A/S). The following protocol was applied: 2-color flexicyte with Hoechst defining the sperm cells to be analyzed; Ex475-Em560/35: exposure time 3000 ms, Ex530-Em675/75: exposure time 500 ms, with a minimum of 5000 analyzed cells (positive for Hoechst). PI intensity as a function of FITC-PSA intensity was plotted

on bi-exponential scales, and specific quadrant gates were used to distinguish four groups:

1. PI-positive and FITC-PSA-positive cells: Acrosome-reacted nonviable sperm cells.
2. PI-negative and FITC-PSA positive cells: Acrosome-reacted viable sperm cells.
3. PI-positive and FITC-PSA-negative cells: Acrosome-intact nonviable sperm cells.
4. PI-negative and FITC-PSA-negative cells: Acrosome-intact viable sperm cells.

Control for spectral overlap between PSA and PI and definition of quadrant gates were carried out by labeling the cells singly with each fluorophore (data not shown). The obtained compensation matrix was applied to all measurements. To account for differences in capacitation between donors, only experiments with an induced positive increment of viable acrosome-reacted sperm cells for both positive controls compared to the negative control were included in the analysis.

#### Assessment of sperm penetration into a viscous medium

Sperm penetration tests with 4000 cP methylcellulose (1% w/v) as an artificial viscous medium were used as in (14). The methylcellulose (1% w/v) was prepared in HTF<sup>+</sup> by adding 10 mg methylcellulose per mL HTF<sup>+</sup> and mixing it by rotation overnight at RT. The methylcellulose (1% w/v) was introduced into glass capillary tubes (borosilicate microslides (VitroTubes) 0.20 mm × 2.0 mm × 10 cm (VitroCom, Mountain Lakes, NJ, USA)) by capillary forces, by placing the glass tubes vertically in a 1.5 mL microfuge tube with 750 μL methylcellulose (1% w/v) for 15 min. Care was taken to prevent air bubbles from entering the glass tubes. One end of the glass tube was sealed with wax (Hounisens laboratorieudstyr A/S, Jystrup, Denmark) and the open end was placed in a semen reservoir of a Kremer sperm penetration meter (R.B.M. Lab., Rødovre, Denmark). Just prior to the insertion of the glass tubes, either chemical UV filters (10 μM), 5 μM progesterone (positive control) or 0.1% DMSO (negative control) were added to a 200 μL non-capacitated sperm sample (10 × 10<sup>6</sup>/mL in HTF<sup>+</sup>) loaded into the semen reservoirs of the Kremer sperm penetration meter. The Kremer sperm penetration meter was tilted at a 45° angle and sperm cells were allowed to migrate into the methylcellulose (1% w/v) for 60 min at 37°C. The glass tube was then removed, wiped to

remove residual sperm cells from the surface of the glass and examined using phase contrast optics on an Olympus BX45 microscope at a total magnification of ×200 (Olympus). The number of sperm cells was counted at 1 cm distance from the base of the tube, with two fields in each of four planes counted. Throughout the study, all samples were analyzed by the same observer. Only experiments with a positive increment in cell density at 1 cm for the positive control compared to the negative control, with more than 50 sperm cells counted at 1 cm for the positive control, and with more than 10 cells counted for the negative control were used for the analysis.

#### Assessment of proportion of hyperactivated sperm cells with computer-assisted semen analysis (CASA)

A suspension of capacitated sperm cells with a sperm cell concentration of 20 × 10<sup>6</sup>/mL and a HSA concentration of 10 mg/mL was divided into equal aliquots and kept at 37°C. Just prior to acquisition of sperm motility data, either chemical UV filters (10 μM), progesterone (5 μM) or a negative control (0.1% DMSO), which matched the DMSO concentration of the chemical UV filters, was added to an aliquot of sperm sample. After mixing, a 4 μL sample was transferred to a 16 μm deep chamber (2 chambers (CASA) slide (CellVision, Oslo, Norway)), preheated to 37°C and placed on the heated motorized stage (37°C) of an Olympus BX41 microscope with a 20× phase contrast objective (Olympus). The microscope was connected to a computer running the Copenhagen Rigshospitalet Image House Sperm Motility Analysis System (CRISMAS), version 8.0.5919 CASA software. Sperm motility data were acquired just as the cells stopped drifting through the slide (took about 1 min) and was commenced <2 min after addition of chemical UV filters and controls. Motility characteristics were obtained at 60 Hz through a Basler camera acA640-120um (Basler AG, Ahrensburg, Germany). At least 200 sperm cells were counted on randomly selected fields in each sample and each sample was assessed in duplicates. Hyperactivated cells were identified using standard criteria: VCL ≥ 150 μm/s, linearity ≤ 50% and ALH ≥ 7 μm (27). As some samples contained clumps of immotile cells, we calculated the percentage of hyperactivated cells out of the total concentration of motile cells for each sample and used the mean value of the duplicates for further analysis.

### Assessment of sperm viability

Concentration of dead sperm cells was determined by image cytometry as in (22, 23), but using phosphate-buffered saline instead of S100 to dilute the sperm sample before running the assay. In this way, only the nonviable cells in the sample are stained with PI and counted.

### Ethical approval

Human semen samples were obtained from healthy volunteers with their prior consent. After delivery, the samples were fully anonymized. Each donor received a fee of 500 DKK (about 75 US dollars) per sample for their inconvenience. All samples were analyzed on the day of delivery and destroyed immediately after the laboratory experiments. Because of the full anonymization and the destruction of the samples immediately after the laboratory experiments, no ethical approval was needed for this work, according to the regional scientific ethical committee of the Capital Region of Denmark.

### Statistical analysis

All data were analyzed using two-way analysis of variance (ANOVA). This properly takes into account and adjusts for the considerable variation between donors as well as between experiments. By including positive and negative controls, the effect of the chemical UV filters can be given relative to a known control. The data were transformed with the natural logarithm to avoid variance heterogeneity and to obtain approximate normality of model residuals.

To display all data from each experiment in a single figure, we normalized the data relative to the positive control (acrosome reaction and viability data) or negative control (sperm penetration and hyperactivation data). *P* values were corrected for multiple comparison type I error inflation by Dunnett's method. To relate the ability of the chemical UV filter to induce a rise in  $[Ca^{2+}]_i$  to the ability to induce acrosome reaction or increase sperm penetration, we used the ability of the chemical UV filter to induce a rise in  $[Ca^{2+}]_i$  as a continuous covariate in the analysis. This results in a correct test for association between acrosome reaction or sperm penetration and the ability to induce a rise in  $[Ca^{2+}]_i$ , while taking into account the considerable variation between donors as well as between experiments. Statistical analyses were performed using proc mixed in SAS, version 9.4 (SAS Institute Inc., Cary, NC, USA).


## Results

### Effect on sperm acrosome reaction

Using an image-cytometer-based assay similar to that in (26), we investigated 29 chemical UV filters allowed in sunscreens in the EU and/or US (Table 1), for their ability to induce acrosome reaction in capacitated human sperm cells after 30 min of incubation. The chemical UV filters were tested at 10  $\mu$ M ( $n=3-5$ ), along with two positive controls (10  $\mu$ M progesterone and 2  $\mu$ M ionomycin) and a negative control (0.2% DMSO). A significant increase in viable acrosome-reacted sperm cells was found after treatment with the UV filters 4-MBC (adjusted *P* value <0.0001), 3-BC (adjusted *P* value <0.0001), Meradimate (adjusted *P* value <0.0001), HMS (adjusted *P* value <0.0001), Octisalate (adjusted *P*-value=0.0036), BCSA (adjusted *P* value=0.0241) and OD-PABA (adjusted *P* value=0.0425). A similar significant increase in viable acrosome-reacted sperm cells was found after treatment with progesterone (adjusted *P* value <0.0001). In order to display all data in a single figure, we calculated the percentage of viable acrosome-reacted sperm cells relative to the ionomycin-induced response from each individual experiment (Fig. 1). Additionally, we calculated the relationship between the ability of the UV filter at 10  $\mu$ M to induce a rise in  $[Ca^{2+}]_i$  (21) and to induce acrosome reaction and found a significant positive association (adjusted *P* value <0.0001) (Fig. 2).

### Effect on penetration into viscous medium


Using sperm penetration tests with methylcellulose (1% w/v) as in (14), we investigated the 29 chemical UV filters for their effect on sperm penetration into a viscous medium. The UV filters were tested at 10  $\mu$ M ( $n=3-6$ ), along with a positive control (5  $\mu$ M progesterone) and a negative control (0.1% DMSO). The increment in cell density at 1 cm was significantly increased after treatment with the UV filter 3-BC (adjusted *P* value=0.0347), similar to the increment observed after treatment with 5  $\mu$ M progesterone (adjusted *P* value=0.0001). In order to display all data in a single figure, we calculated the induced increment in cell density (in % of control) at 1 cm into the viscous medium (Fig. 3). Furthermore, we calculated the relationship between the ability of the UV filter at 10  $\mu$ M to induce a rise in  $[Ca^{2+}]_i$  (21) and to increase sperm penetration into viscous mucous and found a significant positive association (adjusted *P* value <0.0001) (Fig. 4).



**Figure 1**  
Viable acrosome-reacted sperm cells (in % of ionomycin-induced response) (mean  $\pm$  s.d.) after 30-min incubation with negative control (0.2% DMSO), positive control (10  $\mu$ M progesterone) and 10  $\mu$ M UV filters. The UV filters are ordered on the x-axis according to their ability to induce a rise in  $[Ca^{2+}]_i$ , (decreasing from left to right). The UV filters left to the vertical line induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M, whereas those right of the vertical line do not induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M (21). \*\*\*\*Adjusted  $P$  value  $\leq 0.0001$ ; \*\*\*adjusted  $P$  value  $\leq 0.001$ ; \*\*adjusted  $P$  value  $\leq 0.01$  and \*adjusted  $P$  value  $\leq 0.05$ .

### Effect on hyperactivation

Using computer-assisted semen analysis (CASA) we investigated the 29 chemical UV filters for effects on



**Figure 2**  
Scatter plot showing the ability of the chemical UV filter to induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M (in % of the paired progesterone-induced response (5  $\mu$ M)) (mean  $\pm$  s.d.) (21) on the x-axis and the ability to induce acrosome reaction in viable sperm cells (in % of ionomycin-induced response) (mean  $\pm$  s.d.) on the y-axis. The white dots indicate UV filters that induced a significant increment in the amount of viable acrosome-reacted cells. The line is obtained by linear regression and the  $P$  value for the association is obtained from the two-way ANOVA.


hyperactivation in sperm cells. The UV filters were tested at 10  $\mu$ M ( $n=3-4$ ), along with a positive control (5  $\mu$ M progesterone) and a negative control (0.1% DMSO). The percentage of hyperactivated sperm cells (in % of total motile cells) was not significantly changed after treatment with any of the UV filters or with progesterone (adjusted  $P$  values  $>0.8732$ ). In order to display the data in a single figure, we calculated the induced increment in hyperactivation (in % of control) (Supplementary Fig. 1, see section on supplementary data given at the end of this article).

### Effect on sperm viability

Using an image-cytometer-based assay, we tested the 29 chemical UV filters for their effect on sperm viability. We incubated aliquots of non-capacitated sperm cells with the UV filters at 10  $\mu$ M, along with a positive control (0.5% Triton) and a negative control (0.1% DMSO) for 20 h at 37°C. Viability was found to be significantly decreased after treatment with the UV filter Avobenzone (adjusted  $P$  value=0.0051) (Fig. 5).

### Discussion

Here, we investigated the effects of 29 chemical UV filters on the human sperm cell functions acrosome reaction, sperm penetration into a viscous medium and hyperactivation, as well as on sperm viability. We found




**Figure 3** Increment in cell density at 1 cm into a viscous medium (in % of negative control) (mean  $\pm$  s.d.) after treatment of sperm cells with negative control (0.1% DMSO), positive control (5  $\mu$ M progesterone) and 10  $\mu$ M UV filters ( $n=3-6$ ). The UV filters are ordered on the x-axis according to their ability to induce a rise in  $[Ca^{2+}]_i$ , (decreasing from left to right). The UV filters left to the vertical line induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M, whereas those right of the vertical line do not induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M (21). \*\*\*\*Adjusted  $P$  value  $\leq 0.0001$ ; \*adjusted  $P$  value  $\leq 0.05$ .

that only chemical UV filters, which had previously been shown to induce a rise in  $[Ca^{2+}]_i$  in human sperm cells (21), affected sperm cell functions. Seven of these UV filters: 4-MBC, 3-BC, Meradimate, Octisalate, BCSA, HMS and OD-PABA were found to induce acrosome reaction, similar to the response induced by progesterone. In addition, we showed that the UV filter 3-BC increased sperm penetration into a viscous medium, similar to the response induced by progesterone. The ability of the UV filters to induce acrosome reaction and increase sperm penetration was found to be positively associated with the ability of the chemical UV filter to induce a rise in  $[Ca^{2+}]_i$ . None of the UV filters induced a change in the proportion of hyperactivated cells and viability was only decreased after treatment with the UV filter Avobenzene. None of the chemical UV filters that did not induce a rise in  $[Ca^{2+}]_i$  in human sperm cells in our previous study (21) were found to affect sperm function.


Progesterone is a known inducer of acrosome reaction in human sperm cells (15) and a suboptimal induction of acrosome reaction in response to progesterone is associated with reduced male fertility (28, 29, 30, 31). An intact acrosome is required for mouse sperm cells to respond to progesterone-induced chemotaxis (32). Furthermore, only acrosome-intact human sperm cells can bind to the zona pellucida (33), in contrast to what has been found for mouse sperm cells (34, 35). Once bound to the zona pellucida, the human sperm cells must undergo acrosome reaction to penetrate the zona pellucida (36) and fuse with the egg (37). In line with this, a high level of spontaneous acrosome reaction has been associated

with reduced male fertility (38, 39, 40), although the relationship was not found in two other studies (31, 41). This suggests that exposure to chemical UV filters could



**Figure 4** Scatter plot showing the ability of the chemical UV filter to induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M (in % of the paired progesterone-induced response (5  $\mu$ M)) (mean  $\pm$  s.d.) (21) on the x-axis and the increment in cell density at 1 cm into a viscous medium (in % of negative control) (mean  $\pm$  s.d.) on the y-axis. The white dot indicates the UV filter 3-BC, which induced a significant increment in cell density at 1 cm into a viscous medium. The line is obtained by linear regression and the  $P$  value for the association is obtained from the two-way ANOVA.





**Figure 5** Nonviable cells (%) after 20h of incubation with 0.1% DMSO (negative control) and UV filters at 10  $\mu$ M (mean  $\pm$  s.d.). The UV filters are ordered on the x-axis according to their ability to induce a rise in  $[Ca^{2+}]_i$ , (decreasing from left to right). The UV filters left to the vertical line induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M, whereas those right of the vertical line do not induce a rise in  $[Ca^{2+}]_i$  at 10  $\mu$ M (21). \*\*Adjusted *P*-value  $\leq 0.01$ .

impair fertility by inducing premature acrosome reaction in human sperm cells.

In support of our findings, p,p'-DDE has been shown to induce a rise in  $[Ca^{2+}]_i$  via CatSper and acrosome reaction (18). Similarly, in our previous study (17), we showed that the chemical UV filters 4-MBC and 3-BC could induce a rise in  $[Ca^{2+}]_i$  via CatSper and acrosome reaction. Our results here confirmed these findings for 4-MBC and 3-BC. Also, triclosan has been shown to induce a CatSper-independent rise in  $[Ca^{2+}]_i$  and acrosome reaction (17).

In contrast to our findings, diethylstilbestrol (DES) was found neither to induce acrosome reaction nor increase sperm penetration, even though it was found to induce a  $Ca^{2+}$  influx via CatSper (20). Methodological differences might account for these contradicting findings. Unlike in our study, Zou *et al.* added DES to non-capacitated sperm cells and allowed the sperm cells to incubate with DES for 4h before assessing acrosome reaction or sperm penetration. In our study, we on the other hand added the chemical UV filters to already capacitated sperm cells 30min before assessing acrosome reaction and to non-capacitated sperm cells just prior to assessing sperm penetration.

Interestingly, however, DES was found to dose-dependently inhibit both the progesterone-induced rise in  $[Ca^{2+}]_i$ , acrosome reaction and sperm penetration (20). We have previously shown that the UV filters 4-MBC (17), 3-BC and BCSA (21) can competitively inhibit the progesterone-induced rise in  $[Ca^{2+}]_i$ , indicating that these UV filters might similarly be able to inhibit the progesterone-induced acrosome reaction and sperm penetration.

Progesterone is a weak inducer of hyperactivation, inducing only a small increment in the proportion of hyperactivated cells (13, 14), with no relationship between the induced rise in  $[Ca^{2+}]_i$  and hyperactivation response (13). In our study, neither progesterone, nor the chemical UV filters, induced hyperactivation. In our previous study (17), 4-MBC was shown to lower the frequency and enhance the asymmetry of the flagellar beat in a single sperm cell, indicating that 4-MBC could induce hyperactivation. With the experimental setup in our study we could, however, not find an increase in hyperactivation after treatment with 4-MBC on a sperm cell population. Studies have shown that only a given proportion of sperm cells in a population respond to treatment with a  $Ca^{2+}$  signal-inducing EDC (18, 21), probably due to the heterogeneity of sperm samples (42). These findings could explain how hyperactivation can be induced in individual sperm cells, while the proportion of hyperactivated cells in the whole sperm population remains relatively stable.

Most UV filters tested did not affect viability, similar to DES (20) and p,p'-DDE upon one day of incubation (18). Taken together, our data are consistent with the notion that the induced rise in  $[Ca^{2+}]_i$  in human sperm cells on itself does not affect sperm viability, and that the adverse effect of Avobenzene on viability is most likely independent from its effect on  $[Ca^{2+}]_i$ .

Multiple EDCs have been shown to induce a rise in  $[Ca^{2+}]_i$  in human sperm cells through interaction with CatSper (17, 18, 19, 20, 21), as have multiple pharmacological ligands (43). Our findings for the chemical UV filters tested here indicate that other compounds that activate CatSper could similarly affect

sperm function in a progesterone-like manner. EDCs have been shown to act additively to induce a rise in  $[Ca^{2+}]_i$  in human sperm cells (17, 21), suggesting that these EDCs could similarly act additively to induce acrosome reaction and increase sperm penetration.

In conclusion, several chemical UV filters known to mimic the effect of progesterone on  $Ca^{2+}$  signaling in human sperm cells were shown to induce acrosome reaction and sperm penetration in a progesterone-like manner. Exposure to these chemical UV filters could impair fertility by interfering with sperm function, e.g. through induction of premature acrosome reaction. Further studies are needed to confirm our results *in vivo*.

#### Supplementary data

This is linked to the online version of the paper at <https://doi.org/10.1530/EC-17-0156>.

#### Declaration of interest

The authors declare there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

#### Funding

This study was supported by a PhD Internship Scholarship from the Faculty of Health and Medical Sciences, University of Copenhagen, an EDMaRC research grant from the Kirsten and Freddy Johansen's Foundation, and the Innovation Fund Denmark (InnovationsFonden, grant number 14-2013-4).

#### Author contribution statement

Study design: A R, N E S; execution: A R, D L E; analysis: A R, J H P; manuscript drafting: A R, S D, N E S and critical discussion: A R, D L E, K A, J H P, S D, N E S.

#### Acknowledgments

The author would like to thank Ina Lund for her technical assistance with the semen donor corps and the swim-up preparation of the semen samples, as well as for her help with the sperm penetration tests and CASA experiments.

## References

- Skakkebaek NE, Rajpert-De Meyts E, Buck Louis GM, Toppari J, Andersson A-M, Eisenberg ML, Jensen TK, Jørgensen N, Swan SH, Sapra KJ, *et al.* Male reproductive disorders and fertility trends: influences of environment and genetic susceptibility. *Physiological Reviews* 2016 **96** 55–97. (<https://doi.org/10.1152/physrev.00017.2015>)
- Diamanti-Kandaraki E, Bourguignon J-P, Giudice LC, Hauser R, Prins GS, Soto AM, Zoeller RT & Gore AC. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocrine Reviews* 2009 **30** 293–342. (<https://doi.org/10.1210/er.2009-0002>)
- Bergman A, Heindel JJ, Kasten T, Kidd KA, Jobling S, Neira M, Zoeller RT, Becher G, Bjerregaard P, Bornman R, *et al.* The impact of endocrine disruption: a consensus statement on the state of the science. *Environmental Health Perspectives* 2013 **121** A104–A106. (<https://doi.org/10.1289/ehp.1205448>)
- Hull MG, Glazener CM, Kelly NJ, Conway DI, Foster PA, Hinton RA, Coulson C, Lambert PA, Watt EM & Desai KM. Population study of causes, treatment, and outcome of infertility. *BMJ* 1985 **291** 1693–1697. (<https://doi.org/10.1136/bmj.291.6510.1693>)
- Jain T & Gupta RS. Trends in the use of intracytoplasmic sperm injection in the United States. *New England Journal of Medicine* 2007 **357** 251–257. (<https://doi.org/10.1056/NEJMsa070707>)
- Kupka MS, Ferraretti AP, de Mouzon J, Erb K, D'Hooghe T, Castilla JA, Calhaz-Jorge C, De Geyter C, Goossens V & European IVF-Monitoring Consortium, for the European Society of Human Reproduction and Embryology. Assisted reproductive technology in Europe, 2010: results generated from European registers by ESHRE. *Human Reproduction* 2014 **29** 2099–2113. (<https://doi.org/10.1093/humrep/deu175>)
- Publicover S, Harper CV & Barratt C.  $[Ca^{2+}]_i$  signalling in sperm—making the most of what you've got. *Nature Cell Biology* 2007 **9** 235–242. (<https://doi.org/10.1038/ncb0307-235>)
- Publicover SJ, Giojalas LC, Teves ME, de Oliveira GSMM, Garcia AAM, Barratt CLR & Harper CV.  $Ca^{2+}$  signalling in the control of motility and guidance in mammalian sperm. *Frontiers in Bioscience* 2008 **13** 5623–5637. (<https://doi.org/10.2741/3105>)
- Lishko PV, Kirichok Y, Ren D, Navarro B, Chung J-J & Clapham DE. The control of male fertility by spermatozoan ion channels. *Annual Review of Physiology* 2012 **74** 453–475. (<https://doi.org/10.1146/annurev-physiol-020911-153258>)
- Strünker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R & Kaupp UB. The CatSper channel mediates progesterone-induced  $Ca^{2+}$  influx in human sperm. *Nature* 2011 **471** 382–386. (<https://doi.org/10.1038/nature09769>)
- Lishko PV, Botchkina IL & Kirichok Y. Progesterone activates the principal  $Ca^{2+}$  channel of human sperm. *Nature* 2011 **471** 387–391. (<https://doi.org/10.1038/nature09767>)
- Eisenbach M & Giojalas LC. Sperm guidance in mammals – an unpaved road to the egg. *Nature Reviews Molecular Cell Biology* 2006 **7** 276–285. (<https://doi.org/10.1038/nrm1893>)
- Alasmari W, Barratt CLR, Publicover SJ, Whalley KM, Foster E, Kay V, Martins da Silva S & Oxenham SK. The clinical significance of calcium-signalling pathways mediating human sperm hyperactivation. *Human Reproduction* 2013 **28** 866–876. (<https://doi.org/10.1093/humrep/des467>)
- Alasmari W, Costello S, Correia J, Oxenham SK, Morris J, Fernandes L, Ramalho-Santos J, Kirkman-Brown J, Michelangeli F, Publicover S, *et al.*  $Ca^{2+}$  signals generated by CatSper and  $Ca^{2+}$  stores regulate different behaviors in human sperm. *Journal of Biological Chemistry* 2013 **288** 6248–6258. (<https://doi.org/10.1074/jbc.M112.439356>)
- Tamburrino L, Marchiani S, Minetti F, Forti G, Muratori M & Baldi E. The CatSper calcium channel in human sperm: relation with motility and involvement in progesterone-induced acrosome reaction. *Human Reproduction* 2014 **29** 418–428. (<https://doi.org/10.1093/humrep/det454>)
- Brenker C, Goodwin N, Weyand I, Kashikar ND, Naruse M, Krähling M, Müller A, Kaupp UB & Strünker T. The CatSper channel: a polymodal chemosensor in human sperm. *EMBO Journal* 2012 **31** 1654–1665. (<https://doi.org/10.1038/emboj.2012.30>)
- Schiffer C, Müller A, Egeberg DL, Alvarez L, Brenker C, Rehfeld A, Frederiksen H, Wäschle B, Kaupp UB, Balbach M, *et al.* Direct action of endocrine disrupting chemicals on human sperm. *EMBO Reports* 2014 **15** 758–765. (<https://doi.org/10.15252/embr.201438869>)
- Tavares RS, Mansell S, Barratt CLR, Wilson SM, Publicover SJ & Ramalho-Santos J. p,p'-DDE activates CatSper and compromises

- human sperm function at environmentally relevant concentrations. *Human Reproduction* 2013 **28** 3167–3177. (<https://doi.org/10.1093/humrep/det372>)
- 19 Shannon M, Rehfeld A, Frizzell C, Livingstone C, McGonagle C, Skakkebaek NE, Wielogórska E & Connolly L. In vitro bioassay investigations of the endocrine disrupting potential of steviol glycosides and their metabolite steviol, components of the natural sweetener Stevia. *Molecular and Cellular Endocrinology* 2016 **427** 65–72. (<https://doi.org/10.1016/j.mce.2016.03.005>)
  - 20 Zou Q-X, Peng Z, Zhao Q, Chen H-Y, Cheng Y-M, Liu Q, He YQ, Weng SQ, Wang HF, Wang T, *et al.* Diethylstilbestrol activates CatSper and disturbs progesterone actions in human spermatozoa. *Human Reproduction* 2017 **32** 290–298. (<https://doi.org/10.1093/humrep/dew332>)
  - 21 Rehfeld A, Dissing S & Skakkebaek NE. Chemical UV filters mimic the effect of progesterone on Ca(2+) signaling in human sperm cells. *Endocrinology* 2016 **157** 4297–4308. (<https://doi.org/10.1210/en.2016-1473>)
  - 22 Egeberg DL, Kjaerulf S, Hansen C, Petersen JH, Glensbjerg M, Skakkebaek NE, Jørgensen N & Almstrup K. Image cytometer method for automated assessment of human spermatozoa concentration. *Andrology* 2013 **1** 615–623. (<https://doi.org/10.1111/j.2047-2927.2013.00082.x>)
  - 23 Egeberg Palme DL, Johannsen TH, Petersen JH, Skakkebaek NE, Juul A, Jørgensen N & Almstrup K. Validation of image cytometry for sperm concentration measurement: comparison with manual counting of 4010 human semen samples. *Clinica Chimica Acta* 2017 **468** 114–119. (<https://doi.org/10.1016/j.cca.2017.02.014>)
  - 24 ESHRE Andrology Special Interest group. Guidelines on the application of CASA technology in the analysis of spermatozoa. ESHRE Andrology Special Interest Group. European Society for Human Reproduction and Embryology. *Human Reproduction* 1998 **13** 142–145.
  - 25 Sánchez-Cárdenas C, Servín-Vences MR, José O, Treviño CL, Hernández-Cruz A & Darszon A. Acrosome reaction and Ca<sup>2+</sup> imaging in single human spermatozoa: new regulatory roles of [Ca<sup>2+</sup>]<sub>i</sub>. *Biology of Reproduction* 2014 **91** 67. (<https://doi.org/10.1095/biolreprod.114.119768>)
  - 26 Zoppino FCM, Halón ND, Bustos MA, Pavarotti MA & Mayorga LS. Recording and sorting live human sperm undergoing acrosome reaction. *Fertility and Sterility* 2012 **97** 1309–1315. (<https://doi.org/10.1016/j.fertnstert.2012.03.002>)
  - 27 Mortimer ST, Swan MA & Mortimer D. Effect of seminal plasma on capacitation and hyperactivation in human spermatozoa. *Human Reproduction* 1998 **13** 2139–2146. (<https://doi.org/10.1093/humrep/13.8.2139>)
  - 28 Krausz C, Bonaccorsi L, Luconi M, Fuzzi B, Criscuoli L, Pellegrini S, Forti G & Baldi E. Intracellular calcium increase and acrosome reaction in response to progesterone in human spermatozoa are correlated with in-vitro fertilization. *Human Reproduction* 1995 **10** 120–124. (<https://doi.org/10.1093/humrep/10.1.120>)
  - 29 Oehninger S, Blackmore P, Morshedi M, Sueldo C, Acosta AA & Alexander NJ. Defective calcium influx and acrosome reaction (spontaneous and progesterone-induced) in spermatozoa of infertile men with severe teratozoospermia. *Fertility and Sterility* 1994 **61** 349–354. ([https://doi.org/10.1016/S0015-0282\(16\)56530-3](https://doi.org/10.1016/S0015-0282(16)56530-3))
  - 30 Falsetti C, Baldi E, Krausz C, Casano R, Failli P & Forti G. Decreased responsiveness to progesterone of spermatozoa in oligozoospermic patients. *Journal of Andrology* 1993 **14** 17–22.
  - 31 Krausz C, Bonaccorsi L, Maggio P, Luconi M, Criscuoli L, Fuzzi B, Pellegrini S, Forti G & Baldi E. Two functional assays of sperm responsiveness to progesterone and their predictive values in in-vitro fertilization. *Human Reproduction* 1996 **11** 1661–1667. (<https://doi.org/10.1093/oxfordjournals.humrep.a019466>)
  - 32 Guidobaldi HA, Hirohashi N, Cubilla M, Buffone MG & Giojalas LC. An intact acrosome is required for the chemotactic response to progesterone in mouse spermatozoa. *Molecular Reproduction and Development* 2017 **84** 310–315. (<https://doi.org/10.1002/mrd.22782>)
  - 33 Liu DY, Garrett C & Baker HWG. Acrosome-reacted human sperm in insemination medium do not bind to the zona pellucida of human oocytes. *International Journal of Andrology* 2006 **29** 475–481. (<https://doi.org/10.1111/j.1365-2605.2006.00681.x>)
  - 34 Inoue N, Satouh Y, Ikawa M, Okabe M & Yanagimachi R. Acrosome-reacted mouse spermatozoa recovered from the perivitelline space can fertilize other eggs. *PNAS* 2011 **108** 20008–20011. (<https://doi.org/10.1073/pnas.1116965108>)
  - 35 Jin M, Fujiwara E, Kakiuchi Y, Okabe M, Satouh Y, Baba SA, Chiba K & Hirohashi N. Most fertilizing mouse spermatozoa begin their acrosome reaction before contact with the zona pellucida during in vitro fertilization. *PNAS* 2011 **108** 4892–4896. (<https://doi.org/10.1073/pnas.1018202108>)
  - 36 Liu DY & Baker HW. Inhibition of acrosin activity with a trypsin inhibitor blocks human sperm penetration of the zona pellucida. *Biology of Reproduction* 1993 **48** 340–348. (<https://doi.org/10.1095/biolreprod48.2.340>)
  - 37 Wassarman PM, Jovine L & Litscher ES. A profile of fertilization in mammals. *Nature Cell Biology* 2001 **3** E59–E64. (<https://doi.org/10.1038/35055178>)
  - 38 Fénelichel P, Donzeau M, Farahifar D, Basteris B, Ayraud N & Hsi BL. Dynamics of human sperm acrosome reaction: relation with in vitro fertilization. *Fertility and Sterility* 1991 **55** 994–999.
  - 39 Liu DY & Baker HW. Tests of human sperm function and fertilization in vitro. *Fertility and Sterility* 1992 **58** 465–483. ([https://doi.org/10.1016/S0015-0282\(16\)55247-9](https://doi.org/10.1016/S0015-0282(16)55247-9))
  - 40 Takahashi K, Wetzels AM, Goverde HJ, Bastaans BA, Janssen HJ & Rolland R. The kinetics of the acrosome reaction of human spermatozoa and its correlation with in vitro fertilization. *Fertility and Sterility* 1992 **57** 889–894. ([https://doi.org/10.1016/S0015-0282\(16\)54976-0](https://doi.org/10.1016/S0015-0282(16)54976-0))
  - 41 Cummins JM, Pember SM, Jequier AM, Yovich JL & Hartmann PE. A test of the human sperm acrosome reaction following ionophore challenge. Relationship to fertility and other seminal parameters. *Journal of Andrology* 1991 **12** 98–103.
  - 42 Okabe M. The cell biology of mammalian fertilization. *Development* 2013 **140** 4471–4479. (<https://doi.org/10.1242/dev.090613>)
  - 43 Martins da Silva SJ, Brown SG, Sutton K, King LV, Ruso H, Gray DW, Wyatt PG, Kelly MC, Barratt CLR & Hope AG. Drug discovery for male subfertility using high-throughput screening: a new approach to an unsolved problem. *Human Reproduction* 2017 **16** 1–11. (<https://doi.org/10.1093/humrep/dex055>)

Received in final form 22 August 2017

Accepted 5 September 2017

Accepted Preprint published online 5 September 2017