



## **The relationship between genetic risk variants with brain structure and function in bipolar disorder**

Pereira, Licia P; Köhler, Cristiano A; de Sousa, Rafael T; Solmi, Marco; de Freitas, Bárbara P; Fornaro, Michele; Machado-Vieira, Rodrigo; Miskowiak, Kamilla W; Vieta, Eduard; Veronese, Nicola; Stubbs, Brendon; Carvalho, André F

*Published in:*  
Neuroscience & Biobehavioral Reviews

*DOI:*  
[10.1016/j.neubiorev.2017.05.002](https://doi.org/10.1016/j.neubiorev.2017.05.002)

*Publication date:*  
2017

*Document version*  
Peer reviewed version

*Document license:*  
[CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

*Citation for published version (APA):*  
Pereira, L. P., Köhler, C. A., de Sousa, R. T., Solmi, M., de Freitas, B. P., Fornaro, M., ... Carvalho, A. F. (2017). The relationship between genetic risk variants with brain structure and function in bipolar disorder: A systematic review of genetic-neuroimaging studies. *Neuroscience & Biobehavioral Reviews*, 79, 87-109. <https://doi.org/10.1016/j.neubiorev.2017.05.002>

## Accepted Manuscript

Title: The relationship between genetic risk variants with brain structure and function in bipolar disorder: A systematic review of genetic-neuroimaging studies

Authors: Licia P. Pereira, Cristiano A. Köhler, Rafael T. Sousa, Marco Solmi, Bárbara P. de Freitas, Michele Fornaro, Rodrigo Machado-Vieira, Kamilla W. Miskowiak, Eduard Vieta, Nicola Veronese, Brendon Stubbs, André F. Carvalho



PII: S0149-7634(17)30030-1  
DOI: <http://dx.doi.org/doi:10.1016/j.neubiorev.2017.05.002>  
Reference: NBR 2842

To appear in:

Received date: 13-1-2017  
Revised date: 27-4-2017  
Accepted date: 1-5-2017

Please cite this article as: Pereira, Licia P., Köhler, Cristiano A., Sousa, Rafael T., Solmi, Marco, de Freitas, Bárbara P., Fornaro, Michele, Machado-Vieira, Rodrigo, Miskowiak, Kamilla W., Vieta, Eduard, Veronese, Nicola, Stubbs, Brendon, Carvalho, André F., The relationship between genetic risk variants with brain structure and function in bipolar disorder: A systematic review of genetic-neuroimaging studies. *Neuroscience and Biobehavioral Reviews* <http://dx.doi.org/10.1016/j.neubiorev.2017.05.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## The relationship between genetic risk variants with brain structure and function in bipolar disorder: A systematic review of genetic-neuroimaging studies

Licia P. Pereira<sup>1</sup>, Cristiano A. Köhler<sup>1</sup>, Rafael T. Sousa<sup>2</sup>, Marco Solmi<sup>3,4</sup>, Bárbara P. de Freitas<sup>1</sup>, Michele Fornaro<sup>5</sup>, Rodrigo Machado-Vieira<sup>2</sup>, Kamilla W. Miskowiak<sup>6</sup>, Eduard Vieta<sup>7</sup>, Nicola Veronese<sup>4,8</sup>, Brendon Stubbs<sup>9,4</sup>, André F. Carvalho<sup>1,4,\*</sup>

<sup>1</sup> Department of Clinical Medicine and Translational Psychiatry Research Group, Faculty of Medicine, Federal University of Ceará, Fortaleza, CE, Brazil.

<sup>2</sup> Experimental Therapeutics and Pathophysiology Branch, National Institute of Mental Health, NIH, Bethesda, MD, USA.

<sup>3</sup> Department of Neuroscience, University of Padova, Padova, Italy;

<sup>4</sup> Institute for Clinical Research and Education in Medicine (IREM), Padova, Italy;

<sup>5</sup> New York State Psychiatric Institute (NYSPI), Columbia University, New York, NY, USA;

<sup>6</sup> Copenhagen Psychiatric Centre, Copenhagen University Hospital, Rigshospitalet, Denmark;

<sup>7</sup> Bipolar Unit, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain;

<sup>8</sup> National Research Council, Neuroscience Institute, Aging Branch, Padova, Italy;

<sup>9</sup> Physiotherapy Department, South London and Maudsley NHS Foundation Trust, Denmark Hill, London SE5 8AZ, United Kingdom; Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London Box SE5 8AF, United Kingdom; Faculty of Health, Social Care and Education, Anglia Ruskin University, Bishop Hall Lane, Chelmsford CM1 1SQ, United Kingdom.

**Running title:** Genetic-neuroimaging and bipolar disorder

**Word count:** 6356

**Figures/Tables:** 2/3

**Supplementary online material**

\*Corresponding Author:

**André F. Carvalho, MD, PhD**

Department of Clinical Medicine, Faculty of Medicine, Federal University of Ceará

Rua Prof. Costa Mendes, 1608, 4º andar

Fortaleza, 60430-040, Brazil

Phone/Fax: +558532617227

E-mail: andrefc7@terra.com.br or andrefc7@hotmail.com

**Highlights**

- Genetic-neuroimaging paradigms may aid in the understanding of the neurobiology of heterogeneous phenotypes like bipolar disorder;
- A systematic review of gene-imaging studies conducted in samples with bipolar disorder was performed;
- Forty-four studies (N=2,122 participants with bipolar disorder) met inclusion criteria;
- Replicated evidence suggests that individuals with BD who carry the BDNF Val66Met allele may have smaller hippocampi volumes;
- Studies that employed a genome-wide associated approach failed to reveal statistically significant findings.

**Abstract**

Genetic-neuroimaging paradigms could provide insights regarding the pathophysiology of bipolar disorder (BD). Nevertheless, findings have been inconsistent across studies. A systematic review of gene-imaging studies involving individuals with BD was conducted across electronic major databases from inception until January 9<sup>th</sup>, 2017. Forty-four studies met eligibility criteria (N=2,122 BD participants). Twenty-six gene variants were investigated across candidate gene studies and 4 studies used a genome-wide association approach. Replicated evidence (i.e. in >2 studies) suggests that individuals with BD carrying the BDNF Val66Met risk allele could have reduced hippocampal volumes compared to non-carriers. This review underscores the potential of gene-neuroimaging paradigms to provide mechanistic insights for BD. However, this systematic review found a single replicated finding. Suggestions to improve the reproducibility of this emerging field are provided, including the adoption of a trans-diagnostic approach.

**Abbreviations**

Anterior cingulum, AC; Adverse Childhood Experiences, ACE; Anterior cingulate gyrus, ACG; Anterior limb of internal capsule, ALIC; Ankyrin 3, ANK3; Bipolar

disorder, BD; Brain-derived neurotrophic factor, BDNF; Blood oxygen level dependent, BOLD; Calcium voltage-gated channel subunit alpha 1 C, CACNA1C; Corpus callosum, CC; Corpus Callosum, body, CCb; Corpus Callosum, genu, CCg; Cingulate gyrus, CG; 2',3'-Cyclic-nucleotide 3'-phosphodiesterase, CNP; Catechol-O-methyltransferase, COMT; Corona radiata, CR; Corticospinal tract, CST; D-amino acid oxidase, DAAO; D-amino acid oxidase activator, DAOA; Diacylglycerol kinase eta, DGKH; Disrupted in schizophrenia 1, DISC1; Dorsolateral prefrontal cortex, dlPFC; Default mode network, DMN; Docking protein 5, DOK5; Diffusion tensor imaging, DTI; Excitatory amino-acid transporter 2, EAAT2; Erb-B2 Receptor Tyrosine Kinase 2, ERBB2; Fractional anisotropy, FA; Forceps major, FM; Functional MRI, fMRI; Fusiform gyrus, FG; Fronto-occipital fasciculus, FOF; Gamma-Amino Butyric Acid, GABA; Polypeptide N-Acetylgalactosaminyltransferase 7, GALNT7; Gyrfication index, GI; Grey matter, GM; Globus pallidus, GP; Glutamate ionotropic receptor NMDA type subunit 2B, GRIN2B; Glycogen synthase kinase 3 beta, GSK-3 $\beta$ ; Genome-wide association study, GWAS; Risk haplotype at the 5' end of the NRG1 gene, HAP; Healthy controls, HCs; Inferior cerebellar peduncle, ICP; Interleukin-1 beta, IL-1 $\beta$ ; Inferior parietal lobule, IPL; Inferior occipital gyrus, IOG; Lateral ventricles, LV; Longitudinal fasciculus, LF; Minor allele, MA; Middle cerebellar peduncle, MCP; Mean diffusivity, MD; Myelin oligodendrocyte glycoprotein, MOG; Medial prefrontal cortex, mPFC; Magnetic resonance imaging, MRI; Middle temporal gyrus, MTG; Nucleus accumbens (NAc), NA; N-methyl-D-aspartate, NMDA; Neuregulin 1, NRG1; Teneurin transmembrane protein 4, ODZ4; Orbitofrontal cortex, OFC; Posterior cingulate gyrus, PCG; Prefrontal region, PF; Prefrontal cortex, PFC; Polygenic risk score, PGR; Parahippocampal gyrus, PHG; Radial diffusivity, RD; SZ, Schizophrenia; Serotonin-transporter-linked polymorphic region, 5-HTTLPR; Single

nucleotide polymorphism, SNP; Spectrin repeat containing nuclear envelope protein 1, SYNE1; Sterol regulatory element-binding transcription factor 1, SREBF1; Sterol regulatory element-binding transcription factor 2, SREBF2; Superior Temporal Gyrus, STG; Tract-based spatial statistics, TBSSBD; Tumor necrosis factor, TNF; Temporal pole, TP; Thalamic radiation, TR; Uncinate fasciculus, UF; Voxel-based morphometry, VBM; Ventrolateral prefrontal cortex, vIPFC; White matter, WM; Zinc finger protein 804A, ZNF804A

**Key words:** Bipolar disorder; genetic polymorphisms; neuroimaging; magnetic resonance imaging; functional MRI; diffusion tensor imaging, voxel-based morphometry.

## 1. Introduction

Bipolar disorder (BD) may affect approximately 2.4% of the population worldwide, and is associated with significant disability and elevated mortality rates compared to the general population (Grande et al., 2016; Hayes et al., 2015; Merikangas et al., 2011). The pathophysiology of BD has not been completely elucidated, and the current state of knowledge on putative mechanisms underpinning different clinical features and illness trajectories is limited (Craddock and Sklar, 2013; Hasler and Wolf, 2015). Several lines of evidence indicate that hereditary factors play a relevant role in the patho-etiology of BD, with phenotypic concordance rates ranging from 40-70% in monozygotic twins, and 8-10% in first-degree relatives (FDRs) (Kerner, 2014; Smoller and Finn, 2003). Genome-wide significant loci for BD have emerged from meta-analyses of GWAS, while loci near the *TRANK1*, *ANK3*, *ODZ4*, *CACNA1C*, and *NCAN* genes had at least one additional replication (Goes, 2016; Green et al., 2013; Muhleisen et al., 2014). A

recent GWAS identified two additional novel loci associated with bipolar disorder i.e. an inter-genic region on 9p21.3 and markers within *ERBB2* (Hou et al., 2016). In addition, the *CACNA1C* gene differed in expression in the prefrontal cortex of patients with BD compared to controls (Nurnberger et al., 2014). However, identified genome-wide significant signals seem to explain a low proportion of phenotypic variance of BD (Goes, 2016), and a polygenic risk score accounts for only 3% of its phenotypic variance (Group, 2011). It has been proposed that the effects of risk genes for BD could be larger and more evident on intermediate phenotypes neurobiologically linked to the disorder, thus providing an impetus to the emergence of ‘gene imaging’ studies in the literature (Bigos and Weinberger, 2010; Gurung and Prata, 2015; Ivleva et al., 2010).

Precise mechanisms through which genetic variations may influence neural pathways accounting for the phenotypic heterogeneity of BD are yet to be established. Significant efforts have been conducted to identify phenotypic characteristics that are thought to lie more proximal to the genetic factors (i.e. endophenotypes) with the aim that this approach would aid in the identification of biological mechanisms of BD (Gottesman and Gould, 2003; Kurnianingsih et al., 2011). In this context, a large body of literature indicates that BD is associated with significant functional and structural neuroimaging alterations (Kempton et al., 2011; Kupferschmidt and Zakzanis, 2011). Furthermore, meta-analytic evidence indicates that functional and structural neuroimaging abnormalities may be evidence in individuals at-risk for BD (Fusar-Poli et al., 2012), and a recent systematic review indicates that functional and structural neuroimaging abnormalities are also evident in healthy FDRs of patients with BD (Piguet et al., 2015). Altogether this literature provides support to the view that subtler functional and structural neuroimaging abnormalities in at-risk individuals could represent vulnerability markers of BD. ‘Imaging genetics’ has emerged as a field with

an underlying rationale that genetic variations that confer risk to mental disorders may exhibit higher penetrance at such brain functional/structural alterations than at the more distal psychopathological/behavioral levels (Hashimoto et al., 2015; Rasetti and Weinberger, 2011). Hence, an ever-increasing number of studies has attempted to investigate the associations between genetic variations expected to play a pathophysiological role in BD and structural and functional neuroimaging abnormalities. However, different age groups, neuroimaging modalities, treatment-related effects and investigated genes (or polygenic risk scores) are potential confounders which might have contributed to the heterogeneity of studies so far (Kurnianingsih et al., 2011). To overcome such a strong heterogeneity a systematic review of ‘neuroimaging genetics’ studies which considered genes which have been previously found to reach genome-wide significance in schizophrenia and BD was conducted (Gurung and Prata, 2015; Lee et al., 2012). However, this previous systematic review considered studies performed solely in healthy individuals, while only seven studies performed in samples with BD were included (Gurung and Prata, 2015). A comprehensive systematic overview focusing on ‘imaging genetics’ specifically in people with BD is currently lacking.

Therefore, our systematic review aims to provide a comprehensive and up-dated synthesis of all available ‘imaging genetics’ literature in BD. Both structural and functional magnetic resonance imaging studies will be considered. Our goal was two-fold: (1) to summarize and facilitate the integration of findings in this evolving field; and (2) to provide an illustrative structural and functional brain map of significant BD-associated gene risk variants, which are expected to be linked to brain regions with known alterations in BD.

## **2. Methods**



A systematic literature search of genetic variations and functional and structural magnetic resonance imaging (MRI) studies in BD was conducted. We followed the *Preferred Reporting Items for Systematic Reviews and Meta-Analyses* (PRISMA) statement (Moher et al., 2010), using an *a priori* defined but unpublished protocol.

## **2.1. Search strategy**

The EMBASE, PubMed/MEDLINE and PsycINFO electronic databases were searched from inception up to January 9<sup>th</sup>, 2017. The following search string was used: (bipolar disorder OR mania OR bipolar depression) AND (structural magnetic resonance OR functional magnetic resonance OR fMRI OR BOLD fMRI OR magnetic resonance imaging OR magnetic resonance neuroimaging OR tractography) AND (SNPs OR single nucleotide polymorphism OR haplotypes OR gene expression OR gene OR genetic score OR genetic\* OR methylome OR epigenetic\* OR genome OR transcriptome OR polymorphism OR genetic polymorphism OR genome wide OR genome-wide). In addition, the reference lists of eligible articles were hand searched to identify additional eligible references.

## **2.2. Eligibility Criteria**

The articles included in this review fulfilled the following criteria: (1) human studies with participants at any age with a diagnosis of type I BD (BD-I), type II BD (BD-II), or BD not otherwise specified (BD-NOS) using standard diagnostic criteria (DSM-IV, ICD-10 or Research Diagnostic Criteria regardless of the current mood state (euthymic, manic or depressed); (2) combined investigations of genetic factors and brain imaging protocols (structural or functional). The included articles had to investigate imaging-genetic associations of BD patients that were carriers of high-risk alleles compared to either healthy controls (HC) and/or BD patients who were non-carriers of the

investigated risk alleles. No language restrictions were applied. Studies that reported a sub-analysis of a well-defined sample of participants with BD within a broad mood disorder group were also eligible.

Animal and *post-mortem* studies, case series, literature reviews, conference papers, meeting abstracts or meta-analyses were excluded. Studies which included samples with mixed diagnoses were excluded, unless data for participants with BD were separately provided. Articles that used imaging methods other than structural or functional MRI (e.g., magnetic resonance spectroscopy or positron emission tomography) were also excluded.

### **2.3. Study Selection**

Two investigators (LPP and BPF) independently screened the titles and abstracts of retrieved references for eligibility. Next, the full-texts of the selected references were obtained, and the same authors independently reviewed each article for final inclusion in this systematic review. Disagreements were resolved through consensus. Whenever a consensus could not be achieved, a third author (CAK) made the final decision regarding inclusion. The agreement between the two raters was high (83.7%).

### **2.4. Data Extraction**

Two authors (LPP and BPF) independently extracted the data of selected papers using a standardized spreadsheet. The following variables were recorded: first author, year of publication, sample size, age of participants, % of females, diagnostic criteria for BD, genetic assessment (name of the gene, method, SNP and allele groups), imaging

methods and procedures, the experimental paradigm (in case of fMRI), MRI regions of interest (ROI) and the results of the association between the genetic variants and each ROI. Whenever the sample contained BD patients as part of a broader sample that included other psychiatric diagnoses, only data and associations of the BD group was extracted. The agreement between the two raters was 89.6%.

## **2.5. Data synthesis**

Due to the anticipated heterogeneity and paucity of homogenous studies, meta-analysis of included studies was not feasible. Thus, we synthesized the included studies with a best evidence synthesis. First, we considered structural imaging studies and candidate genes and GWAS relationships. Second, we considered the relationship between functional imaging studies and candidate genes and GWAS studies. We considered evidence to be replicated or consistent when a relationship was evident between 2 studies between a candidate gene/GWAS and a particular structural and/or functional neuroimaging abnormality.

## **3. Results**

### **3.1 Search Results**

The literature search found 873 records, and 9 additional references were found through searching the reference lists of included articles. After the removal of duplicates, 632 unique references were screened. Five hundred and seventy-one references were excluded after title/abstract screening. Of the 61 full-texts assessed, 17 were excluded due to: (1) not an original study ( $k = 2$ ); (2) no data for BD participant was provided ( $k = 9$ ), (3) not investigating samples with BD ( $k = 1$ ), (4) no genetic measure ( $k = 1$ ), (5) using other neuroimaging method not specified in the inclusion criteria ( $k = 2$ ), (6) not investigating genetic-imaging associations in BD ( $k = 1$ ) or (7) article not available ( $k =$

1). Therefore, forty-four genetic-neuroimaging studies met inclusion criteria for this qualitative systematic review. Figure 1 presents the flowchart of study selection. The studies excluded during full-text review and reasons for exclusion are presented in Supplementary Table S1 that accompanies the online version of this article.

<Please insert Figure 1 here>

### **3.2 Overview of included studies**

All included studies are described in Tables 1 and 2 (structural MRI,  $k = 28$ ) and Tables 3 and 4 (fMRI,  $k = 16$ ). Forty studies investigated 26 candidate risk genes for BD and 4 studies used a genome-wide significance analysis. The studies altogether included 2,122 participants with BD [BD group; age =  $38.6 \pm 13.6$  years (mean  $\pm$  SD); 56.6% female] and 2,389 healthy participants (HC group; age =  $35.9 \pm 12.4$  years (mean  $\pm$  SD); 53.0% female). All studies included only adult samples except for three studies that included only pediatric samples (Barzman et al., 2014; Liu et al., 2010; Zeni et al., 2016). Twenty-eight studies investigated structural changes using either VBM or DTI, and 16 studies used functional MRI to investigate changes in brain activity associated. The functional studies were based on several tasks, including emotional processing of faces ( $k = 8$ ), Posner emotional task ( $k = 1$ ), verbal fluency tasks ( $k = 4$ ), and working memory ( $k = 2$ ). Emotional tasks included contrasts of the task-related activity and baseline, and within neutral and affective content. The other tasks compared task-related activity with baseline.

<Please insert Table 1 here>

<Please insert Table 2 here>

<Please insert Table 3 here>

<Please insert Table 4 here>

### 3.3. Structural imaging studies

#### 3.3.1. Candidate Genes

Twenty-six studies investigated associations of 19 candidate genes with structural imaging data (see Table 1 for studies using VBM and Table 2 for studies using DTI). Except for the study by Zeni et al. (2016), all other studies included an adult sample. Eighteen studies investigated structural measures [total/regional brain volumes, cortical thickness and white matter (WM) integrity] using VBM, 6 studies focused on DTI metrics [e.g. fractional anisotropy (FA)] and 2 studies used both methods. The most frequently investigated genes were *BDNF* (5 studies, all VBM), *CACNA1C* (5 studies, 4 VBM and 1 DTI), *ANKK1* (3 studies, 1 VBM and 2 VBM/DTI combined), *5-HTTLPR* (3 studies, 2 VBM and 1 DTI), *ZNF804A* (3 studies, 1 VBM and 2 DTI), and *GSK-3 $\beta$*  (2 studies, 1 VBM and 1 DTI) (Table 1 and Table 2). The remaining genes (*EAAT2*, *DGKH*, *NRG1*, *HAP*, *CNP*, *MOG*, *IL-1B*, *ODZ4*, *SYNE1*, *DAOA*, *GRIN2B*, *SREBF1* and *SREBF2*) were investigated by a single study.

Six studies included only participants with BD (i.e. carriers vs. non-carriers of genetic risk variants) (Benedetti et al., 2013; Benedetti et al., 2015a; Benedetti et al., 2015b; Benedetti et al., 2014; Poletti et al., 2016; Poletti et al., 2014). Three of those studies investigated *5-HTTLPR* (Benedetti et al., 2015a), *GSK-3 $\beta$*  (Benedetti et al., 2013) or *SREBF1/2* (Poletti et al., 2016) using DTI. Benedetti et al. (2015a) found that carriers of the *5-HTTLPR* S (i.e., short) allele had increased radial and mean diffusivity in several brain white matter tracts, including the cingulum gyrus, corpus callosum (body and genu) and corona radiata compared to non-carriers. Significant increases in axial diffusivity measures were observed in carriers of the less active GSK3- $\beta$

rs334558\*C gene-promoter variant in 70 participants with an index bipolar depressive episode across several white matter fiber tracts (Benedetti et al., 2013). Interestingly, lithium treatment (which inhibits GSK-3 $\beta$ ) was also associated with similar changes in axial diffusivity, which points to a better integrity of axon and myelin sheaths (Benedetti et al., 2013). Poletti et al. (2016) found that carriers of the *SREBF2* rs1052717 polymorphism A/A genotype had increased radial diffusivity and reduced FA compared to G carriers in the cingulum, corpus callosum, superior and inferior longitudinal fasciculi, and anterior thalamic radiation. The remaining 3 studies investigated variations in the *5-HTTLPR* (Benedetti et al., 2014), *GSK-3 $\beta$*  (Benedetti et al., 2015b) or *EAAT2* (Poletti et al., 2014) genes using VBM. All three studies did not verify any significant genetic-imaging associations in BD.

The other 19 studies included a HC comparison group. Fourteen of these studies found significant associations of brain structural changes and genetic variants, in both grey and white matter. These included associations of the *BDNF* (k = 4), *5-HTTLPR* (k = 1), *CACNAC1* (k = 1), *DGKH* (k = 1), *NRG1* and *HAP<sub>ICE</sub>* haplotype (k = 1), *IL-1 $\beta$*  (k = 1) with brain volumes using VBM, and also *ANK3* (k = 2), *5-HTTLPR* (k = 1), *GSK-3 $\beta$*  (k = 1) and *GRIN2B* (k = 1) with white matter integrity using DTI. See section 3.5.1 for details.

### 3.3.2. Genome-Wide Association Studies

Two studies used Genome-Wide Association Studies (GWAS) to identify genes associated with BD, and then investigated associations with structural changes using VBM (Bakken et al., 2011; Oertel-Knochel et al., 2015) (Table 1). Oertel-Knochel et al. (2015) investigated 7 SNPs obtained from a GWAS study in SZ (*MIR137*, *CCDC68*, *CNNM2*, *NT5C2*, *MMP16*, *CSMD1* and *PCGEM1*) to identify genetic variants

associated with structural brain changes across the psychosis spectrum. No statistically significant association was observed for the group that included only participants with BD. Bakken et al. (2011) examined associations of 597,198 SNPs with average cortical thickness using the PLINK analytic tool (Purcell et al., 2007) to fit an additive linear model with minor allele counts, sex, age and diagnosis, using a conservative Bonferroni correction for genome-wide significance. No statistically significant imaging-genetic associations were found in the BD group.

### **3.4. Functional Imaging studies**

#### *3.4.1. Candidate Genes*

Fifteen studies investigated associations of variations in 11 candidate genes and blood oxygen level dependent (BOLD) fMRI. The most frequently investigated genes were *CACNA1C* (4 studies) and *ANK3*, *DAAO* and *DISC1* (2 studies each) (Tables 3 and 4). The remaining genes (*TNF*, *G72*, *BclI*, *COMT*, *DOK5*, *5-HTTLPR* and *NRG1*) were investigated in a single study.

Barzman et al. (2014) investigated a small sample of pediatric BD patients, and found that the expression of 11 TNF-related genes in peripheral blood mononuclear cells of participants with BD significantly correlated with activation of the amygdala or anterior cingulate gyrus during the affective Posner task.

All other studies included only adults, and included a HC group for comparison. Eight of these studies found significant gene  $\times$  brain activity associations in BD patients. The *CACNA1C* gene was associated with increased amygdala activation in the face recognition paradigm ( $k = 2$ ). The *ANK3* gene was associated with increased activity in the cingulate cortex during a working memory task ( $k = 1$ ). Both the *ANK3* and *CACNA1C* genes were associated with reduced activation of the vIPFC during the

emotional facial processing task ( $k = 1$ ). The DISC1 gene was associated with decreased activation of the IPL and left CG during a verbal initiation and sentence completion task ( $k = 1$ ). Also in a verbal fluency paradigm, the DAOA ( $k = 1$ ) genotype was associated with a greater deactivation of the left precuneus in BD patients, while the NRG1 genotype was associated with increased activation of the right posterior OFC. Finally, the 5-HTTLPR was associated with lower ventral anterior CG activity during emotional processing of faces ( $k = 1$ ). See section 3.5.2 for details.

#### *3.4.2. Genome-Wide Association Studies*

Liu et al. (2010) investigated a sample of adolescents with BD and HCs of similar age. These authors performed a GWAS, and found that the rs2023454 SNP of the *DOK5* gene was associated with right amygdala activation under contrast to hostility faces although no significant differences between and within BD and HC samples were observed. Dima et al. (2016) calculated a polygenic risk score (PGR) from genes that were associated with BD in a GWAS. Although the PGR was associated with changes in brain activity during a facial processing task and a working memory task in both the BD and HC groups, no statistically significant differences emerged between the two groups or within the BD group as a function of the PGR.

### **3.5. Significant genetic-neuroimaging associations in BD**

The statistically significant associations reported across genetic-neuroimaging studies using candidate genes are shown in Table 5. In addition, a brief synopsis of possible biological functions of gene products is provided.

#### *3.5.1. Structural VBM and DTI studies*



Statistically significant structural neuroimaging alterations in BD patients were associated with genetic variations of the *BDNF*, *5-HTTLPR*, *CACNA1C*, *DGKH*, *NRG1*, *IL-1 $\beta$* , *ANK3*, and *GRIN2B* genes (Table 5). Nevertheless, there was a lack of replicated evidence.

<Please insert Table 5 here>

Two studies provided evidence that BD patients who carry the Met allele of the *BDNF* gene may present several structural alterations encompassing several brain areas namely the left and right hippocampus (Cao et al., 2016; Chepenik et al., 2009). A four-year prospective study found that individuals BD participants who were carriers of one or more *BDNF* Met alleles had significantly greater losses in gyrification indexes, an effect that correlated with gray matter loss in the left hemisphere (Mirakhur et al., 2009). Matsuo et al. (2009) observed smaller bilateral anterior cingulate gyrus volumes in BD patients with Val/Met compared to those with Val/Val *BDNF* genotypes, while in both the BD and HC groups participants with the Val/Met *BDNF* genotype had smaller left and right gray matter volumes of the dorsolateral prefrontal cortex.

Increased volumes of the left amygdala were observed in carriers of the S allele of the *5-HTTLPR* gene both in BD and HC groups (Scherk et al., 2009a).

Genetic variations in the *CACNA1C* genes were not associated with significant structural changes in three VBM studies (Soeiro-de-Souza et al., 2012; Tesli et al., 2013; Wolf et al., 2014), whereas Perrier et al. (2011a) found that euthymic BD patients carrying the *CACNA1C* rs1006737 risk allele had a smaller volume of the left putamen compared HCs.

A significantly increased volume of the left amygdala was associated with the *DGKH* haplotype (rs994856/rs9525580/rs9525584 GAT) in 30 euthymic patients with type I BD but not in HCs (Kittel-Schneider et al., 2015). The risk genotype (TT) of the

*NRG1* SNP8NRG221533 was associated with reduced white matter volumes in the fornix, cingulum and para-hippocampal gyrus in a type I BD sample (Cannon et al., 2012). In the same study, BD participants carrying one or two copies of the *HAP<sub>ICE</sub>* haplotypes of the *NRG1* gene had greater white matter volume than those carrying none in the fornix, caudate and cingulum (Cannon et al., 2012). Papiol et al. (2008) found that a -511C/T SNP (rs16944) of the IL-1 $\beta$  gene was associated with whole-brain and left dlPFC gray matter deficits in a sample of 20 participants with BD in a VBM study. Two studies found that distinct variations of the *ANK3* gene were associated with DTI findings (reduced FA) compatible with widespread white matter deficits in several brain regions, such as the forceps minor, the uncinate fasciculus, the anterior cingulate gyrus, the dorsolateral frontal cortex, the left temporoparietal WM, and in posterior dorsomedial WM (Lippard et al., 2016; Ota et al., 2016). Finally, compared to the G allele of the *GRIN2B* gene, brain FA values were significantly lower in BD patients with risk T allele in left and right frontal regions, left parietal region, left and right occipital regions and the left cingulate gyrus (Kuswanto et al., 2013).

### 3.5.2. *fMRI studies*

Functional neuroimaging alterations were associated with genetic variations in the *CACNA1C*, *ANK3*, *DISC1*, *TNF*, *DAOA*, *5-HTTLPR* and *NRG1* genes (Table 5).

The most frequent regions with functional alterations significantly associated with genetic variations in BD were: (1) the right anterior CG (ACG), where variation in the *ANK3* and *TNF* genes were associated with greater activation in working memory tasks (*ANK3*) or Posner task (*TNF*), whereas the *5-HTTLPR* S allele was significantly associated with lower activation during an emotional processing task; (2) the left ACG, where polymorphisms in the *TNF* gene were associated with increased activation and the *5-HTTLPR* S allele with decreased activation during emotional processing tasks; (3)

left amygdala, where variation in the *CACNA1C* and *TNF* genes were associated with greater activation during emotional processing tasks; (4) left CG, where polymorphisms in *DISC1* were associated with lower activation during a verbal fluency task; (5) left para-hippocampal gyrus, where *ANK3* polymorphisms were associated with greater activation; and during a working memory task; and (6) the left vIPFC, where variations in both *CACNA1C* and *5-HTTLPR* genes were associated with greater activation during emotional processing tasks.

The fMRI paradigms that were most frequently used across functional neuroimaging studies with statistically significant genetic-imaging findings were emotional faces task (8 studies) and the verbal fluency test (2 studies). Three studies found associations of the *CACNA1C* risk allele A with an increase in activation of either left or right amygdala (Jogia et al., 2011; Tesli et al., 2013), and a hypoactivation of the vIPFC (Dima et al., 2013; Jogia et al., 2011), with one study reported both alterations (Jogia et al., 2011) in the emotional faces task. The remaining studies found decreased activation of the vIPFC in association with the *ANK3* rs10994336 polymorphism risk allele T (Delvecchio et al., 2015) or a decreased activation of the ventral anterior cingulate gyrus related to the *5HTTLPR* S risk allele (Shah et al., 2009). Studies that employed the verbal fluency test were inconsistent regarding both genetic variations and activated ROIs (Mechelli et al., 2012; Mechelli et al., 2008).

### *3.5.3. Illustrative brain map of significant replicated gene-neuroimaging findings*

Figure 2 summarized replicated gene-imaging findings in BD patients in comparison to healthy controls. A difference was considered statistically significant ( $p < 0.05$ ) only if the neuroimaging findings of BD patients carrying the risk allele were different from the HCs or BD subjects not carrying the risk allele (i.e., a gene x diagnosis interaction).

Two fMRI studies found that individuals with BD carrying the A variant of the *CACNA1C* Rs1006737 polymorphism had decreased activity in the right dorsal ventrolateral prefrontal cortex during the emotional faces paradigm. However, samples across those two investigations appeared to overlap (Dima et al., 2013; Jogia et al., 2011), and thus this association was not regarded as a true replication. Furthermore, two VBM studies found that subjects BD who were carriers of the Met allele of the BDNF Val66Met polymorphism had decreased volumes of the left and right hippocampi (Cao et al., 2016; Chepenik et al., 2009).

<Please insert Figure 2 here>

### **3.6. Methodological considerations**

The minority of included studies enrolled only euthymic BD participants ( $k = 10$ ; 22.7%), while the mood status of participants with BD was clearly described in 21 (47.7%) studies. Twenty studies (45.5%) controlled results for the effects of medication or otherwise included only drug-free BD participants, while most included studies controlled findings for multiple comparisons ( $k = 33$ ; 75.0%). A healthy control group was included in 36 (81.8%) studies. The median (IQR) sample sizes for VBM, DTI and fMRI studies were 80 (72-84), 153.5 (87.25-172) and 80.5 (69.5-87.25). A whole-brain analysis was conducted in 15 (34.0%) studies, while 18 (40.9%) studies performed only a priori defined ROI-based analyses, and 4 (9.1%) studies carried out both types of analyses. Twenty-eight studies used a 1.5T magnetic field, 13 studies used 3.0T, 1 study 4.0T and 2 did not specify the magnetic field of the scanner.

## **4. Discussion**

The aim of this systematic review was to assess the extant literature reporting ‘imaging genetics’ findings in BD. We included both structural and functional MRI studies. The most frequently reported genes (at least 2 studies) with statistically significant neuroimaging alterations in BD patients were *CACNA1C*, *ANK3*, *BDNF*, *5-HTTLPR*, *NRG1* and *DAOA*. Of those genes, loci close to the *CACNA1C* and *ANK3* genes have reached genome-wide significant associations with BD, and associations were replicated in at least one independent dataset (Goes, 2016). To our knowledge this effort represents the largest evidence-based synthesis to date of this field.

Our findings suggest that the effects of genetic variants on intermediate neuroimaging phenotypes could be independent (i.e., pleiotropic) of effects on the clinical phenotype per se (Gottesman and Gould, 2003), which is consistent with the view that an endophenotypic approach may aid in the search of biological pathways underpinning heterogeneous mental disorders like BD (Miskowiak et al., 2016). The findings reviewed herein may at least partly explain that although at the population level BD is associated with a significant degree of both “cold” and “hot” (i.e., emotion-laden) (Miskowiak and Carvalho, 2014; Roiser et al., 2009) cognitive deficits, recent meta-analyses point to a significant degree of heterogeneity (Bora and Pantelis, 2016; Bortolato et al., 2015; Bourne et al., 2013). Nevertheless, a certain degree of uncertainty lies on the precise pathways which could be influenced by those gene products with relevance to the underlying neurobiology of subsets of individuals with BD. Furthermore, one cannot exclude the possibility that those risk genetic variants are not inherently causal, but instead may be passed in linkage disequilibrium with causative ones. We observed that although several candidate gene studies reported significant associations with structural/functional neuroimaging findings, whilst the few studies that followed a GWAS methodology did not report statistically significant findings

(Bakken et al., 2011; Liu et al., 2010; Oertel-Knochel et al., 2015). Evidence indicates that the literature on structural and functional neuroimaging studies could be limited by an excess of significance bias (i.e., there is an excess of statistically significant findings), which may undermine the reproducibility of the field as a whole (Fusar-Poli et al., 2014; Ioannidis et al., 2014). In addition, a selective reporting of outcomes (i.e., only those genes with statistically significant findings are reported) could result in a type I error (Ioannidis et al., 2014). Moreover, sample sizes varied across studies, and due to the few studies available it is difficult to estimate the statistical power of individual studies. This aspect may also undermine the reproducibility of gene-imaging studies as discussed in detail elsewhere (Carter et al., 2016). Therefore, we focused our discussion on candidate genes with at least two statistically significant findings. Furthermore, we contextualized the main findings of our review with data derived from the preclinical and neuropsychological literature.

#### *4.1. The CACNA1C gene*

The *CACNA1C* gene encodes the L-type voltage-dependent calcium channel 1C subunit, and at least two GWAS have implicated its rs1006737 SNP as a risk variant associated with BD (Sklar et al., 2008). This association has been consistently replicated since then (Goes, 2016). Notwithstanding Perrier et al. (2011a) observed a significantly reduced volume of the left putamen in a sample of BD patients carrying the rs1006737 SNP risk allele compared to HCs. However, two subsequent VBM studies failed to replicate those findings (Soeiro-de-Souza et al., 2012; Wolf et al., 2014).

Significant within-group differences were observed in BD who were carriers of the risk allele of the *CACNA1C* gene after recognition of negative/fearful faces (compared to neutral faces) in the facial affect recognition task. For example, carriers of the risk allele had a higher activation of the left amygdala in one study (Tesli et al.,

2013), while another study found higher activation of the amygdala bilaterally in the same task (Jogia et al., 2011). Furthermore, two studies found an hyperactivation of the left ventrolateral prefrontal cortex in the same experimental paradigm (Dima et al., 2013; Jogia et al., 2011). Nevertheless, another study did not report significant functional brain abnormalities related to this risk allele in the facial emotion recognition task (Radua et al., 2013). Methodological differences across studies may explain those discrepant findings. For example, Radua et al. (2013) did not explicitly exclude BD participants with co-occurring somatic and mental disorders. Furthermore, there was an overlap in samples included in the studies carried out by Jogia et al. (2011) and Dima et al. (2013). Ou et al. (2015) postulated that the lack of significant differences in neuroimaging studies between participants with BD and HCs as a function of the *CACNA1C* risk allele could be due to differences in the prevalence of cardiovascular risk factors. Therefore, differences in eligibility criteria across studies may at least in part explain contrasting results across studies. Nevertheless, those functional brain abnormalities as a function of the presence of the AA/AG *CACNA1C* risk alleles are consistent with the neuropsychological literature. Hence although less unanimous than the neuroimaging literature, BD patients who carry those risk alleles may score poorer on speed of processing and digit span tests compared to non-carriers (consistent with a lower efficiency of the left ventrolateral prefrontal cortex). In addition, carriers of the risk allele may display impaired recognition of emotional faces (disgust, sadness, happiness, and anger) [see Ou et al. (2015) for a review].

#### 4.2. *The ANK3 gene*

The *ANK3* gene codes for Ankyrin G, a protein which may be involved in the stabilization and localization of ion channels and cell adhesion molecules to nodes of Ranvier and initial segments of axon (Gasser et al., 2012). Furthermore, an elegant

preclinical study found a lack of voltage-gated sodium channels in GABAergic parvalbumin interneurons in mice deficient of exon 1b of the *ANK3* (Lopez et al., 2016). Consistently, mice exhibited an *ANK3* gene dose-dependent phenotype characterized by manic-like behavior, epilepsy, and sudden death (Lopez et al., 2016). In addition, Ankyrin G has been implicated in neurodevelopment, and in the onset of myelination (Ching et al., 1999), and also in the regulation of neurogenesis (Durak et al., 2015; Leussis et al., 2012).

Replicated findings indicate that different risk alleles of the *ANK3* gene may impact white matter structure in BD. For example, BD patients who were carriers of the risk allele rs10761482 SNP had decreased FA in the forceps minor (Ota et al., 2016), while BD patients who were carriers of the risk allele of the rs9804190 had decreased FA in the uncinate fasciculus and the cingulate gyrus bilaterally among other regions compared to CC homozygotes (Lippard et al., 2016). These findings are consistent with a recent meta-analysis of DTI studies in BD which evidences widespread white matter in this illness compared to controls (Nortje et al., 2013). Greater widespread abnormalities in individuals with BD carrying risk variants of the *ANK3* gene also provide support for a putative role of Ankyrin G in myelination. In addition, risk alleles of this gene could lead to accelerated brain aging in BD (Rizzo et al., 2014). Furthermore, the risk C-allele of rs10761482 SNP was significantly associated with worse performance on verbal comprehension, logical memory and processing speed in BD patients in one study (Hori et al., 2014), while another study found that the risk allele of the rs10994336 SNP was associated with reduced sensitivity in target detection and increased errors of commission during sustained attention in both patients with BD and HCs (Ruberto et al., 2011). Altogether, these data suggest that different risk alleles of the *ANK3* gene could have a deleterious effect on WM structure in BD, which could



be related to neurocognitive deficits. This hypothesis is further supported by a study that found that the risk allele of the rs10994336 SNP was associated with hyperactivation of the right anterior cingulate cortex and left posterior cingulate cortex in patients with BD compared to HCs in the N-back test, which measures executive function (Delvecchio et al., 2015).

#### 4.3. *The BDNF gene*

The brain-derived neurotrophic factor (*BDNF*) gene is located on chromosome 11p14.1. The BDNF protein is a member of the neurotrophin superfamily, which supports neuronal survival, neural differentiation during development, and has been implicated in the regulation of activity dependent-synaptic plasticity in mature neurons (Duman and Monteggia, 2006; Hempstead, 2015). This neurotrophin is abundantly expressed in the hippocampus (Duman and Monteggia, 2006). The BDNF rs6265 SNP has been frequently investigated and an alteration at nucleotide 196 (G/A) which produces a Val66Met substitution (Notaras et al., 2015a). This SNP may result in a diminished cellular trafficking and packaging of the mature BDNF protein into the secretory vesicles, thus reducing depolarization-induced release of this neurotrophin (Notaras et al., 2015a). Furthermore, carriers of this risk SNP could produce the Met BDNF prodomain in larger amounts, with may have opposing effects (i.e., a negative impact in neuron architecture remodeling) via an activation of the p75 and sortilin-related VPS10 domain containing receptor 2 (SorCS2) receptors (Hempstead, 2015). Consistently, this systematic review found replicated evidence that carriers of the BDNF met allele exhibit smaller hippocampal volumes (Cao et al., 2016; Chepenik et al., 2009). In one study BD carriers of this risk allele presented smaller hippocampal volumes compared to HCs (Cao et al., 2016), whereas in carriers of the Met allele had smaller hippocampus regardless of diagnostic group (i.e., BD or HC) (Chepenik et al., 2009). Zeni et al.

(2016) studied a sample of pediatric patients with BD, and found that the Met allele of the BDNF gene had no influence on hippocampal volumes. A potential explanation for this finding is that this SNP could influence hippocampal volume over time as suggested by a previous study (McIntosh et al., 2007). A previous meta-analysis found that neuropsychiatric patients with either the Val/Val genotype or Met-carriers had significantly smaller hippocampal volumes compared to HCs with the same genotypes (Harrisberger et al., 2015). Therefore, it is possible that the Met risk allele could mediate within group differences in BD samples, but not differences between participants with BD and HCs.

A recent meta-analysis suggests that the *BDNF* Val66Met SNP is not associated with BD (Gonzalez-Castro et al., 2015), although this association could be significant in European populations (Li et al., 2016). This highlights that the effects of the Met allele of the *BDNF* gene could be more readily demonstrated at the neuroimaging or neuropsychological level than at the diagnostic level. Hence, several studies suggest that individuals with BD carrying the Met allele could have worse cognitive function in several domains including memory (Cao et al., 2016; Rybakowski et al., 2003; Rybakowski et al., 2006; Tramontina et al., 2009), although this association has not been unanimously demonstrated across studies (Rolstad et al., 2016; Rosa et al., 2014). These discrepancies may be related to the influence of concomitant medication (Grande et al., 2014). Furthermore, the involvement of BDNF in the pathophysiology of BD is supported by a recent meta-analysis which found that peripheral levels of this protein could be a biomarker of illness activity (Fernandes et al., 2015). Finally, preclinical evidence points to a role for BDNF in BD (de Souza Gomes et al., 2015; Macedo et al., 2012).

#### 4.4. The 5-HTTLPR gene

A functional polymorphism (*5-HTTLPR*) in the promoter of serotonin transporter gene (*SLC6A4*) has been described in 1996 (Lesch et al., 1996). Since then, the impact of this polymorphism in a range of mental disorders and intermediate phenotypes have been a focus of substantial research efforts [see Jonassen and Landro (2014) for a review]. In addition, evidence suggests that methylation of the serotonin transporter gene may provide an epigenetic marker of exposure to life adversities (Provenzi et al., 2016). Notwithstanding the *5-HTTLPR* polymorphism does not seem to affect the methylation status of the *SLC6A4* gene, preliminary evidence suggests a possible interaction of methylation status and the short (S) allele in the development of stress-related mood disorders (Olsson et al., 2010).

Benedetti et al. (2014) observed that the S allele of the *5-HTTLPR* mediated the effect of early life stress on gray matter volumes in the right prefrontal cortex in a sample with BD. Furthermore, the S allele was also associated with higher right amygdala volumes in both patients with BD and HCs. Notwithstanding a previous meta-analysis suggests that the S allele could lead to amygdala hyperactivation in emotional paradigms (Murphy et al., 2013), our systematic review did not find a study to replicate this finding in BD. Nevertheless, one study found that S carriers had lower ventral anterior cingulate cortex activation compared to L/L participants during processing of happy and fear faces; this effect was evident in both the HC and BD groups (Shah et al., 2009). Clearly effects of the ‘S’ *5-HTTLPR* on intermediate phenotypes in BD deserve further investigation.

#### 4.5. *The NRG1 gene*

Evidence indicates that neuregulin 1 and its cognate receptor ErbB4 play significant roles in the regulation of synaptic transmission, myelin formation, and neuronal and glial cell survival (Mei and Nave, 2014). Although variations in *NRG1* gene were initially

associated with schizophrenia [see Mostaid et al. (2016) for a review], subsequent studies pointed to a possible association with BD (Cao et al., 2014; Georgieva et al., 2008; Green et al., 2005; Gutierrez-Fernandez et al., 2014), notwithstanding this findings has not been supported thus far by GWAS (Goes, 2016). In keeping with this view, a study found aberrant cleavage of the neuregulin 1 in the *post mortem* hippocampus of individuals with BD (Marballi et al., 2012). We found evidence that a risk *NRG1* SNP (SNP8NRG221533) and its HAP<sub>ICE</sub> haplotype was associated with greater white matter in the fornix, cingulum, para-hippocampal gyrus, and the corpus callosum (Cannon et al., 2012). In addition, a functional neuroimaging study found that individuals with BD carrying the high-risk SNP (rs35753505) of the *NRG1* gene displayed hyperactivation of the right posterior orbitofrontal cortex compared to non-carriers (Mechelli et al., 2012). Clearly the impact of high-risk variants of the *NRG1* gene on structural and functional brain abnormalities in individuals with BD require further study.

#### 4.8. Limitations

The findings of this systematic review should be interpreted within its limitations. First, the methodological quality of included studies varied. For example, the mood state of participants with BD varied across studies. In addition, some studies did not include a HC group, while few studies did not control results for multiple comparisons. Second, several confounding variables should be considered (e.g., differences in length of illness, number of previous affective episodes, and exposure to mood stabilizing medications). For example, it has been suggested that hippocampal volumes in BD may vary as a function of the number of affective episodes in a subset of patients with neuroprogressive forms of the illness (Cao et al., 2017; Lim et al., 2013). Furthermore, the large majority of studies included in this systematic review enrolled adult samples.

For example, a meta-analysis indicates that hyperactivation of the amygdala across emotional face recognition fMRI studies is more evident in BD-youths than among BD-adults (Wegbreit et al., 2014). Third, although our findings indicate that the effects of genetic variants in the risk of BD may be more readily reflected as at the brain structure/function level than in the disease per se, this notion has been challenged by some experts. For example, Flint and Munafò (2007) provides meta-analytic evidence that the effect sizes of illness-related genetic variants on intermediate phenotypes may not necessarily be larger than the ones observed for the illness phenotype. Fourth, we included structural and functional MRI studies, but not other imaging tools (e.g., positron emission tomography). Fifth, although we found promising replicated findings, several significant associations deserve replication. For example, converging evidence from both preclinical and GWAS studies have implicated ANK3 as a putative risk gene for BD, while recent gene imaging studies offered promising initial results (Lippard et al., 2016). In addition, the reproducibility of this field deserves careful examination (Carter et al., 2016). Sixth, the use of pre-defined ROI-based analyses could bias some of the converging findings of this review. For example, the study by Chepenik et al. (2009) in which the Met allele of the BDNF Val66Met polymorphism was associated with bilateral hippocampi reduction in individuals with BD restricted their analyses to this brain structure.

#### *4.9. Implications*

This systematic review open several research implications. It has been increasingly recognized that neurobiological abnormalities span conventional diagnostic categories in psychiatry. This fact motivated the NIMH to launch the Research Domain Criteria (RDoC) initiative (Cuthbert and Insel, 2013), with an attempt to provide a complimentary research classification system for mental disorders built upon

dimensions of neurobiology and observable behavior, and moving towards precision psychiatry (Fernandes et al., 2017; Vieta, 2015). Consistent with this assumption several genetic risk variants seem to overlap across major mental disorders (Gatt et al., 2015). Furthermore, a recent study investigated a large panel of brain-based biomarkers and included participants across the psychotic spectrum (schizophrenia, schizoaffective disorder, and BD), and found three distinct psychotic biotypes that did not respect diagnostic categories (Clementz et al., 2016). Consistently, the only replicated finding observed in this review is also apparent in similar studies involving schizophrenia samples. For example, the Val66Met BDNF polymorphism has also been associated with reduced hippocampi volume in schizophrenia (Notaras et al., 2015b), while a recent systematic review found that several putative risk genes for both BD and schizophrenia may influence brain structure and function in healthy control samples (Gurung and Prata, 2015). Therefore, future efforts to replicate the findings of this systematic review could include participants with different diagnostic categories, and a better control of potential confounding variables (e.g. concomitant medication and substance use. In addition, the *a priori* publication of research protocols in the field of ‘imaging genetics’ could improve its reproducibility and reduce the risk of selective outcome reporting (Carter et al., 2016).

## **5. Conclusion**

This review synthesis indicates that the ‘gene-imaging’ research paradigm may aid in the identification of intermediate phenotypes, and therefore could provide more consistent biological mechanistic insights for BD. Variants in the *CACNA1C*, *ANKK1*, and *BDNF* genes yielded the most consistent findings thus far, by applying neuroimaging paradigms to GWAS-emerging candidate genes. Future research efforts should include samples with different diagnostic categories, and the development of

collaborative consortia with a priori published protocols could enhance the impact of future efforts. However, highly robust findings are unlikely if the only source of candidate genes are GWAS and gene polymorphisms, and neuroimaging studies are particularly difficult in bipolar patients because of the influence of complex medication regimes.

### **Role of funding source**

This study received no funding.

### **Contributors**

LPP, CAK and AFC designed the protocol and searched the literature. LPP and BPF screened the studies and extracted the data. LPP, CAK and AFC analyzed the data. LPP, CAK, RTS and AFC wrote the first draft of the manuscript. MS, BPF, MF, RMV, KWM, EV, NV and BS contributed to the interpretation and discussion of the findings, and to the writing of the manuscript. All authors have read and approved the final version of this manuscript for submission. All authors have participated sufficiently in the work to take responsibility for its content.

### **Conflict of interest**

The authors declare no conflicts of interest that could influence this work.

### **Acknowledgements**

CAK is supported by a postdoctoral fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brazil). The Lundbeckfonden and the Weimann Foundation provide support for half of KWM's salary as senior research psychologist at

Copenhagen Affective Disorders Research Centre and University of Copenhagen, Denmark, which enables her to do fulltime research until 2020. AFC is the recipient of a research fellowship award from the Conselho de Desenvolvimento Científico e Tecnológico (CNPq; Brazil).

## References

- Bakken, T.E., Bloss, C.S., Roddey, J., Joyner, A.H., Rimol, L.M., Djurovic, S., Melle, I., Sundet, K., Agartz, I., Andreassen, O.A., Dale, A.M., Schork, N.J., 2011. Association of genetic variants on 15q12 with cortical thickness and cognition in schizophrenia. *Archives of general psychiatry* 68, 781-790.
- Barzman, D., Eliassen, J., McNamara, R., Abonia, P., Mossman, D., Durling, M., Adler, C., DelBello, M., Lin, P.I., 2014. Correlations of inflammatory gene pathways, corticolimbic functional activities, and aggression in pediatric bipolar disorder: a preliminary study. *Psychiatry research* 224, 107-111.
- Benedetti, F., Bollettini, I., Barberi, I., Radaelli, D., Poletti, S., Locatelli, C., Pirovano, A., Lorenzi, C., Falini, A., Colombo, C., Smeraldi, E., 2013. Lithium and GSK3-beta promoter gene variants influence white matter microstructure in bipolar disorder. *Neuropsychopharmacology* 38, 313-327.
- Benedetti, F., Bollettini, I., Poletti, S., Locatelli, C., Lorenzi, C., Pirovano, A., Smeraldi, E., Colombo, C., 2015a. White matter microstructure in bipolar disorder is influenced by the serotonin transporter gene polymorphism 5-HTTLPR. *Genes, brain, and behavior* 14, 238-250.
- Benedetti, F., Poletti, S., Radaelli, D., Locatelli, C., Pirovano, A., Lorenzi, C., Vai, B., Bollettini, I., Falini, A., Smeraldi, E., Colombo, C., 2015b. Lithium and GSK-3beta promoter gene variants influence cortical gray matter volumes in bipolar disorder. *Psychopharmacology* 232, 1325-1336.
- Benedetti, F., Riccaboni, R., Poletti, S., Radaelli, D., Locatelli, C., Lorenzi, C., Pirovano, A., Smeraldi, E., Colombo, C., 2014. The serotonin transporter genotype modulates the relationship between early stress and adult suicidality in bipolar disorder. *Bipolar disorders* 16, 857-866.
- Bergmann, O., Haukvik, U.K., Brown, A.A., Rimol, L.M., Hartberg, C.B., Athanasiu, L., Melle, I., Djurovic, S., Andreassen, O.A., Dale, A.M., Agartz, I., 2013. ZNF804A and cortical thickness in schizophrenia and bipolar disorder. *Psychiatry Research - Neuroimaging* 212, 154-157.
- Bigos, K.L., Weinberger, D.R., 2010. Imaging genetics--days of future past. *NeuroImage* 53, 804-809.
- Bora, E., Pantelis, C., 2016. Social cognition in schizophrenia in comparison to bipolar disorder: A meta-analysis. *Schizophrenia research* 175, 72-78.



- Bortolato, B., Miskowiak, K.W., Kohler, C.A., Vieta, E., Carvalho, A.F., 2015. Cognitive dysfunction in bipolar disorder and schizophrenia: a systematic review of meta-analyses. *Neuropsychiatric disease and treatment* 11, 3111-3125.
- Bourne, C., Aydemir, O., Balanza-Martinez, V., Bora, E., Brissos, S., Cavanagh, J.T., Clark, L., Cubukcuoglu, Z., Dias, V.V., Dittmann, S., Ferrier, I.N., Fleck, D.E., Frangou, S., Gallagher, P., Jones, L., Kieseppa, T., Martinez-Aran, A., Melle, I., Moore, P.B., Mur, M., Pfennig, A., Raust, A., Senturk, V., Simonsen, C., Smith, D.J., Bio, D.S., Soeiro-de-Souza, M.G., Stoddart, S.D., Sundet, K., Szoke, A., Thompson, J.M., Torrent, C., Zalla, T., Craddock, N., Andreassen, O.A., Leboyer, M., Vieta, E., Bauer, M., Worhunsky, P.D., Tzagarakis, C., Rogers, R.D., Geddes, J.R., Goodwin, G.M., 2013. Neuropsychological testing of cognitive impairment in euthymic bipolar disorder: an individual patient data meta-analysis. *Acta psychiatrica Scandinavica* 128, 149-162.
- Callicott, J.H., Straub, R.E., Pezawas, L., Egan, M.F., Mattay, V.S., Hariri, A.R., Verchinski, B.A., Meyer-Lindenberg, A., Balkissoon, R., Kolachana, B., Goldberg, T.E., Weinberger, D.R., 2005. Variation in DISC1 affects hippocampal structure and function and increases risk for schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America* 102, 8627-8632.
- Cannon, D.M., Walshe, M., Dempster, E., Collier, D.A., Marshall, N., Bramon, E., Murray, R.M., McDonald, C., 2012. The association of white matter volume in psychotic disorders with genotypic variation in NRG1, MOG and CNP: a voxel-based analysis in affected individuals and their unaffected relatives. *Translational psychiatry* 2, e167.
- Cao, B., Bauer, I.E., Sharma, A.N., Mwangi, B., Frazier, T., Lavagnino, L., Zunta-Soares, G.B., Walss-Bass, C., Glahn, D.C., Kapczinski, F., Nielsen, D.A., Soares, J.C., 2016. Reduced hippocampus volume and memory performance in bipolar disorder patients carrying the BDNF val66met met allele. *Journal of affective disorders* 198, 198-205.
- Cao, B., Passos, I.C., Mwangi, B., Amaral-Silva, H., Tannous, J., 2017. Hippocampal subfield volumes in mood disorders.
- Cao, L., Deng, W., Guan, L., Yang, Z., Lin, Y., Ma, X., Li, X., Liu, Y., Ye, B., Lao, G., Chen, Y., Liang, H., Wu, Y., Ou, Y., Huang, W., Liu, W., Wang, Q., Wang, Y., Zhao, L., Li, T., Hu, X., 2014. Association of the 3' region of the neuregulin 1 gene with bipolar I disorder in the Chinese Han population. *Journal of affective disorders* 162, 81-88.
- Carter, C.S., Bearden, C.E., Bullmore, E.T., Geschwind, D.H., Glahn, D.C., Gur, R.E., Meyer-Lindenberg, A., Weinberger, D.R., 2016. Enhancing the Informativeness and Replicability of Imaging Genomics Studies. *Biological psychiatry*.
- Chakirova, G., Whalley, H.C., Thomson, P.A., Hennah, W., Moorhead, T.W., Welch, K.A., Giles, S., Hall, J., Johnstone, E.C., Lawrie, S.M., Porteous, D.J., Brown, V.J., McIntosh, A.M., 2011. The effects of DISC1 risk variants on brain activation in controls, patients with bipolar disorder and patients with schizophrenia. *Psychiatry research* 192, 20-28.
- Chepenik, L.G., Fredericks, C., Papademetris, X., Spencer, L., Lacadie, C., Wang, F., Pittman, B., Duncan, J.S., Staib, L.H., Duman, R.S., Gelernter, J., Blumberg, H.P., 2009. Effects of the brain-derived neurotrophic growth factor val66met variation on hippocampus morphology in bipolar disorder. *Neuropsychopharmacology* 34, 944-951.

- Ching, W., Zanazzi, G., Levinson, S.R., Salzer, J.L., 1999. Clustering of neuronal sodium channels requires contact with myelinating Schwann cells. *Journal of neurocytology* 28, 295-301.
- Clementz, B.A., Sweeney, J.A., Hamm, J.P., Ivleva, E.I., Ethridge, L.E., Pearlson, G.D., Keshavan, M.S., Tamminga, C.A., 2016. Identification of Distinct Psychosis Biotypes Using Brain-Based Biomarkers. *Am J Psychiatry* 173, 373-384.
- Craddock, N., Sklar, P., 2013. Genetics of bipolar disorder. *Lancet (London, England)* 381, 1654-1662.
- Cuthbert, B.N., Insel, T.R., 2013. Toward the future of psychiatric diagnosis: the seven pillars of RDoC. *BMC medicine* 11, 126.
- de Souza Gomes, J.A., de Souza, G.C., Berk, M., Cavalcante, L.M., de Sousa, F.C., Budni, J., de Lucena, D.F., Quevedo, J., Carvalho, A.F., Macedo, D., 2015. Antimanic-like activity of candesartan in mice: Possible involvement of antioxidant, anti-inflammatory and neurotrophic mechanisms. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* 25, 2086-2097.
- Delvecchio, G., Dima, D., Frangou, S., 2015. The effect of ANK3 bipolar-risk polymorphisms on the working memory circuitry differs between loci and according to risk-status for bipolar disorder. *American Journal of Medical Genetics, Part B: Neuropsychiatric Genetics* 168, 188-196.
- Dima, D., Breen, G., 2015. Polygenic risk scores in imaging genetics: Usefulness and applications. *Journal of psychopharmacology (Oxford, England)* 29, 867-871.
- Dima, D., de Jong, S., Breen, G., Frangou, S., 2016. The polygenic risk for bipolar disorder influences brain regional function relating to visual and default state processing of emotional information. *NeuroImage. Clinical* 12, 838-844.
- Dima, D., Jogia, J., Collier, D., Vassos, E., Burdick, K.E., Frangou, S., 2013. Independent modulation of engagement and connectivity of the facial network during affect processing by CACNA1C and ANK3 risk genes for bipolar disorder. *JAMA psychiatry* 70, 1303-1311.
- Duman, R.S., Monteggia, L.M., 2006. A neurotrophic model for stress-related mood disorders. *Biological psychiatry* 59, 1116-1127.
- Durak, O., de Anda, F.C., Singh, K.K., Leussis, M.P., Petryshen, T.L., Sklar, P., Tsai, L.H., 2015. Ankyrin-G regulates neurogenesis and Wnt signaling by altering the subcellular localization of beta-catenin. *Molecular psychiatry* 20, 388-397.
- Fernandes, B.S., Molendijk, M.L., Kohler, C.A., Soares, J.C., Leite, C.M., Machado-Vieira, R., Ribeiro, T.L., Silva, J.C., Sales, P.M., Quevedo, J., Oertel-Knochel, V., Vieta, E., Gonzalez-Pinto, A., Berk, M., Carvalho, A.F., 2015. Peripheral brain-derived neurotrophic factor (BDNF) as a biomarker in bipolar disorder: a meta-analysis of 52 studies. *BMC medicine* 13, 289.
- Fernandes, B.S., Williams, L.M., Steiner, J., Leboyer, M., Carvalho, A.F., Berk, M., 2017. The new field of 'precision psychiatry'. *BMC medicine* 15, 80.
- Flint, J., Munafò, M.R., 2007. The endophenotype concept in psychiatric genetics. *Psychological medicine* 37, 163-180.
- Fusar-Poli, P., Howes, O., Bechdolf, A., Borgwardt, S., 2012. Mapping vulnerability to bipolar disorder: a systematic review and meta-analysis of neuroimaging studies. *Journal of psychiatry & neuroscience : JPN* 37, 170-184.
- Fusar-Poli, P., Radua, J., Frascarelli, M., Mechelli, A., Borgwardt, S., Di Fabio, F., Biondi, M., Ioannidis, J.P., David, S.P., 2014. Evidence of reporting biases in voxel-

- based morphometry (VBM) studies of psychiatric and neurological disorders. *Human brain mapping* 35, 3052-3065.
- Gargus, J.J., 2009. Genetic calcium signaling abnormalities in the central nervous system: seizures, migraine, and autism. *Annals of the New York Academy of Sciences* 1151, 133-156.
- Gasser, A., Ho, T.S., Cheng, X., Chang, K.J., Waxman, S.G., Rasband, M.N., Dib-Hajj, S.D., 2012. An ankyrinG-binding motif is necessary and sufficient for targeting Nav1.6 sodium channels to axon initial segments and nodes of Ranvier. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 32, 7232-7243.
- Gatt, J.M., Burton, K.L., Williams, L.M., Schofield, P.R., 2015. Specific and common genes implicated across major mental disorders: a review of meta-analysis studies. *Journal of psychiatric research* 60, 1-13.
- Georgieva, L., Dimitrova, A., Ivanov, D., Nikolov, I., Williams, N.M., Grozeva, D., Zaharieva, I., Toncheva, D., Owen, M.J., Kirov, G., O'Donovan, M.C., 2008. Support for neuregulin 1 as a susceptibility gene for bipolar disorder and schizophrenia. *Biological psychiatry* 64, 419-427.
- Goes, F.S., 2016. Genetics of Bipolar Disorder: Recent Update and Future Directions. *The Psychiatric clinics of North America* 39, 139-155.
- Gonzalez-Castro, T.B., Nicolini, H., Lanzagorta, N., Lopez-Narvaez, L., Genis, A., Pool Garcia, S., Tovilla-Zarate, C.A., 2015. The role of brain-derived neurotrophic factor (BDNF) Val66Met genetic polymorphism in bipolar disorder: a case-control study, comorbidities, and meta-analysis of 16,786 subjects. *Bipolar disorders* 17, 27-38.
- Gottesman, I.I., Gould, T.D., 2003. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry* 160, 636-645.
- Grande, I., Berk, M., Birmaher, B., Vieta, E., 2016. Bipolar disorder. *Lancet (London, England)* 387, 1561-1572.
- Grande, I., Magalhaes, P.V., Chendo, I., Stertz, L., Fries, G.R., Cereser, K.M., Cunha, A.B., Goi, P., Kunz, M., Udina, M., Martin-Santos, R., Frey, B.N., Vieta, E., Kapczinski, F., 2014. Val66Met polymorphism and serum brain-derived neurotrophic factor in bipolar disorder: an open-label trial. *Acta psychiatrica Scandinavica* 129, 393-400.
- Green, E.K., Hamshere, M., Forty, L., Gordon-Smith, K., Fraser, C., Russell, E., Grozeva, D., Kirov, G., Holmans, P., Moran, J.L., Purcell, S., Sklar, P., Owen, M.J., O'Donovan, M.C., Jones, L., Jones, I.R., Craddock, N., 2013. Replication of bipolar disorder susceptibility alleles and identification of two novel genome-wide significant associations in a new bipolar disorder case-control sample. *Molecular psychiatry* 18, 1302-1307.
- Green, E.K., Raybould, R., Macgregor, S., Gordon-Smith, K., Heron, J., Hyde, S., Grozeva, D., Hamshere, M., Williams, N., Owen, M.J., O'Donovan, M.C., Jones, L., Jones, I., Kirov, G., Craddock, N., 2005. Operation of the schizophrenia susceptibility gene, neuregulin 1, across traditional diagnostic boundaries to increase risk for bipolar disorder. *Archives of general psychiatry* 62, 642-648.
- Group, P.G.C.B.D.W., 2011. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nature genetics* 43, 977-983.
- Gurung, R., Prata, D.P., 2015. What is the impact of genome-wide supported risk variants for schizophrenia and bipolar disorder on brain structure and function? A systematic review. *Psychological medicine* 45, 2461-2480.
- Gutierrez-Fernandez, A., Palomino, A., Gonzalez-Pinto, A., Ugarte, A., Hernanz, M., Mendibil, B., Etxebeste, M., Pacheco, L., Gonzalez-Garcia, G., Matute, C., 2014. Novel

- association of Neuregulin 1 gene with bipolar disorder but not with schizophrenia. *Schizophrenia research* 159, 552-553.
- Ham, B.J., Greenberg, T., Chase, H.W., Phillips, M.L., 2016. Impact of the glucocorticoid receptor Bcl i polymorphism on reward expectancy and prediction error related ventral striatal reactivity in depressed and healthy individuals. *Journal of Psychopharmacology* 30, 48-55.
- Harrisberger, F., Smieskova, R., Schmidt, A., Lenz, C., Walter, A., Wittfeld, K., Grabe, H.J., Lang, U.E., Fusar-Poli, P., Borgwardt, S., 2015. BDNF Val66Met polymorphism and hippocampal volume in neuropsychiatric disorders: A systematic review and meta-analysis. *Neuroscience and biobehavioral reviews* 55, 107-118.
- Hashimoto, R., Ohi, K., Yamamori, H., Yasuda, Y., Fujimoto, M., Umeda-Yano, S., Watanabe, Y., Fukunaga, M., Takeda, M., 2015. Imaging genetics and psychiatric disorders. *Current molecular medicine* 15, 168-175.
- Hasler, G., Wolf, A., 2015. Toward stratified treatments for bipolar disorders. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* 25, 283-294.
- Hayes, J.F., Miles, J., Walters, K., King, M., Osborn, D.P., 2015. A systematic review and meta-analysis of premature mortality in bipolar affective disorder. *Acta psychiatrica Scandinavica* 131, 417-425.
- Heils, A., Teufel, A., Petri, S., Stober, G., Riederer, P., Bengel, D., Lesch, K.P., 1996. Allelic variation of human serotonin transporter gene expression. *Journal of neurochemistry* 66, 2621-2624.
- Hempstead, B.L., 2015. Brain-Derived Neurotrophic Factor: Three Ligands, Many Actions. *Transactions of the American Clinical and Climatological Association* 126, 9-19.
- Hori, H., Yamamoto, N., Teraishi, T., Ota, M., Fujii, T., Sasayama, D., Matsuo, J., Kinoshita, Y., Hattori, K., Nagashima, A., Ishida, I., Koga, N., Higuchi, T., Kunugi, H., 2014. Cognitive effects of the ANK3 risk variants in patients with bipolar disorder and healthy individuals. *Journal of affective disorders* 158, 90-96.
- Hou, L., Bergen, S.E., Akula, N., Song, J., Hultman, C.M., Landen, M., Adli, M., Alda, M., Arda, R., Arias, B., Aubry, J.M., Backlund, L., Badner, J.A., Barrett, T.B., Bauer, M., Baune, B.T., Bellivier, F., Benabarre, A., Bengesser, S., Berrettini, W.H., Bhattacharjee, A.K., Biernacka, J.M., Birner, A., Bloss, C.S., Brichant-Petitjean, C., Bui, E.T., Byerley, W., Cervantes, P., Chillotti, C., Cichon, S., Colom, F., Coryell, W., Craig, D.W., Cruceanu, C., Czerski, P.M., Davis, T., Dayer, A., Degenhardt, F., Del Zompo, M., DePaulo, J.R., Edenberg, H.J., Etain, B., Falkai, P., Foroud, T., Forstner, A.J., Frisen, L., Frye, M.A., Fullerton, J.M., Gard, S., Garnham, J.S., Gershon, E.S., Goes, F.S., Greenwood, T.A., Grigoriu-Serbanescu, M., Hauser, J., Heilbronner, U., Heilmann-Heimbach, S., Herms, S., Hipolito, M., Hitturlingappa, S., Hoffmann, P., Hofmann, A., Jamain, S., Jimenez, E., Kahn, J.P., Kassem, L., Kelsoe, J.R., Kittel-Schneider, S., Kliwicki, S., Koller, D.L., Konig, B., Lackner, N., Laje, G., Lang, M., Lavebratt, C., Lawson, W.B., Leboyer, M., Leckband, S.G., Liu, C., Maaser, A., Mahon, P.B., Maier, W., Maj, M., Manchia, M., Martinsson, L., McCarthy, M.J., McElroy, S.L., McInnis, M.G., McKinney, R., Mitchell, P.B., Mitjans, M., Mondimore, F.M., Monteleone, P., Muhleisen, T.W., Nievergelt, C.M., Nothen, M.M., Novak, T., Nurnberger, J.I., Jr., Nwulia, E.A., Osby, U., Pfennig, A., Potash, J.B., Propping, P., Reif, A., Reininghaus, E., Rice, J., Rietschel, M., Rouleau, G.A., Rybakowski, J.K., Schalling, M., Scheftner, W.A., Schofield, P.R., Schork, N.J., Schulze, T.G., Schumacher, J., Schweizer, B.W., Severino, G., Shekhtman, T., Shilling, P.D., Simhandl, C., Slaney, C.M., Smith, E.N., Squassina, A.,

- Stamm, T., Stopkova, P., Streit, F., Strohmaier, J., Szelinger, S., Tighe, S.K., Tortorella, A., Turecki, G., Vieta, E., Volkert, J., Witt, S.H., Wright, A., Zandi, P.P., Zhang, P., Zollner, S., McMahon, F.J., 2016. Genome-wide association study of 40,000 individuals identifies two novel loci associated with bipolar disorder. *Human molecular genetics* 25, 3383-3394.
- Ioannidis, J.P., Munafo, M.R., Fusar-Poli, P., Nosek, B.A., David, S.P., 2014. Publication and other reporting biases in cognitive sciences: detection, prevalence, and prevention. *Trends in cognitive sciences* 18, 235-241.
- Ivleva, E.I., Morris, D.W., Moates, A.F., Suppes, T., Thaker, G.K., Tamminga, C.A., 2010. Genetics and intermediate phenotypes of the schizophrenia--bipolar disorder boundary. *Neuroscience and biobehavioral reviews* 34, 897-921.
- Jogia, J., Ruberto, G., Lelli-Chiesa, G., Vassos, E., Maieru, M., Tatarelli, R., Girardi, P., Collier, D., Frangou, S., 2011. The impact of the CACNA1C gene polymorphism on frontolimbic function in bipolar disorder. *Molecular psychiatry* 16, 1070-1071.
- Jonassen, R., Landro, N.I., 2014. Serotonin transporter polymorphisms (5-HTTLPR) in emotion processing: implications from current neurobiology. *Progress in neurobiology* 117, 41-53.
- Kempton, M.J., Salvador, Z., Munafo, M.R., Geddes, J.R., Simmons, A., Frangou, S., Williams, S.C., 2011. Structural neuroimaging studies in major depressive disorder. Meta-analysis and comparison with bipolar disorder. *Archives of general psychiatry* 68, 675-690.
- Kerner, B., 2014. Genetics of bipolar disorder. *Appl Clin Genet* 7, 33-42.
- Kittel-Schneider, S., Wobrock, T., Scherk, H., Schneider-Axmann, T., Trost, S., Zilles, D., Wolf, C., Schmitt, A., Malchow, B., Hasan, A., Backens, M., Reith, W., Falkai, P., Gruber, O., Reif, A., 2015. Influence of DGKH variants on amygdala volume in patients with bipolar affective disorder and schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 265, 127-136.
- Kupferschmidt, D.A., Zakzanis, K.K., 2011. Toward a functional neuroanatomical signature of bipolar disorder: quantitative evidence from the neuroimaging literature. *Psychiatry research* 193, 71-79.
- Kurnianingsih, Y.A., Kuswanto, C.N., McIntyre, R.S., Qiu, A., Ho, B.C., Sim, K., 2011. Neurocognitive-genetic and neuroimaging-genetic research paradigms in schizophrenia and bipolar disorder. *J Neural Transm (Vienna)* 118, 1621-1639.
- Kuswanto, C.N., Sum, M.Y., Thng, C.R., Zhang, Y.B., Yang, G.L., Nowinski, W.L., Sitoh, Y.Y., Low, C.M., Sim, K., 2013. GRIN2B gene and associated brain cortical white matter changes in bipolar disorder: a preliminary combined platform investigation. *BioMed research international* 2013, 635131.
- Lee, K.W., Woon, P.S., Teo, Y.Y., Sim, K., 2012. Genome wide association studies (GWAS) and copy number variation (CNV) studies of the major psychoses: what have we learnt? *Neuroscience and biobehavioral reviews* 36, 556-571.
- Lelli-Chiesa, G., Kempton, M.J., Jogia, J., Tatarelli, R., Girardi, P., Powell, J., Collier, D.A., Frangou, S., 2011. The impact of the Val158Met catechol-O-methyltransferase genotype on neural correlates of sad facial affect processing in patients with bipolar disorder and their relatives. *Psychological medicine* 41, 779-788.
- Lesch, K.P., Bengel, D., Heils, A., Sabol, S.Z., Greenberg, B.D., Petri, S., Benjamin, J., Muller, C.R., Hamer, D.H., Murphy, D.L., 1996. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science (New York, N.Y.)* 274, 1527-1531.

- Leussis, M.P., Madison, J.M., Petryshen, T.L., 2012. Ankyrin 3: genetic association with bipolar disorder and relevance to disease pathophysiology. *Biology of mood & anxiety disorders* 2, 18.
- Li, M., Chang, H., Xiao, X., 2016. BDNF Val66Met polymorphism and bipolar disorder in European populations: A risk association in case-control, family-based and GWAS studies. *Neuroscience and biobehavioral reviews* 68, 218-233.
- Lim, C.S., Baldessarini, R.J., Vieta, E., Yucel, M., Bora, E., Sim, K., 2013. Longitudinal neuroimaging and neuropsychological changes in bipolar disorder patients: review of the evidence. *Neuroscience and biobehavioral reviews* 37, 418-435.
- Linke, J., Witt, S.H., King, A.V., Nieratschker, V., Poupon, C., Gass, A., Hennerici, M.G., Rietschel, M., Wessa, M., 2011. Genome-wide supported risk variant for bipolar disorder alters anatomical connectivity in the human brain. *NeuroImage* 59, 3288-3296.
- Lippard, E.T., Jensen, K.P., Wang, F., Johnston, J.A., Spencer, L., Pittman, B., Gelernter, J., Blumberg, H.P., 2016. Effects of ANK3 variation on gray and white matter in bipolar disorder. *Molecular psychiatry*.
- Liu, X., Akula, N., Skup, M., Brotman, M.A., Leibenluft, E., McMahon, F.J., 2010. A genome-wide association study of amygdala activation in youths with and without bipolar disorder. *Journal of the American Academy of Child and Adolescent Psychiatry* 49, 33-41.
- Lopez, A.Y., Wang, X., Xu, M., Maheshwari, A., Curry, D., Lam, S., Adesina, A.M., Noebels, J.L., Sun, Q.Q., Cooper, E.C., 2016. Ankyrin-G isoform imbalance and interneuronopathy link epilepsy and bipolar disorder. *Molecular psychiatry*.
- Lu, B., Gottschalk, W., 2000. Modulation of hippocampal synaptic transmission and plasticity by neurotrophins. *Prog Brain Res* 128, 231-241.
- Macedo, D.S., Medeiros, C.D., Cordeiro, R.C., Sousa, F.C., Santos, J.V., Morais, T.A., Hyphantis, T.N., McIntyre, R.S., Quevedo, J., Carvalho, A.F., 2012. Effects of alpha-lipoic acid in an animal model of mania induced by D-amphetamine. *Bipolar disorders* 14, 707-718.
- Mallas, E., Carletti, F., Chaddock, C.A., Shergill, S., Woolley, J., Picchioni, M.M., McDonald, C., Touloupoulou, T., Kravariti, E., Kalidindi, S., Bramon, E., Murray, R., Barker, G.J., Prata, D.P., 2016a. The impact of CACNA1C gene, and its epistasis with ZNF804A, on white matter microstructure in health, schizophrenia and bipolar disorder. *Genes, Brain and Behavior*.
- Mallas, E.J., Carletti, F., Chaddock, C.A., Woolley, J., Picchioni, M.M., Shergill, S.S., Kane, F., Allin, M.P., Barker, G.J., Prata, D.P., 2016b. Genome-wide discovered psychosis-risk gene ZNF804A impacts on white matter microstructure in health, schizophrenia and bipolar disorder. *PeerJ* 4, e1570.
- Marballi, K., Cruz, D., Thompson, P., Walss-Bass, C., 2012. Differential neuregulin 1 cleavage in the prefrontal cortex and hippocampus in schizophrenia and bipolar disorder: preliminary findings. *PloS one* 7, e36431.
- Matsuo, K., Walss-Bass, C., Nery, F.G., Nicoletti, M.A., Hatch, J.P., Frey, B.N., Monkul, E.S., Zunta-Soares, G.B., Bowden, C.L., Escamilla, M.A., Soares, J.C., 2009. Neuronal correlates of brain-derived neurotrophic factor Val66Met polymorphism and morphometric abnormalities in bipolar disorder. *Neuropsychopharmacology* 34, 1904-1913.
- McIntosh, A.M., Moorhead, T.W., McKirdy, J., Sussmann, J.E., Hall, J., Johnstone, E.C., Lawrie, S.M., 2007. Temporal grey matter reductions in bipolar disorder are

- associated with the BDNF Val66Met polymorphism. *Molecular psychiatry* 12, 902-903.
- Mechelli, A., Prata, D., Papagni, S.A., Tognin, S., Kambeitz, J., Fu, C., Picchioni, M., Walshe, M., Touloupoulou, T., Bramon, E., Murray, R., McGuire, P., 2012. Genetic vulnerability to psychosis and cortical function: epistatic effects between DAAO and G72. *Current pharmaceutical design* 18, 510-517.
- Mechelli, A., Prata, D.P., Fu, C.H., Picchioni, M., Kane, F., Kalidindi, S., McDonald, C., Demjaha, A., Kravariti, E., Touloupoulou, T., Murray, R., Collier, D.A., McGuire, P.K., 2008. The effects of neuregulin1 on brain function in controls and patients with schizophrenia and bipolar disorder. *NeuroImage* 42, 817-826.
- Mei, L., Nave, K.A., 2014. Neuregulin-ERBB signaling in the nervous system and neuropsychiatric diseases. *Neuron* 83, 27-49.
- Merikangas, K.R., Jin, R., He, J.P., Kessler, R.C., Lee, S., Sampson, N.A., Viana, M.C., Andrade, L.H., Hu, C., Karam, E.G., Ladea, M., Medina-Mora, M.E., Ono, Y., Posada-Villa, J., Sagar, R., Wells, J.E., Zarkov, Z., 2011. Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Archives of general psychiatry* 68, 241-251.
- Mirakhor, A., Moorhead, T.W., Stanfield, A.C., McKirdy, J., Sussmann, J.E., Hall, J., Lawrie, S.M., Johnstone, E.C., McIntosh, A.M., 2009. Changes in gyrification over 4 years in bipolar disorder and their association with the brain-derived neurotrophic factor valine(66) methionine variant. *Biological psychiatry* 66, 293-297.
- Miskowiak, K.W., Carvalho, A.F., 2014. 'Hot' cognition in major depressive disorder: a systematic review. *CNS & neurological disorders drug targets* 13, 1787-1803.
- Miskowiak, K.W., Kjaerstad, H.L., Meluken, I., Petersen, J.Z., Maciel, B.R., Kohler, C.A., Vinberg, M., Kessing, L.V., Carvalho, A.F., 2016. The search for neuroimaging and cognitive endophenotypes: A critical systematic review of studies involving unaffected first-degree relatives of individuals with bipolar disorder. *Neuroscience and biobehavioral reviews*.
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., 2010. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *International journal of surgery (London, England)* 8, 336-341.
- Mostaid, M.S., Lloyd, D., Liberg, B., Sundram, S., Pereira, A., Pantelis, C., Karl, T., Weickert, C.S., Everall, I.P., Bousman, C.A., 2016. Neuregulin-1 and schizophrenia in the genome-wide association study era. *Neuroscience and biobehavioral reviews* 68, 387-409.
- Muhleisen, T.W., Leber, M., Schulze, T.G., Strohmaier, J., Degenhardt, F., Treutlein, J., Mattheisen, M., Forstner, A.J., Schumacher, J., Breuer, R., Meier, S., Herms, S., Hoffmann, P., Lacour, A., Witt, S.H., Reif, A., Muller-Myhsok, B., Lucae, S., Maier, W., Schwarz, M., Vedder, H., Kammerer-Ciernioch, J., Pfennig, A., Bauer, M., Hautzinger, M., Moebus, S., Priebe, L., Czerski, P.M., Hauser, J., Lissowska, J., Szeszenia-Dabrowska, N., Brennan, P., McKay, J.D., Wright, A., Mitchell, P.B., Fullerton, J.M., Schofield, P.R., Montgomery, G.W., Medland, S.E., Gordon, S.D., Martin, N.G., Krasnow, V., Chuchalin, A., Babadjanova, G., Pantelejeva, G., Abramova, L.I., Tiganov, A.S., Polonikov, A., Khusnutdinova, E., Alda, M., Grof, P., Rouleau, G.A., Turecki, G., Laprise, C., Rivas, F., Mayoral, F., Kogevinas, M., Grigoriou-Serbanescu, M., Propping, P., Becker, T., Rietschel, M., Nothen, M.M., Cichon, S., 2014. Genome-wide association study reveals two new risk loci for bipolar disorder. *Nature communications* 5, 3339.

- Murphy, S.E., Norbury, R., Godlewska, B.R., Cowen, P.J., Mannie, Z.M., Harmer, C.J., Munafo, M.R., 2013. The effect of the serotonin transporter polymorphism (5-HTTLPR) on amygdala function: a meta-analysis. *Molecular psychiatry* 18, 512-520.
- Nortje, G., Stein, D.J., Radua, J., Mataix-Cols, D., Horn, N., 2013. Systematic review and voxel-based meta-analysis of diffusion tensor imaging studies in bipolar disorder. *Journal of affective disorders* 150, 192-200.
- Notaras, M., Hill, R., van den Buuse, M., 2015a. The BDNF gene Val66Met polymorphism as a modifier of psychiatric disorder susceptibility: progress and controversy. *Molecular psychiatry* 20, 916-930.
- Notaras, M., Hill, R., van den Buuse, M., 2015b. A role for the BDNF gene Val66Met polymorphism in schizophrenia? A comprehensive review. *Neuroscience and biobehavioral reviews* 51, 15-30.
- Nurnberger, J.I., Jr., Koller, D.L., Jung, J., Edenberg, H.J., Foroud, T., Guella, I., Vawter, M.P., Kelsoe, J.R., 2014. Identification of pathways for bipolar disorder: a meta-analysis. *JAMA psychiatry* 71, 657-664.
- Oertel-Knochel, V., Lancaster, T.M., Knochel, C., Stablein, M., Storchak, H., Reinke, B., Jurcoane, A., Kniep, J., Prvulovic, D., Mantripragada, K., Tansey, K.E., O'Donovan, M.C., Owen, M.J., Linden, D.E., 2015. Schizophrenia risk variants modulate white matter volume across the psychosis spectrum: evidence from two independent cohorts. *NeuroImage. Clinical* 7, 764-770.
- Olsson, C.A., Foley, D.L., Parkinson-Bates, M., Byrnes, G., McKenzie, M., Patton, G.C., Morley, R., Anney, R.J., Craig, J.M., Saffery, R., 2010. Prospects for epigenetic research within cohort studies of psychological disorder: a pilot investigation of a peripheral cell marker of epigenetic risk for depression. *Biological psychology* 83, 159-165.
- Ota, M., Hori, H., Sato, N., Yoshida, F., Hattori, K., Teraishi, T., Kunugi, H., 2016. The effects of ankyrin 3 gene risk variants on brain structures in patients with bipolar disorder and healthy subjects. *Psychiatry and clinical neurosciences*.
- Ou, X., Crane, D.E., MacIntosh, B.J., Young, L.T., Arnold, P., Ameis, S., Goldstein, B.I., 2015. CACNA1C rs1006737 genotype and bipolar disorder: Focus on intermediate phenotypes and cardiovascular comorbidity. *Neuroscience and biobehavioral reviews* 55, 198-210.
- Papagni, S.A., Mechelli, A., Prata, D.P., Kambeitz, J., Fu, C.H., Picchioni, M., Walshe, M., Toulopoulou, T., Bramon, E., Murray, R.M., Collier, D.A., Bellomo, A., McGuire, P., 2011. Differential effects of DAAO on regional activation and functional connectivity in schizophrenia, bipolar disorder and controls. *NeuroImage* 56, 2283-2291.
- Papiol, S., Molina, V., Desco, M., Rosa, A., Reig, S., Sanz, J., Palomo, T., Fanas, L., 2008. Gray matter deficits in bipolar disorder are associated with genetic variability at interleukin-1 beta gene (2q13). *Genes, brain, and behavior* 7, 796-801.
- Perrier, E., Pompei, F., Ruberto, G., Vassos, E., Collier, D., Frangou, S., 2011a. Initial evidence for the role of CACNA1C on subcortical brain morphology in patients with bipolar disorder. *European psychiatry : the journal of the Association of European Psychiatrists* 26, 135-137.
- Perrier, E., Pompei, F., Ruberto, G., Vassos, E., Collier, D., Frangou, S., 2011b. Initial evidence for the role of CACNA1C on subcortical brain morphology in patients with bipolar disorder. *European Psychiatry* 26, 135-137.




- Piguet, C., Fodouliau, L., Aubry, J.M., Vuilleumier, P., Houenou, J., 2015. Bipolar disorder: Functional neuroimaging markers in relatives. *Neuroscience and biobehavioral reviews* 57, 284-296.
- Poletti, S., Aggio, V., Bollettini, I., Falini, A., Colombo, C., Benedetti, F., 2016. SREBF-2 polymorphism influences white matter microstructure in bipolar disorder. *Psychiatry Research - Neuroimaging* 257, 39-46.
- Poletti, S., Locatelli, C., Radaelli, D., Lorenzi, C., Smeraldi, E., Colombo, C., Benedetti, F., 2014. Effect of early stress on hippocampal gray matter is influenced by a functional polymorphism in EAAT2 in bipolar disorder. *Progress in neuro-psychopharmacology & biological psychiatry* 51, 146-152.
- Prata, D.P., Mechelli, A., Picchioni, M., Fu, C.H.Y., Kane, F., Kalidindi, S., McDonald, C., Kravariti, E., Touloupoulou, T., Bramon, E., Walshe, M., Murray, R., Collier, D.A., McGuire, P.K., 2011. No association of Disrupted-in-Schizophrenia-1 variation with prefrontal function in patients with schizophrenia and bipolar disorder. *Genes, Brain and Behavior* 10, 276-285.
- Provenzi, L., Giorda, R., Beri, S., Montirosso, R., 2016. SLC6A4 methylation as an epigenetic marker of life adversity exposures in humans: A systematic review of literature. *Neuroscience and biobehavioral reviews* 71, 7-20.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J., Sham, P.C., 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *American journal of human genetics* 81, 559-575.
- Radua, J., Surguladze, S.A., Marshall, N., Walshe, M., Bramon, E., Collier, D.A., Prata, D.P., Murray, R.M., McDonald, C., 2013. The impact of CACNA1C allelic variation on effective connectivity during emotional processing in bipolar disorder. *Molecular psychiatry* 18, 526-527.
- Rasetti, R., Weinberger, D.R., 2011. Intermediate phenotypes in psychiatric disorders. *Current opinion in genetics & development* 21, 340-348.
- Rizzo, L.B., Costa, L.G., Mansur, R.B., Swardfager, W., Belangero, S.I., Grassi-Oliveira, R., McIntyre, R.S., Bauer, M.E., Brietzke, E., 2014. The theory of bipolar disorder as an illness of accelerated aging: implications for clinical care and research. *Neuroscience and biobehavioral reviews* 42, 157-169.
- Roiser, J.P., Cannon, D.M., Gandhi, S.K., Taylor Tavares, J., Erickson, K., Wood, S., Klaver, J.M., Clark, L., Zarate, C.A., Jr., Sahakian, B.J., Drevets, W.C., 2009. Hot and cold cognition in unmedicated depressed subjects with bipolar disorder. *Bipolar disorders* 11, 178-189.
- Rolstad, S., Sellgren Majkowitz, C., Joas, E., Ekman, C.J., Palsson, E., Landen, M., 2016. Polymorphisms of BDNF and CACNA1C are not associated with cognitive functioning in bipolar disorder or healthy controls. *Cognitive neuropsychiatry* 21, 271-278.
- Rosa, A.R., Singh, N., Whitaker, E., de Brito, M., Lewis, A.M., Vieta, E., Churchill, G.C., Geddes, J.R., Goodwin, G.M., 2014. Altered plasma glutathione levels in bipolar disorder indicates higher oxidative stress; a possible risk factor for illness onset despite normal brain-derived neurotrophic factor (BDNF) levels. *Psychological medicine* 44, 2409-2418.
- Ruberto, G., Vassos, E., Lewis, C.M., Tatarelli, R., Girardi, P., Collier, D., Frangou, S., 2011. The cognitive impact of the ANK3 risk variant for bipolar disorder: initial evidence of selectivity to signal detection during sustained attention. *PloS one* 6, e16671.

- Rybakowski, J.K., Borkowska, A., Czerski, P.M., Skibinska, M., Hauser, J., 2003. Polymorphism of the brain-derived neurotrophic factor gene and performance on a cognitive prefrontal test in bipolar patients. *Bipolar disorders* 5, 468-472.
- Rybakowski, J.K., Borkowska, A., Skibinska, M., Szczepankiewicz, A., Kapelski, P., Leszczynska-Rodziewicz, A., Czerski, P.M., Hauser, J., 2006. Prefrontal cognition in schizophrenia and bipolar illness in relation to Val66Met polymorphism of the brain-derived neurotrophic factor gene. *Psychiatry and clinical neurosciences* 60, 70-76.
- Scherk, H., Backens, M., Schneider-Axmann, T., Usher, J., Kemmer, C., Reith, W., Falkai, P., Gruber, O., 2009a. Cortical neurochemistry in euthymic patients with bipolar I disorder. *The world journal of biological psychiatry : the official journal of the World Federation of Societies of Biological Psychiatry* 10, 285-294.
- Scherk, H., Gruber, O., Menzel, P., Schneider-Axmann, T., Kemmer, C., Usher, J., Reith, W., Meyer, J., Falkai, P., 2009b. 5-HTTLPR genotype influences amygdala volume. *Eur Arch Psychiatry Clin Neurosci* 259, 212-217.
- Shah, M.P., Wang, F., Kalmar, J.H., Chepenik, L.G., Tie, K., Pittman, B., Jones, M.M., Constable, R.T., Gelernter, J., Blumberg, H.P., 2009. Role of variation in the serotonin transporter protein gene (SLC6A4) in trait disturbances in the ventral anterior cingulate in bipolar disorder. *Neuropsychopharmacology* 34, 1301-1310.
- Sklar, P., Smoller, J.W., Fan, J., Ferreira, M.A., Perlis, R.H., Chambert, K., Nimgaonkar, V.L., McQueen, M.B., Faraone, S.V., Kirby, A., de Bakker, P.I., Ogdie, M.N., Thase, M.E., Sachs, G.S., Todd-Brown, K., Gabriel, S.B., Sougnez, C., Gates, C., Blumenstiel, B., Defelice, M., Ardlie, K.G., Franklin, J., Muir, W.J., McGhee, K.A., MacIntyre, D.J., McLean, A., VanBeck, M., McQuillin, A., Bass, N.J., Robinson, M., Lawrence, J., Anjorin, A., Curtis, D., Scolnick, E.M., Daly, M.J., Blackwood, D.H., Gurling, H.M., Purcell, S.M., 2008. Whole-genome association study of bipolar disorder. *Molecular psychiatry* 13, 558-569.
- Smoller, J.W., Finn, C.T., 2003. Family, twin, and adoption studies of bipolar disorder. *American journal of medical genetics. Part C, Seminars in medical genetics* 123c, 48-58.
- Soeiro-de-Souza, M.G., Otaduy, M.C., Dias, C.Z., Bio, D.S., Machado-Vieira, R., Moreno, R.A., 2012. The impact of the CACNA1C risk allele on limbic structures and facial emotions recognition in bipolar disorder subjects and healthy controls. *Journal of affective disorders* 141, 94-101.
- Tesli, M., Egeland, R., Sonderby, I.E., Haukvik, U.K., Bettella, F., Hibar, D.P., Thompson, P.M., Rimol, L.M., Melle, I., Agartz, I., Djurovic, S., Andreassen, O.A., 2013. No evidence for association between bipolar disorder risk gene variants and brain structural phenotypes. *Journal of affective disorders* 151, 291-297.
- Tramontina, J.F., Yates, D., Magalhaes, P.V., Trentini, C., Sant'anna, M.K., Fries, G.R., Bock, H., Saraiva-Pereira, M.L., Kapczinski, F., 2009. Brain-derived neurotrophic factor gene val66met polymorphism and executive functioning in patients with bipolar disorder. *Revista brasileira de psiquiatria (Sao Paulo, Brazil : 1999)* 31, 136-140.
- Vieta, E., 2015. [Personalised medicine applied to mental health: Precision psychiatry]. *Revista de psiquiatria y salud mental* 8, 117-118.
- Wegbreit, E., Cushman, G.K., Puzia, M.E., Weissman, A.B., Kim, K.L., Laird, A.R., Dickstein, D.P., 2014. Developmental meta-analyses of the functional neural correlates of bipolar disorder. *JAMA psychiatry* 71, 926-935.


- Winterer, G., Konrad, A., Vucurevic, G., Musso, F., Stoeter, P., Dahmen, N., 2008. Association of 5' end neuregulin-1 (NRG1) gene variation with subcortical medial frontal microstructure in humans. *NeuroImage* 40, 712-718.
- Wolf, C., Mohr, H., Schneider-Axmann, T., Reif, A., Wobrock, T., Scherk, H., Kraft, S., Schmitt, A., Falkai, P., Gruber, O., 2014. CACNA1C genotype explains interindividual differences in amygdala volume among patients with schizophrenia. *European Archives of Psychiatry and Clinical Neuroscience* 264, 93-102.
- Zeni, C.P., Mwangi, B., Cao, B., Hasan, K.M., Walss-Bass, C., Zunta-Soares, G., Soares, J.C., 2016. Interaction between BDNF rs6265 Met allele and low family cohesion is associated with smaller left hippocampal volume in pediatric bipolar disorder. *Journal of affective disorders* 189, 94-97.
- Zuliani, R., Moorhead, T.W., Job, D., McKirdy, J., Sussmann, J.E., Johnstone, E.C., Lawrie, S.M., Brambilla, P., Hall, J., McIntosh, A.M., 2009. Genetic variation in the G72 (DAOA) gene affects temporal lobe and amygdala structure in subjects affected by bipolar disorder. *Bipolar disorders* 11, 621-627.

## Figure legends

Figure 1. PRISMA flowchart of the study selection process.



**Figure 2. Figure 2. Brain map representing the approximate locations of replicated gene-neuroimaging findings in BD patients compared to healthy control groups. VBM studies (k=2) found decreased volumes of the left and right hippocampi in carriers of the Met allele of the *BDNF* Val66Met polymorphism (yellow).**



**Table 1. Studies investigating the association of genetic polymorphisms and brain structure in BD using voxel based morphometry (VBM) in magnetic resonance imaging (MRI).**

Author	Gene	Subjects, n	Genetic Polymorphisms, n	Gender, female, n (%)	Age, years, mean $\pm$ SD	Methods	Statistically significant difference?*	Main findings
<b>Cao et al. (2016)</b>	<i>BDNF</i>	BD, 48 HC, 60	Met carriers, 13 Met carriers, 19	32 (66.6) 39 (65.0)	41.0 $\pm$ 12.6 40.5 $\pm$ 12.9	MRI 1.5T VBM ROIs: Hippocampal cortical/subcortical volume	<b>Yes</b>	BD patients carrying the <i>BDNF</i> met allele had smaller hippocampal volumes compared to HCs.
<b>Zeni et al. (2016)</b>	<i>BDNF</i> rs6265	BD, 29 HC, 22	Val/Val, 19 Met carriers, 10 Val/Val, 12 Met carriers, 10	11 (57.8) 3 (30.0) 3 (25.0) 5 (50.0)	14.8 $\pm$ 2.2 11.8 $\pm$ 2.5 12.7 $\pm$ 2.6 13.4 $\pm$ 3.4	MRI VBM ROI: Hippocampal cortical volume	<b>No†</b>	No significant differences between BD patients and HCs in left or right hippocampal volumes.
<b>Chepenik et al. (2009)</b>	<i>BDNF</i>	BD, 20 HC, 18	Val/Val, 12 Val/Met, 7 Met/Met, 1 Val/Val, 12 Val/Met, 8	11 (55.0) 12 (66.6)	21-56*** 18-58***	MRI 1.5T VBM ROI: Hippocampal cortical volume	<b>Yes</b>	Both hippocampal volumes were significantly smaller in participants with BD compared to HCs, and the <i>BDNF</i> met allele was associated with smaller hippocampal volumes in both diagnostic groups.
<b>Mirakhur et al. (2009)</b>	<i>BDNF</i>	BD, 18 HC, 18	One or more Met alleles, 6 One or more Met alleles, 4	10 (55.5) 9 (50.0)	38.4 $\pm$ 8.4 36.7 $\pm$ 13.2	MRI 1.5T VBM Cortical volume, gyrification index	<b>Yes</b>	Individuals with BD carrying one or more <i>BDNF</i> met alleles showed greater losses in GI, an effect that correlated with GM loss in the left hemisphere
<b>Matsuo et al. (2009)</b>	<i>BDNF</i>	BD, 42 HC, 42	Val/Val, 24 Val/Met, 18 Val/Val, 29 Val/Met, 13	19 (79.1) 14 (77.8) 19 (65.5) 8 (61.5)	36.1 $\pm$ 9.3 35.2 $\pm$ 9.7 35.4 $\pm$ 10.5 34.2 $\pm$ 10.4	MRI 1.5T VBM ROIs: dlPFC, ACG, and hippocampus GM volumes	<b>Yes</b>	Anterior CG GM volumes significantly smaller in Val/Met BD compared to Val/Val BD Smaller left dlPFC GM volumes in Val/Met compared to Val/Val subjects within BD and HC groups.
<b>Benedetti et al. (2014)</b>	<i>5-HTTLPR</i>	BD, 136	L/L, 45 L/S, 68 S/S, 23	32 (71.1) 42 (61.7) 19 (82.6)	45.9 $\pm$ 10.9 46.1 $\pm$ 12.2 48.5 $\pm$ 13.6	MRI 3.0T VBM ROI: Hippocampal GM	<b>No</b>	Exposure to early stress correlated with GM volumes in the right prefrontal cortex (Brodmann area 46) in S carriers only.
<b>Scherk et al. (2009b)</b>	<i>5-HTTLPR</i>	BD, 37	L/L, 8 S carriers, 29	4 (50.0) 15 (51.7)	39.8 $\pm$ 14.1 45.1 $\pm$ 11.8	MRI 1.5T VBM	<b>Yes</b>	S carriers showed a relatively increased volume of the right amygdala compared to homozygous

		HC, 37	L/L, 18 S carriers, 19	15 (83.3) 7 (36.8)	39.6 ± 15.1 42.3 ± 10.8	ROI: Amygdala GM volume		L-allele carriers irrespective of diagnostic status.
<b>Wolf et al. (2014)</b>	<i>CACNA1C</i>	BD, 28  HC, 16	A/A+A/G, 16 G/G, 12  A/A+A/G, 8 G/G, 8 (rs1006737)	6 (37.5) 7 (58.3)  4 (50.0) 6 (75.0)	43.9 ± 13.0  33.7 ± 13.4	MRI 1.5T VBM ROI: Amygdala GM volume	<b>No†</b>	The authors found that the amygdala volume was influenced by <i>CACNA1C</i> genotype in BD but did not reach statistical significance.
<b>Soeiro-de-Souza et al. (2012)</b>	<i>CACNA1C</i>	BD, 39  HC, 40	Met/Met, 4 Val/Met, 20 Val/Val, 15 Met/Met, 3 Val/Met, 15 Val/Val, 22	24 (61.5)  20 (50.0)	32.9 ± 10.9  25.9 ± 5.8	MRI 3.0T VBM ROI: Amygdala and hippocampal GM volumes	<b>No†</b>	The <i>CACNA1C</i> genotype had no significant influence on amygdala or hippocampus volumes in either groups.
<b>Perrier et al. (2011b)</b>	<i>CACNA1C</i> Rs1006737	BD, 41  HC, 50	A/A+A/G, 24 G/G, 17  A/A+A/G, 22 G/G, 28	10 (41.6) 11 (64.7)  7 (31.8) 16 (57.1)	44.4 ± 12.3 44.1 ± 11.5  35.6 ± 12.7 34.4 ± 13.7	MRI 1.5T VBM ROI: Basal ganglia, hypothalamus and amygdala GM volumes	<b>Yes</b>	BD patients carrying the risk allele had smaller left putamen than healthy controls.
<b>Benedetti et al. (2015b)</b>	<i>GSK-3β</i>	BD, 150	A/A, 62 A/G, 68 G/G, 20	40 (64.5) 14 (20.5) 14 (70.0)	46.2 ± 11.5 44.9 ± 11.6 46.9 ± 10.8	MRI 3.0T VBM Whole-brain GM volume	<b>No</b>	<i>GSK-3β</i> rs334558*G carriers and the long-term administration of lithium were synergistically associated with increased GM volumes in the right frontal lobe, including the boundaries of subgenual and OFC.
<b>Poletti et al. (2014)</b>	<i>EAAT2</i>	BD, 86	G/G, 14 G/T, 43 T/T, 29	56 (65.1)	46.4 ± 12.7 45.3 ± 11.0 45.0 ± 12.6	MRI 3.0T VBM ROI: Hippocampal GM volume	<b>No</b>	The effect of SLC1A2-181A > C revealed itself only among patients exposed to lower levels of ACE, with T/T homozygotes showing the lowest and G/G the highest hippocampal GM volume.
<b>Kittel-Schneider et al. (2015)</b>	<i>DGKH</i>	BD, 30  HC, 18	No GAT, 15 ≥ 1 GAT, 15  No GAT, 13 ≥ 1 GAT, 5	9 (60.0) 6 (40.0)  10 (76.9) 2 (40.0)	42.1 ± 12.4 45.8 ± 12.4  31.0 ± 11.3 39.4 ± 15.5	MRI 1.5T VBM ROI: Amygdala GM volume	<b>Yes</b>	There was a significant association of the <i>DGKH</i> risk haplotype with increased left amygdala volume in BD, but not in HCs.
<b>Cannon et al. (2012)</b>	<i>NRG1</i>  <i>HAP</i>	BD, 33	T/T, 9 C carriers, 18 Arh1, 8	6 (67.0) 11 (61.0) 5 (63.0)	37.0 ± 11.0 42.0 ± 11.0 44.0 ± 11.0	MRI 1.5T VBM Whole-brain WM volume	<b>Yes</b>	The <i>NRG1</i> SNP and the <i>HAP</i> <sub>ICE</sub> haplotype were associated with abnormal WM volume in the BD group in the fornix, cingulum and PHG circuit

	<i>CNP</i> <i>MOG</i> <i>NRG1</i> <i>HAP</i> <i>CNP</i> <i>MOG</i>	HC, 39	Arh0, 23 G/G, 16 A carriers, 14 G/G, 19 C carriers, 11 T/T, 10 C carriers, 27 Arh1, 11 Arh0, 26 G/G, 21 A carriers, 16 G/G, 21 C carriers, 15	14 (61.0) 10 (63.0) 9 (64.0) 13 (68.0) 6 (55.0) 8 (80.0) 11 (41.0) 4 (34.0) 15 (58.0) 9 (43.0) 8 (50.0) 10 (48.0) 7 (47.0)	41.0 ± 12.0 40.0 ± 12.0 41.0 ± 12.0 41.0 ± 12.0 40.0 ± 10.0 44.0 ± 15.0 40.0 ± 15.0 36.0 ± 13.0 43.0 ± 16.0 39.0 ± 14.0 45.0 ± 17.0 45.0 ± 12.0 34.0 ± 16.0			<i>NRG1</i> (SNP8NRG221533): C carriers > T/T in CG/PHG, splenium and anterior CC <i>HAP</i> <sub>ICE</sub> haplotype: Arh0 < Arh1 in Fornix, caudate, posterior CG <i>MOG</i> (rs2857766): HC G/G > C carriers in CST, MCP, ICP
<b>Papiol et al. (2008)</b>	<i>IL-1β</i>	BD, 20 HC, 45	Non carriers, 8 Allele*2 carriers, 20 Not reported	10 (50.0) 21 (46.6)	43.4 ± 11.7 29.4 ± 9.0	MRI 1.5T VBM Whole-brain WM and GM volumes; ROIs: dlPFC GM, STG GM, hippocampus GM and LV	<b>Yes</b>	A -511C/T polymorphism (rs16944) of <i>IL-1β</i> gene was associated with whole-brain and left dlPFC GM deficits in BD patients.
<b>Tesli et al. (2013)</b>	<i>ANK3</i> <i>CACNA1C</i> <i>ODZ4</i> <i>SYNE1</i>	BD, 121 HC, 219	<i>ANK3</i> (rs9804190, rs10994336 rs10994397) <i>rs1938526</i> ** <i>CACNA1C</i> (rs1006737, rs4765913)** <i>ODZ4</i> (rs12576775, rs2175420)** <i>SYNE1</i> (rs9371601)**	71 (58.6) 102 (46.5)	35.8 ± 11.5 35.9 ± 9.7	MRI 1.5T VBM Whole-brain GM volume	<b>No</b>	There were no significant associations between risk SNPs and structural brain alterations in BD.
<b>Zuliani et al. (2009)</b>	<i>DAOA</i>	BD, 38	M23 CC, 10 M23 CT, 16 M23 TT, 12 M24 AA, 10 M24 AT, 18 M24 TT, 9	7 (70.0) 7 (43.7) 5 (41.6) 7 (70.0) 7 (38.8) 4 (44.4)	38.9 ± 11.0 41.1 ± 11.4 37.5 ± 7.7 41.7 ± 10.9 40.5 ± 11.1 37.1 ± 7.8	MRI 1.5T VBM Whole-brain; ROIs: temporal lobe and amygdala-hippocampal complex GM volumes	<b>Yes</b>	Both M23 and M24 were associated with reductions of GM density within left TP (CC<CT<TT) in the BD group. M23 was also associated with reductions in right amygdala GM density.



		HC, 81	M23 CC, 25 M23 CT, 38 M23 TT, 18 M24 AA, 26 M24 AT, 38 M24 TT, 17	12 (48.0) 17 (44.7) 10 (55.5) 13 (50.0) 16 (42.1) 10 (58.8)	34.4 ± 10.3 35.0 ± 11.6 32.6 ± 8.5 34.5 ± 10.4 35.0 ± 11.6 33.3 ± 8.8			
<b>Bergmann et al. (2013)</b>	<i>ZNF804A</i>	BD, 85  HC, 152	rs13393271, 81 rs359878, 85 rs13393271, 148 rs359878, 151	43 (53.0)  73 (48.0)	36.1 ± 11.0  35.9 ± 9.6	MRI 1.5 T VBM Whole-brain cortical thickness	<b>No</b> ††	There were no associations between any of the SNPs and cortical thickness measures in HC or BD groups.
<b>Oertel-Knochel et al. (2015)</b>	7 risk SZ SNPs from a GWAS	BD, 20 HC, 38	<i>MIR137</i> (rs1625579) <i>CCDC68</i> (rs12966547) <i>CNNM2</i> (rs7914558) <i>NT5C2</i> (rs111915801) <i>MMP16</i> (rs7004633) <i>CSMD1</i> (rs10503253) <i>PCGEM1</i> (rs17662626)**	11 (55.0) 20 (52.6)	39.0 ± 12.1 37.1 ± 11.1	MRI 3.0 T VBM Whole-brain WM volume	<b>No</b>	Increased additive genetic risk for SZ was associated with reduced white matter volume in a group of participants consisting of healthy individuals, SZ first-degree relatives, SZ patients and BD patients, but not in diagnostic groups separately.
<b>Bakken et al. (2011)</b>	GWAS	BD, 97 HC, 181	None	53 (55.0) 87 (48.0)	35.7 ± 11.1 35.9 ± 9.5	MRI 1.5T VBM Whole-brain cortical thickness	<b>No</b>	No SNP associations were genome-wide significant in the BD group.

\* Risk between BD carriers of the risk allele and healthy controls or BD non-carriers of the risk allele. Significant interactions considering only clinical symptoms scores or medication use were not considered.

\*\* Frequencies or N not reported

\*\*\* Mean and SD not reported

† Small sample size was considered a limitation in the original report

†† Limitations were not reported

**Table 2. Studies investigating the association of genetic polymorphisms and brain structure in BD using diffusion tensor imaging (DTI) in magnetic resonance imaging (MRI).**

Author	Gene	Subjects, n	Genetic Polymorphisms, n	Gender, female, n (%)	Age, years, mean $\pm$ SD	Methods	Statistically significant difference?*	Main findings
<i>Diffusion Tensor Imaging (DTI) and VBM</i>								
<b>Ota et al. (2016)</b>	<i>ANK3</i> rs10761482	BD, 43  HC, 229	C/C, 24 T carriers, 19 C/C, 133 T carriers, 96	8 (33.3) 13 (68.4) 94 (70.6) 74 (77.0)	40.8 $\pm$ 9.9 35.9 $\pm$ 7.9 45.8 $\pm$ 15.8 45.3 $\pm$ 15.6	MRI 1.5T DTI/VBM Whole-brain GM volume and WM FA	<b>Yes (DTI)</b>	Decreased FA was found in the forceps minor in non-T-allele BD patients compared with the T-carrier BD group. No main effect of genetic variations were found on the GM volume and the genotype-by-diagnosis interaction.
<b>Lippard et al. (2016)</b>	<i>ANK3</i> rs9804190	BD, 90  HC, 97	C/C, 52 T carriers, 38 C/C, 56 T carriers, 41	35 (67.0) 26 (68.0) 28 (50.0) 23 (53.0)	27.5 $\pm$ 12.2 26.5 $\pm$ 11.1 23.9 $\pm$ 9.0 28.6 $\pm$ 12.9	MRI 3.0 T DTI/VBM Whole-brain analysis ROIs: amygdala and OFC GM; whole-brain and UF FA	<b>Yes (DTI)</b>	BD subjects carrying the T (risk) allele showed decreased FA compared with other subgroups, independent of age within the UF. Compared with BD CC homozygotes, BD T-carriers had lower FA in the UF and anterior CG bilaterally, in dorsomedial frontal WM, in left temporoparietal WM and in posterior dorsomedial WM, among others.
<i>DTI</i>								
<b>Benedetti et al. (2015a)</b>	<i>5-HTTLPR</i>	140, BD	L/L, 47 S carriers, 93	32 (68.0) 36 (38.7)	46.4 $\pm$ 20.7 30.3 $\pm$ 10.2	MRI 3.0T DTI Whole-brain analysis	<b>Yes</b>	S carriers showed significantly increased radial and mean diffusivity in several brain WM tracts (right posterior CG, left anterior CG, CCB, CCg and right posterior CR)
<b>Benedetti et al. (2013)</b>	<i>GSK-3<math>\beta</math></i>	70, BD	T/T, 26 C carriers, 44	19 (73.0) 31 (70.0)	45.7 $\pm$ 11.8 45.7 $\pm$ 11.4	MRI 3.0T DTI Whole-brain analysis	<b>Yes</b>	The rs334558*C carriers and the long-term use of lithium were associated with increased axial diffusivity in several WM fiber tracts (CC, FM, anterior CG and posterior CG bilaterally, including its hippocampal part, left superior and inferior LF, left inferior FOF, left posterior TR, bilateral superior and posterior CR, and bilateral CST).
<b>Kuswanto et al. (2013)</b>	<i>GRIN2B</i>	BD, 14	G/G, 1 G/T, 2 T/T, 11	4 (18.1)	36.9 $\pm$ 12.2	MRI 3.0T DTI Whole-brain analysis	<b>Yes</b>	Compared to G allele, brain FA values were significantly lower in BD patients carrying the T allele in bilateral frontal regions, left parietal region, left occipital region, right occipital

		HC, 22	G/G, 3 G/T, 5 T/T, 14	11 (50.0)	32.7 ± 12.3			region and left CG.
<b>Mallas et al. (2016a)</b>	<i>CACNA1C</i> rs1006737	BD, 43  HC, 124	A/A, 15 A/G, 15 G/G, 13 A/A, 17 A/G, 51 G/G, 56	25 (58.1)  57 (46.0)	41.1 ± 12.3  35.8 ± 13.4	MRI 1.5T DTI Whole-brain analysis	No†	There was no significant main effect of the <i>CACNA1C</i> genotype on FA. In BD patients, the <i>ZNF804A</i> rs1344706 risk genotype increased the magnitude of the effect of the <i>CACNA1C</i> risk genotype, but the association was no longer significant after controlling for age.
<b>Mallas et al. (2016b)</b>	<i>ZNF804A</i> rs1344706	BD, 43  HC, 124	A/A, 19 A/C, 16 C/C, 8 A/A, 59 A/C, 51 C/C, 14	25 (58.1)  57 (46.0)	41.0 ± 12.3  35.7 ± 13.4	MRI 1.5T DTI Whole-brain analysis	No	No areas with a significant diagnosis by genotype interaction were found.
<b>Poletti et al. (2016)</b>	<i>SREBF1</i> rs11868035  <i>SREBF2</i> rs1052717	BD, 93	A/A, 10 A/G, 45 G/G, 38 A/A, 27 A/G, 39 G/G, 27	5 (50.0) 29 (64.4) 28 (73.7) 19 (70.4) 25 (64.1) 18 (66.7)	44.8 ± 13.8 46.1 ± 12.1 43.8 ± 9.2 43.8 ± 12.1 45.7 ± 11.3 45.3 ± 10.3	MRI 3.0T DTI Whole-brain analysis	Yes	No effect on DTI measures of WM integrity was observed for <i>SREBF1</i> polymorphism. The <i>SREBF2</i> rs1052717 polymorphism A/A genotype had increased radial diffusivity compared to A/G and G/G, and the A/A genotype had reduced FA compared to G carriers in cingulum, corpus callosum, superior and inferior longitudinal fasciculi, and anterior thalamic radiation.

\* Risk between BD carriers of the risk allele and healthy controls or BD non-carriers of the risk allele. Significant interactions considering only clinical symptoms scores or medication use were not considered.

† Sample size was not mentioned as a limitation in the original report.

**Table 3. Studies investigating the association of genetic polymorphisms and brain activity in BD using functional magnetic resonance imaging (fMRI) during emotional tasks.**

Author	Gene	Subjects	Genetic Polymorphism, n	Gender, female, n (%)	Age, years, mean $\pm$ SD	Methods	Statistically significant difference?*	Main findings
<b>Tesli et al. (2013)</b>	<i>CACNA1C</i>	BD, 66 HC, 123	A/A+A/G, 34 G/G, 32 A/A+A/G, 71 G/G, 52	18 (52.9) 20 (62.5) 31 (43.7) 23 (44.2)	34.2 $\pm$ 10.0 35.5 $\pm$ 11.4 34.0 $\pm$ 7.8 35.3 $\pm$ 10.4	fMRI 1.5T Emotional stimuli (negative faces) paradigm ROI: Bilateral amygdala	<b>Yes</b>	Carriers of the risk allele had increased activation in the left amygdala in the BD group.
<b>Radua et al. (2013)</b>	<i>CACNA1C</i>	BD, 20 HC, 20	Rs1006737 GG, AG and AA Letter to the editor, N not reported	8 (40.0) 10 (50.0)	43.0 $\pm$ 14.0 42.0 $\pm$ 12.0	fMRI ( <i>Tesla not reported</i> ) Fearful faces ROIs: MFG, left putamen and left amygdala	<b>No††</b>	MTG out-degree connectivity gradually decreased with the number of <i>CACNA1C</i> risk alleles (GG<AG<AA) in BD and HC groups.
<b>Jogia et al. (2011)</b>	<i>CACNA1C</i>	BD, 41 HC, 50	GG, 17 AG, 19 AA, 5 GG, 28 AG, 18 AA, 4	21 (51.2) 23 (46.0)	44.3 $\pm$ 11.9 34.9 $\pm$ 13.2	fMRI 1.5T Facial affect recognition task (fearful vs neutral faces) ROIs: PFC, ACG, amygdala and hippocampus	<b>Yes</b>	Independent of diagnostic group, the right amygdala showed greater activation during fear-face recognition relative to neutral faces in AA/AG compared to GG individuals. The right vIPFC expressed reduced activation in individuals with the high-risk allele compared with those with the low-risk variant in BD patients.
<b>Dima et al. (2013)</b>	<i>CACNA1C</i>  <i>ANK3</i>	BD, 41 HC, 46 BD, 41 HC, 46	A/A+A/G, 17 G/G, 24 A/A+A/G, 25 G/G, 21 T/T+C/T, 16 C/C, 25 T/T+C/T, 14 C/C, 32	11 (64.7) 10 (41.6) 9 (36.0) 9 (42.8) 7 (43.7) 14 (56.0) 7 (50.0) 14 (43.7)	44.4 $\pm$ 12.3 44.1 $\pm$ 11.6 36.3 $\pm$ 10.4 38.1 $\pm$ 13.4 42.0 $\pm$ 10.7 43.3 $\pm$ 2.3 40.6 $\pm$ 12.2 39.3 $\pm$ 12.3	fMRI 1.5T Facial affect paradigm ROIs: IOG, FG, amygdala, vIPFC and whole-brain analysis	<b>Yes</b>	BD carriers of either genetic risk variant exhibited pronounced reduction in vIPFC activation compared to HCs.
<b>Barzman et al. (2014)</b>	<i>TNF</i>	BD, 10	<i>TNF</i> gene expression levels, 10	5 (50.0)	15.0 $\pm$ 1.0	fMRI 4.0T Posner Task	<b>Yes</b>	Expression of 11 <i>TNF</i> -related genes were significantly correlated with activation of amygdala or anterior CG

						(Selective attention task to negative emotions) Whole-brain analysis		during the affective Posner Task.
<b>Lelli-Chiesa et al. (2011)</b>	<i>COMT</i>	BD, 40 HC, 50	Val/Val, 11 Val/Met+Met/Met, 29 Val/Val, 15 Val/Met+Met/Met, 35	21 (52.5) 24 (48.0)	44.0 ± 11.9 34.9 ± 13.2	fMRI 1.5T Sad facial affect discrimination task ROIs: Amygdala and ventral PFC	<b>No††</b>	No significant diagnosis × genotype interaction was detected in the BD group.
<b>Liu et al. (2010)</b>	GWAS	BD, 39 HC, 29	rs2023454 SNP <i>DOK5</i> gene	24 18	15.0 ± 2.8 13.0 ± 2.9	fMRI 3.0T Face-processing paradigm ROI: Amygdala	<b>No†</b>	No significant main effect of diagnosis and no significant diagnosis × genotype interaction was detected. In both the BD and HC samples the rs2023454 SNP of the <i>DOK5</i> gene was significantly associated with right amygdala activation under the hostility contrast.
<b>Shah et al. (2009)</b>	<i>5-HTTLPR</i>	BD, 30 HC, 48	L/L, 10 S carriers, 20 L/L, 14 S carriers, 34	7 (70.0) 13 (65.0) 6 (42.8) 22 (64.7)	36.9 ± 16.9 29.4 ± 11.0 26.1 ± 9.3 29.1 ± 10.3	fMRI 3.0T Emotional face paradigm ROI: Ventral ACG	<b>Yes</b>	During fear and happy face processing, ventral ACG activation was significantly lower in the BD compared to the HC group, and in S carriers compared to L/L individuals within both HC and BD groups
<b>Dima et al. (2016)</b>	Polygenic risk score (GWAS)**	BD, 41 HC, 46	0.37 (0.04) 0.32 (0.06)	21 (51.2) 21 (45.7)	44.3 ± 11.9 40.3 ± 13.2	fMRI 1.5T Facial affect paradigm; 2-back task Whole-brain analysis	<b>No</b>	The PGR was associated with task-related changes in the BOLD signal in both the BD and HC groups. No significant correlation between the PGR score and the signal changes observed in BD patients during the facial affect paradigm was found (increased activation of ACG and decreased activation of the right superior frontal gyrus). In the 2-back task, no effect of group was noted and no correlation of the PGR score and group was observed in the task-related activation.

\* Comparisons between BD carriers of the risk allele and healthy controls or BD non-carriers of the risk allele. Significant interactions considering only clinical symptoms scores or medication use were not considered.

\*\* Polygenic risk score (PGR) is calculated in each individual by aggregating variation across GWAS loci nominally associated with BD into a quantitative score (Dima and Breen, 2015). The study used a PGR based on 16,691 SNPs with  $p < 0.1$ . It is presented as mean (SD).

† Sample size was considered a limitation in the original report

†† Sample size was not mentioned as a limitation in the original report

**Table 4. Studies investigating the association of genetic polymorphisms and brain activity in BD using functional magnetic resonance imaging (fMRI) during verbal and working memory tasks.**

Author	Gene	Subjects	Genetic Polymorphism, n	Gender, female, n (%)	Age, years, mean $\pm$ SD	Methods	Statistically significant difference?*	Main findings
<b>Delvecchio et al. (2015)</b>	<i>ANK3</i>	BD, 41	T/T+C/T, 16	7 (43.7)	42.0 $\pm$ 10.7	fMRI 1.5T	<b>Yes</b>	For the <i>ANK3</i> rs10994336, the risk T-allele was associated with increased activation in the right ACG and left PCG in BD patients compared to HCs. For the <i>ANK3</i> rs9804190, the risk C-allele homozygotes showed increased activation in right ACG in BD patients compared to HCs.
			C/C, 25 (Rs10994336)	14 (56.0)	43.3 $\pm$ 12.3	N-back task		
			C/C, 21	13 (61.9)	43.5 $\pm$ 12.5	Whole-brain analysis		
			T/T+C/T, 20 (Rs9810490)	9 (45.0)	44.8 $\pm$ 9.5			
		HC, 46	T/T+C/T, 14	7 (50.0)	40.6 $\pm$ 12.2			
			C/C, 32 (Rs10994336)	14 (43.7)	39.3 $\pm$ 12.3			
			C/C, 28	14 (50.0)	40.1 $\pm$ 13.2			
			T/T+C/T, 18 (Rs9810490)	8 (44.4)	40.1 $\pm$ 11.9			
<b>Prata et al. (2011)</b>	<i>DISC1</i>	BD, 35	Ser/Ser, 17	11 (61.1)	38.1 $\pm$ 13.3	fMRI 1.5T	<b>No</b>	No significant effect of Cys704Ser was detected.
			Cys Carriers, 18	10 (55.5)	40.9 $\pm$ 11.2	Verbal fluency test		
		HC, 53	Ser/Ser, 26	12 (46.1)	31.7 $\pm$ 11.1	ROIs: Left middle/superior frontal gyrus and whole-brain analysis		
			Cys Carriers, 27	15 (55.5)	37.8 $\pm$ 9.9			
<b>Chakirova et al. (2011)</b>	<i>DISC1</i>	BD, 36	T/T, 16	14 (38.8)	39.3 $\pm$ 10.8	fMRI 1.5T	<b>Yes</b>	Decreased activation in BD carriers of SNP rs821633 in the right IPL and left CG compared to non-carriers.
			C/C+C/T, 20			Verbal initiation and Sentence completion tasks		
		HC, 34	T/T, 16	25 (75.7)	37.3 $\pm$ 12.1	Whole-brain analysis		
			C/C+C/T, 18 (rs821633)					
<b>Mechelli et al. (2012)</b>	<i>DAOA</i> (rs746187)	BD, 33	A/A, 10	7 (70.0)	34.6 $\pm$ 13.1	fMRI	<b>Yes</b>	<i>DAOA</i> AA genotype was associated with greater deactivation (i.e. repetition > verbal fluency) during task performance than the AG/GG genotype in patients with
			A/G+G/G, 23	15 (65.2)	39.2 $\pm$ 11.7	1.5T		

	<i>DAAO</i> (rs2111902)	HC, 47	A/A, 22 A/G+G/G, 25 T/T, 27 T/G+G/G, 20 T/T, 16 T/G+G/G, 17	11 (50.0) 12 (48.0) 14 (51.8) 9 (45.0) 10 (62.5) 11 (64.7)	32.9 ± 8.2 34.8 ± 11.8 34.5 ± 10.3 33.2 ± 10.3 39.1 ± 11.0 38.5 ± 12.0	Verbal fluency paradigm Whole-brain analysis		BD, but not in HC in left precuneus. There were no regions showing a significant diagnosis by <i>DAAO</i> genotype interaction.
<b>Papagni et al. (2011)</b>	<i>DAAO</i>	BD, 33  HC, 48	C/C, 19 C/T+T/T, 14 C/C, 29 C/T+T/T, 19	11 (57.8) 9 (64.2) 14 (48.2) 10 (52.6)	40.1 ± 12.2 37.0 ± 11.9 34.9 ± 10.5 33.5 ± 10.9	fMRI 1.5T Verbal fluency paradigm Whole-brain analysis	<b>No</b>	No significant effects of diagnosis or of genotype in comparisons involving BD patients.
<b>Ham et al. (2016)</b>	<i>Bcll</i>	BD, 26  HC, 32	G/G, 11 C carriers, 15 G/G, 18 C carriers, 14	11 (100.0) 10 (66.6) 12 (66.6) 10 (71.4)	29.6 ± 8.5 33.6 ± 9.0 34.5 ± 4.7 32.5 ± 6.0	fMRI 3.0T Reward paradigm Whole-brain analysis	<b>No</b> †	No significant main effects of genotype, diagnosis or reward condition involving BD patients solely.
<b>Mechelli et al. (2008)</b>	<i>NRG1</i>	BD, 29  HC, 45	T/T, 16 C/T, 13 T/T, 25 C/T, 20	12 (75.0) 8 (61.5) 12 (48.0) 12 (60.0)	35.1 ± 12.6 41.1 ± 9.8 35.7 ± 9.7 34.6 ± 12.2	fMRI 1.5T Verbal fluency task Whole-brain analysis	<b>Yes</b>	The high-risk variant of <i>NRG1</i> was associated with greater deactivation in the left precuneus in both HC and BD. Right posterior OFC expressed increased activation in individuals with the high-risk variant in the BD group.

\* Comparisons between BD carriers of the risk allele and healthy controls or BD non-carriers of the risk allele. Significant interactions considering only clinical symptoms scores or medication use were not considered.

† Sample size was considered a methodological limitation in the original report.

†† Sample size was not mentioned as a limitation in the original report.



**Table 5. Summary of findings of reported genes and statistically significant differences between BD patients with high-risk genetic polymorphisms and healthy controls (HCs).**

Gene	Name	Function	Reported Polymorphisms	N of studies	Polymorphism with positive results	Method	Neuroimaging finding		Functional paradigm or DTI parameter used	
							Increased	Decreased		
<i>CACNA1C</i>	L-type calcium channel $\alpha_1C$ subunit	Voltage-dependent $Ca^{2+}$ channels rapidly increase intracellular $Ca^{2+}$ concentration after depolarization, initiating a host of responses, including neurotransmitter release and changes in gene expression (Gargus, 2009)	rs1006737, G to A	4	Rs1006737 (Risk allele A)	VBM		L Putamen (v)		
					Rs1006737 (Risk allele A)	fMRI	L Amygdala (f)		Emotional face task	
					Rs1006737 (Risk allele A)	fMRI	R Amygdala (f)	R vIPFC (f)		Emotional face task
					Rs1006737 (Risk allele A)	fMRI		R vIPFC / L vIPFC (f)		Emotional face task
<i>ANK3</i>	Ankyrin 3	Encodes ankyrin 3, a large protein involved in coordinated assembly of ion transporters and cell adhesion molecules at axon initial segments and nodes of Ranvier in myelinated nerves (Linke et al., 2011)	rs10994336 rs9804190 rs10761482	4	rs10761482 (cytosine [C] / thymine [T]; (risk allele C)	DTI		R Forceps minor / L Forceps minor	FA	
					rs9804190 (T carriers [risk])	DTI		R UF / L UF / R Dorsal ACG / L Dorsal ACG / L Temporoparietal region / L Dorsal CG / L Parietooccipital region / R Parietal region WM	FA	
					rs10994336 (risk allele T)	fMRI	R ACG / L PCG (f)		N-back task	

					rs9804190, risk allele C)	fMRI	R ACG (f)		N-back task
					rs10994336; risk allele T	fMRI		R viPFC/ L viPFC (f)	Emotional face task
<b><i>BDNF</i></b>	Brain-derived neurotrophic factor	Small protein of the neurotrophin family that regulates various brain functions; it has been implicated in modulation of hippocampal plasticity and hippocampal-dependent memory (Lu and Gottschalk, 2000)	Val66Met (rs6265)	3	Met carriers	VBM		R Hippocampus/ L Hippocampus	
					Met carriers	VBM		R Hippocampus/ L Hippocampus	
					Met carriers	VBM		R ACG/L ACG/ L dlPFC	
<b><i>5-HTTLPR</i></b>	Serotonin transporter polymorphism	Polymorphism in the upstream regulatory region of the gene – a 44-bp deletion/insertion ( <i>5-HTTLPR</i> ) located at the 5'-flanking regulatory region of the gene coding for the serotonin transporter (SLC6A4) on chromosome 17q11.2. In vitro studies evidenced that the basal activity of the long (l) variant was more than twice that of the short (s) form of the 5-HTTLPR, suggesting that serotonin transporter gene transcription is modulated by variants of the 5-HTTLPR with the s allele corresponding to low serotonin uptake activity (Heils et al., 1996)	Long (l) and short (s) variants (risk)	3	s carriers	VBM	R Amygdala (v)		
					s carriers	DTI	R PCG/ L ACG/CCb/CCg/ R posterior CR		RD and MD
					s carriers	fMRI		R Ventral ACG/ L Ventral ACG	Emotional face task
<b><i>NRG1</i></b>	Neuregulin-1	Encodes a family of signaling proteins in various tissues of the body with <i>NRG1</i> expression being highest in the brain. In the nervous system, <i>NRG1</i> proteins have been implicated in numerous functions, including neuronal migration, synapse formation and receptor expression as well as myelination by regulating	SNP8NRG243177 (rs6994992) SNP8NRG221533 (rs35753505)	2	SNP8NRG221533 (rs35753505) C carriers (high risk)	VBM	L CG/ L PHG/CCs/ACC		
					SNP8NRG221533 (rs35753505) C carriers (high risk)	fMRI	R posterior OFC		Verbal fluency task

		oligodendrocyte proliferation and differentiation (Winterer et al., 2008)							
<b>G72</b>	D-amino acid oxidase activator	<i>G72</i> has been associated with modulation of NMDA receptor function and with regulation of mitochondrial function and dendritic branching (Zuliani et al., 2009)	SNPs M23C/T and M24A/T	2	M23 and M24 T carriers	VBM		L TP/ R Amygdala	
			SNP rs746187A/G		SNP rs746187 A/A	fMRI		L Precuneus	Verbal fluency

<b>DISC1</b>	Disrupted-in-schizophrenia 1	Expressed predominantly within the hippocampus and codes for a protein with a globular N-terminus domain, a coiled C-terminus domain, and several coiled-coil domains. The functional role of DISC1 is largely unknown, but these distinct domains allow DISC1 protein to interact with both centrosomal and cytoskeletal proteins as well as with membrane associated and signal transduction proteins (Callicott et al., 2005)	Various risk variants (rs1538979, rs821577, rs821633, rs821616 [Ser704Cys], rs6675281 [Leu607Phe] and rs1411771) (Chakirova et al., 2011)	1	rs821633, allele C	risk fMRI		R L CG	IPL/ Verbal initiation and Sentence completion tasks
<b>DGKH</b>	Diacylglycerol kinase eta (DGK $\eta$ )	The DGK $\eta$ enzyme plays an important role in the inositol triphosphate second messenger pathway by catalyzing the metabolism of diacylglycerol (DAG) to phosphatidic acid. DAG is an activator of many isoforms of protein kinase C (PKC). Therefore, <i>DGKH</i> regulates the activity of PKC isoforms which play a key role in various signaling pathways (Kittel-Schneider et al., 2015)	Risk haplotypes (rs9315885, rs1012053, rs1170191, TAC), risk haplotype (rs994856/rs9525580/rs9525584 GAT) and/or risk polymorphisms in <i>DGKH</i> (rs994856, rs9525580, rs9525584, rs9315885)	1	rs994856/ rs9525580/rs9525584 584 GAT (risk haplotype)	VBM	L Amygdala		
<b>HAP1CE</b>	NRG1 HAPICE (deCODE) haplotype	It is a core haplotype of NRG1 consisting of five SNPs and two microsatellites (Cannon et al., 2012)	Arh0 (no copies of the haplotype): no risk Arh1 (1 or 2 copies of the haplotype): risk	1	Arh1: risk	VBM	R L L PCG	Fornix/ Caudate/	
<b>IL-1<math>\beta</math></b>	Interleukin-1 beta	Encodes for interleukin-1 beta (IL-1 $\beta$ ), pro-inflammatory cytokine which has an important role in the induction of the dopaminergic phenotype in mesencephalic neuronal precursors as well as in the regulation of dendrite growth in developing cortical neurons (Papiol et al., 2008)	--511 AvaI polymorphic site (rs16944) of <i>IL-1B</i> gene. Allele*1 (511C) of <i>IL-1B</i> gene completes an AvaI restriction site, while allele*2 (511T) gives an intact product	1	Allele*2 carriers	VBM	L dIPFC		
<b>GRIN2B</b>	NMDA receptor subunit 2B	Encodes the NR2B subunit of the NMDA glutamate receptor. This subunit is expressed in the cortical and medial temporal parts of the brain, striatum, and olfactory bulb	Risk variant rs890 G/T	1	T allele	DTI	R L L L R	Frontal region/ Frontal region/ Parietal region/ Occipital region/ Occipital region/	FA

(Kuswanto et al., 2013)

L CG

<i>TNF</i>	Tumor necrosis factor	Encodes Tumor necrosis factor-alpha (TNF $\alpha$ ), a cytokine involved in both systemic and neuro-inflammation and in the acute phase reaction, and may influence neuronal and neurochemical processes associated with aggression in preclinical and clinical studies (Barzman et al., 2014)	<i>TNF</i> family genes 1 expression (Barzman et al., 2014)	<i>TNF</i> family genes expression levels (11)	fMRI	R L L amygdala	ACG/ ACG/	Posner Task (Frustrative non-reward task)
------------	-----------------------	--	---	--	------	----------------------	--------------	---