



Københavns Universitet



Guide to handling of tropical and subtropical forest seed

Schmidt, Lars Holger

Publication date:
2000

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Schmidt, L. H. (2000). *Guide to handling of tropical and subtropical forest seed*. Danida Forest Seed Centre.

SEED DOCUMENTATION

Contents	14.1 Introduction	1
	14.2 Documentation and Certification	3
	14.3 Codes and Accession Number System	3
	14.4 Seed Documentation Systems	5
	14.4.1 Seed source records	6
	14.4.2 Collection and seed handling records	7
	14.4.3 Seed testing records	8
	14.4.4 Seed stock and dispatch records	8
	14.4.5 Customer information	9
	14.5 Updating and Maintenance of Data	9
	14.6 Data Management Systems	10
	14.6.1 Manual systems	10
	14.6.2 Computer database systems	11
	REFERENCES	12
	APPENDICES	13
	Appendix A14.1	
	Forms and registers for seed documentation	15
	Appendix A14.2	
	Examples of seed documentation forms used by seed centres	27
	Appendix A14.3	
	Examples of screen pictures and data entry forms in a computerized seed documentation system	32

Other Chapters of the book Guide to Handling of Tropical and Sub-Tropical Forest Seed by Lars Schmidt soon available on www.dfsc.dk

- Chapter 1: Introduction
- Chapter 2: Seed Biology, Development and Ecology
- Chapter 3: Planning and Preparation of Seed Collections
- Chapter 4: Seed Collection
- Chapter 5: Fruit and Seed Handling between Collection and Processing
- Chapter 6: Seed Processing
- Chapter 7: Phytosanitary Problems and Seed Treatment
- Chapter 8: Seed Storage
- Chapter 9: Dormancy and Pretreatment
- Chapter 10: Germination and Seedling Establishment
- Chapter 11: Seed Testing
- Chapter 12: Genetic Implications of Seed Handling
- Chapter 13: Microsymbiont Management
- Chapter 15: Trade and Transfer of Forest Seed

SEED DOCUMENTATION

14.1 Introduction

High physiological and genetic seed quality is tightly linked to a good record and documentation system. From a genetic point of view, seed without information on its origin, history and potential performance poses a risk and may be almost worthless. Seed records have different purposes for the different people involved, and the documentation system and contents including the level of detail vary accordingly. For example:

1. Seed documentation may be used as a planning tool for future seed procurement. Records on seed collection from various seed sources may provide useful information on productivity, seasonality and periodicity in seeding pattern, thereby saving time on initial surveys prior to collection. Success or failure of achieving good seed quality may often be ascribed to particular handling procedures. Records on seed handling correlated with seed quality may thus indicate sensitive aspects in the handling process and suggest improved procedures. Far too much re-invention and re-exploration is made because previous mistakes were not recorded. Poulsen and Thomsen (1999) have proposed a thorough seed procurement record and documentation system with the major purpose of improving procurement methods by analysing documented practices. The system covers all handling aspects from maturity criteria, collection and processing to testing and storage.
2. Internal database. In small units such as local forest departments, where seed collection, nursery operation and planting are under the same head, documentation may seem redundant. However, human memory has limitations, staff members move or resign, and the life time of trees, which can be important when tracing seed sources, tends to be longer than service in the forest sector.
3. Information and quality documentation for end user (customer). Documentation is a link between supplier and end user. It contains information on both seed source (origin) and test results. Good seed documentation becomes particularly important when seeds are dispatched to end users who are not in direct contact with those who collected

and handled the seed. This is particularly so in international transfer of seed (see chapter 15). Seed source details, genetic base and physiological quality are factors of major importance for the seed buyer.

4. Requirement from official authorities. Type and details of required seed documentation vary from one country to another and usually depend on quantity, type of disposal, and species. Small non-commercial seed lots distributed within the country e.g. in connection with research are rarely subject to any official requirement of documentation. Commercial seed lots are usually subject to the same type of accounting as any other commercial product. Seed transferred over international borders may be subject to particular regulations and documentation requirements. The topic is discussed in chapter 15; a few points should be summarised:
 - a. Customs regulations. Normally only in importing countries. Commercial seed lots may be subject to general or specific import regulations e.g. on paying tax and duty, and official invoices are required.
 - b. The importing country may require that imported 'living plant material' is free of pest and diseases. Where an official accredited laboratory carries out the examination and/or treatment, the seed lot is issued with an official 'Phytosanitary Certificate'.
 - c. Export of forest seed has so far rarely been subject to regulations by the exporting countries. However, with the increasing concern of national right to genetic resources, export regulations are likely to become more common in the future, at least for some species and in some countries.

Seed suppliers usually design their own forms and documentation system. Examples of a number of forms are presented in the appendices to this chapter. It should be emphasised that both too much and too little information may be a nuisance. It is important that all relevant information on a seed lot is available, but not all information need to be provided to any user. For example, a label adhering to a submitted seed sample need only contain basic information on species, provenance, quantity etc., while additional information is submitted on a separate form. It is therefore important that each seed lot be allocated a unique seed identity number, which will appear on any information form of the same seed lot, and from which specific seed lot information can be retrieved.

Seed documentation is usually filled in on pre-designed forms during collection, processing or testing. However, several computer programmes are available which facilitate record keeping and documentation. Specially designed integrated systems allow all kinds of information to be stored and retrieved in the same system. Database systems are continuously under development, and electronic storage and data management systems are still becoming more common.

14.2 Documentation and Certification

Reliability of documentation is always a matter of confidence. If a seed collector states that they have collected seeds from 20 widely spaced, good phenotype trees in a superior stand, there is rarely a way to check whether it is true. Unfortunately, much seed is traded above its real quality on that account. A good documentation system is a way to avoid lapse, failures or direct dishonesty in seed procurement, and in turn achieve confidence from customers. The signature by the officer in charge of individual operations should be a guarantee of both appropriate methodology applied (e.g. in testing) and reliability of data. Certification is a further guarantee issued by an official, governmental accredited authority, e.g. a seed testing station. Certification does not necessarily mean that the seeds have a higher quality than those with more informal documentation, only that quality or authenticity is verified by an official authority. Therefore, certification is generally used only for special valuable seeds or in connection with special requirements, e.g.:

1. Genetic quality. Seed collected from trees of proven genetic superiority, e.g. from seed orchards or superior ('plus') trees in a natural stand with controlled pollination. Certificates may state 'Certificate of Origin' or seeds may simply be referred to as 'certified tree seed' (For. Com. 1994).
2. Physiological quality. Where seed testing is carried out according to the ISTA (or AOSA) rules (see chapter 8) by special accredited seed testing laboratories, an ISTA certificate of seed quality may be issued. An ISTA seed quality certificate is hence an assurance that seeds have been tested according to the rules.
3. Health and diseases. Governments often require imported seeds to have been certified by an official authority, stating that the seeds are not infected or carrying diseases. Such a 'Phytosanitary Certificate' states that seeds have been examined and 'to the best of our knowledge' been found free of special pests and fungi.

Some examples of official certificates are presented in appendix A14.2.

14.3 Codes and Accession Number System

The purpose of a seed coding and numbering system is to create a quick reference system across all handling procedures. It saves both time and space in writing when a species, seed source or seed lot is referred to with few letters or reference figures rather than writing a full detailed explanation every time it is encountered. Codes and accession numbers (in some systems called 'reference' or 'identity' numbers) make the use of computerised database management possible, and allow e.g. lots of information to be compressed into column form, and hence several columns on a summary data sheet (table A14.2C). In practical seed handling, a seed source and seed-lot reference or accession number is used. Species codes are used mainly in computer-based documentation. In seed trading, supplier and customer codes are normally used in connection with a full seed documentation system (Lauridsen 1994). Seed suppliers would often

create their individual system of documentation and data management, but some initial considerations should be made:

Codes and reference numbers should be unique, meaning that two species, two seed sources or two seed lots do not have the same code or number. Species may be referred to by three or four first letters of genus and species name, all in capital letters. Three is usually sufficient both to be unique and to recognise the species name from the code, but it becomes easier with four. However, even four is not a safe identification code. For example EUCAMICR may mean both *Eucalyptus microtheca* and *E. microcorys*. If one happens to have both these species in supply, they must be distinguished by other means. In fact there are very few examples where 4+4 letters can cause confusion.

Seed-source reference number may consist of numbers, letters or a combination. Letters may refer to species, location, type of seed-source etc. Letters referring to species may cause confusion in case several species are collected from the same seed source. As an example of a seed source reference system, the National Tree Seed Programme (NTSP) in Tanzania uses a system consisting of three components (Rasmussen 1992):

- two capital letters, referring to one of the three regional seed centres (RC) responsible for the seed source.
- a three digit serial number referring to seed source identity within the particular RC.
- one capital letter referring to type of seed source e.g. Z (= seed collection zone), I (= identified stand), S (= selected stand), A (= seed production area), P (= provenance seed stand), and O (= seed orchard).

For example, seed source number MO149S thus refers to a seed source located in the area under the Morogoro RC, it is number 149 of their seed sources, and is a selected stand.

Seeds delivered at the central processing station (forest office, seed centre etc.) may already have been given a provisional or temporary seed-lot number for identification in the field. At the beginning of the handling procedure the seed lot is given a new seed-lot number according to a pre-decided system, which may differ from place to place. As with seed source the identification or accession number may contain letters, figures or both, which may be related to species, seed source, year of collection etc. (Lauridsen 1994).

Australian Tree Seed Centre uses a continuous 5 digit number for each species (ATSC 1995 (see table A14.2C)). Danida Forest Seed Centre uses a system where seed lots are numbered in sequence as they enter the seed bank, the year of accession being indicated. A seed lot with identity number 5320/92 means a bulked seed lot number 5320 received in 1992 (Lauridsen 1994). At the NTSP in Tanzania the seed-lot accession number consists of three components: 1) the seed source number (according to the system described above), 2) a

two digit number indicating year of collection, and 3) a capital letter indicating number of collection in the particular seed source. Seed-lot accession number MO149S/91B thus refers to the second seed collection (indicated by B) in 1991 (indicated by /91) from above mentioned Morogoro seed source number 149, which is a selected stand (Rasmussen 1992).

As the above examples show, the options are vast and the individual seed supplier must decide on a system, which is appropriate under the particular operation conditions. The most important is that identity or reference numbers are unique, so that two seed sources or seed lots will not have the same number. Once the seed lot number has been allocated, it should be entered on all seed forms and labels subsequently used for that seed lot.

Codes and reference numbers for seed suppliers and customers may include letters for country code, region, project etc.

14.4 Seed Documentation Systems

Complete seed documentation systems consist of a variable number of standard forms, which are filled in during seed handling. Examples of seed documentation systems are given by Bowen (1980), Rasmussen (1992), Lauridsen (1994), and ATSC (1995). Willan (1985) gives examples of forms used by forest seed centres and suppliers all over the world. A compilation and modification of some of these forms appears in appendix A14.1 of this chapter. Additional examples of filled in forms used by various seed centres are shown in appendix A14.2. Although computer software has replaced much of the cumbersome manual recording and registering, predesigned forms remain essential for recording in the field or laboratory. Information and records filled in forms may be transferred to computer systems to facilitate both statistical calculation and other data management. Computer data management will be further discussed in section 14.6. A discussion of some of the components of the seed documentation forms of appendix A14.1 is given in section 14.4.1 to 14.4.5 below. An overview of the forms is presented in table 14.1.

Table 14.1. Overview over seed record and documentation forms. Figures in parentheses refer to appendix A14.1. (compare also table 14.3).

Seed source records	Seed source information form (A14.1A) List of seed sources (A14.1B) Country map of seed sources (A14.1C)
Collection and seed handling records	Collection and handling records (A14.1D) Seed procurement accounting form (A14.1E)
Seed testing records	Seed sampling form (A14.1F) Seed test forms (A14.1G1+2)
Seed stock and dispatch records	Seed stock and dispatch form (14.1H)
Customer information	Seed lot summary data sheet (A14.1I) Labels for containers (A14.1J)

14.4.1 Seed source records

Seed source information is recorded on forms like the example in appendix A14.1A. The records provide information on the origin of the seed, i.e. from where the seeds were collected including the ecological conditions of the site and type of seed source. Ecological information such as altitude, rainfall, temperature, and soil type, is essential for matching the right seed source with the particular planting environment. Also seed zone and provenance contain information about site conditions. Seed zones are the largest units in seed collection. The classification of seed zones is based on climatic factors (temperature, precipitation), physiographic structure (topography, geology, soils) and geographical elements (vegetation). It is envisaged that the genetic variation within a seed zone is less than between two different zones (Albrecht 1993). Seed zones serve as a broad guideline for transferring seeds for national plantation programmes (Barner and Willan 1985). The provenance name refers more specifically to where the species grows. A seed zone (sometimes called region of provenance) may thus contain several identified provenances. A provenance name normally refers to a relatively well identifiable place on a map, e.g. a lake, hill or nearest larger town. The provenance concept (see chapter 3) implies that there is a reasonable uniformity and exchange of genetic material within the population that makes up a provenance. Hence, provenance contains two types of information 1. ecological and 2. genetic uniformity.

The ecological information of provenances may be a more specific guideline on provenance choice for particular planting sites than seed zone. For example, if one were to plant *Pinus caribaea* at a lowland moist site, it would probably be advisable to purchase seed from a lowland moist provenance. The provenance name refers to where the species grows, no matter whether it is a natural forest or it was introduced. However, for exotic species the origin should, if possible be indicated. If a seed source in Salima, Malawi, was established from a Petford provenance of *Eucalyptus camaldulensis*, it is a Salima provenance, but the origin of Petford is indicated.

The type of stand gives information on the genetic history and sometimes quality of the seed source. Neither natural stands nor most plantations have been subject to selection for quality. A natural stand often has a wider genetic base than a plantation unless the natural stand is a small isolated group of trees, e.g. a small relict of a previous large stand. Both natural stands and plantations can be upgraded to selected or certified seed sources or seed stands. The latter involves some selection since inferior individuals must be eliminated to upgrade the average genetic quality. Where a seed source has been established based on genetic results from an improvement programme, the seed source is referred to as a seed orchard. Both generation (1st, 2nd etc.) and mode of establishment (clonal or seedling) are indicated in the seed source record (see also discussion on seed sources in chapter 3).

Much information on seed-source forms has only relevance to the seed collector or supplier. That pertains to e.g. information on ownership, accessibility, size, age and productivity of the stand. Also details

on flowering and fruiting periods, potential local labour availability and other pure collection details are of little interest for the seed customer, while being highly relevant for the seed supplier. Only information that refers directly or indirectly to seed quality is transferred to the form submitted to the customer (appendix A14.1I).

Seed sources are conveniently listed in an easily overviewed table (A14.1B). Listing may be done according to species or seed source. The latter is sometimes more appropriate where seed sources contain and serve as source for more species e.g. many mixed natural forests. Computer databases can easily shift between seed source listing and species listing. Eventually, seed sources are conveniently plotted on a national or regional map indicating seed-source reference number (appendix A14.1C).

14.4.2 Collection and seed handling records

For each individual seed lot, procedures and conditions during seed collection and processing are recorded (A14.1D). As the seed-source reference number appears in the form, collection and handling records can be linked to the particular seed source where the seed was collected.

The collection form contains two measures on genetic quality viz. genetic base and selection of seed trees. Genetic base is indicated by the number of and distance between parent trees of the seed lot. This information is important if the plants to be raised may be considered as seed source some time in the future (see discussion on genetic base, chapter 3). Information on phenotypic selection of seed trees applies in particular to natural stands and plantations, since selection is already implicit where seeds are collected from seed stands or seed orchards. Phenotypic selection of seed trees is usually done during collection (section 3.3).

Information on field handling, transport and possible field processing are particularly important for the further processing at the central processing station. It is envisaged that the first part of the seed-handling and processing form is filled in and signed by the person responsible for the collection, before the seed lot incl. form is handed over to the processing station.

Many details on collection and handling appearing on the form have relevance only to those involved in seed collection and processing. The information may be used by the seed supplier as a guideline during the succeeding processing of the particular seed lot, and to improve procedures for future seed lots. Seed-handling records are especially pertinent where different people are involved in different processes. For example, if field collection records tell that a seed lot was collected during moist weather, the processing officer may give priority to rapid processing of this seed lot. It is therefore an advantage that the seed-collection and handling form follows the seed lot during seed handling and is filled in gradually during handling (Poulsen and Thomsen 1999), see also appendix A6.2.

Expenses of seed collection and processing are recorded on form

14.4.3 Seed testing records

A14.1E. The accounting system helps to estimate the actual procurement expenses, which form a major component during calculation of seed price (chapter 15). This information may also be used to identify areas where gains in efficiency can be made. It is also useful during planning when demand is matched to potential sites for collection.

Samples for seed testing are frequently drawn from the bulk seed lot and submitted to the seed-testing laboratory. A seed sampling form (A14.1F) should accompany the samples. The form is a documentation that sampling was undertaken according to the rules of representativeness as stated in section 11.4.

Seed testing documentation contains results of standard seed testing procedures such as seed weight, moisture content, purity and viability/germination plus sometimes special test (chapter 11). For laboratory tests special working forms are used (form A14.1G1 and A14.1G2). The results are summarised in a seed testing section of the seed-lot summary data form (A14.1I). If other viability test than germination is carried out (cutting, X-ray, TTZ etc.), the form should state which type of test has been carried out. Since viability declines with time, the date of testing is important. Regular viability tests are pertinent for long-time storage, and for seed with short viability. Obviously the latest record should be the one submitted to customers.

Testing for seed-borne pests and pathogens may be carried out in connection with general seed testing or by special laboratories. The latter is often a requirement during international transfer, where the phytosanitary condition must be documented on a Phytosanitary Certificate (A15.2). A phytosanitary certificate always states that the seed is found free of special pathogens, since otherwise it would not be issued. Often it is issued only after treatment of the seed. The document will then state which type of pesticide has been used. This should also always be indicated on more informal seed documentation forms.

14.4.4 Seed stock and dispatch records

These records are essential for management of seed stores, sales or other modes of disposal. In addition the seed-stock records contain the basic information to be used in seed catalogues (chapter 15).

The seed stock and dispatch record contains a list of species, provenance, seed lot number, date of collection etc. (appendix A14.1H). In large seed stores it may be advantageous to indicate also the storage location in order to find seeds quickly. Seed stock currently changes as new seed lots are collected and enter into the stock, and seed lots are dispatched for sale or local nursery sowing. It is essential that both outgoing and incoming seed lots be consciously registered to keep trace of the seed stock. Where seed of different age is available in the stock, the 'first-in - first-out' principle is adopted. Data of incoming and outgoing stock are essential also for planning future seed collection.

Computer databases are particularly useful for stock and dispatch records. Even where the documentation system is mainly manual, tables formed in word processing programmes may be useful for record keeping since new lines can be added as new seed lots appear or parts of seed lots are dispatched.

14.4.5 Customer information

Labels are attached to the seed sample submitted to the customer. The label contains only basic information such as species, seed lot number, quantity etc. as indicated in form A14.1J. Submitted seed lots may be accompanied by copies of seed handling and testing forms, or information may be compiled into a separate summary data form (A14.1I). In addition to the essential seed lot information, the summary form contains brief recommendations on handling from receipt to sowing.

14.5 Updating and Maintenance of Data

Data management consists of collecting, storing and distribution of data. As any new seed lot involves several forms and registers, one would soon end up with piles of paper, if it were all kept in files. Much information may be relevant only to few people and for short time. Both ledger systems and computer databases tend to accumulate vast numbers of outdated data, and a system of erasing irrelevant data soon becomes an important task in data management. In general, information should only be kept as long as it is needed, and distributed only to those for whom it is relevant. Supplier as well as customer must consider which type of information need to be maintained and what can be eliminated. Seed testing records would, for instance, normally be of no use once the seed has been sown. Of utmost importance is that the seed origin of any plantation can be traced back to its seed source. If a seed lot happens to give rise to an exceptionally good (or bad!) performance, it is important that the customer (and supplier) is able to trace the source from where the seeds were collected.

Generally, seed-source information is kept as long as the (potential) seed source is there, whether seed is still collected from there or not. However, current information is obviously only updated in those seed sources regularly used. The seed collector must be aware of this since e.g. changed ownership or administration may necessitate new arrangements on collection. If appropriate measures were taken during collection to assure the genetic quality (number, spacing and performance of mother trees), and the seed supplier is likely to keep seed source records, then information on seed source may be redundant for the tree planter. However, in practice such information tends to get lost for various reasons during the lifetime of a forest plantation. Therefore it is advisable that the tree planter keeps essential seed-source information that will enable him/her to trace the seed source.

For the seed supplier, information on a particular seed lot becomes irrelevant once the last seed has been dispatched. In some computer database systems a seedlot will automatically be deleted if its final

value is nil. However, the seed supplier should keep records on dispatched seed lots, both as a service to the customer in case some records are lost, and in their own interest since it will help to identify particularly valuable (or poor!) seed sources and provenances.

A simple system of records is shown in table 14.2.

Table 14.2.

Example of seed records maintained by the seed supplier (upper) and customer (lower). If plantation E.Rift/2 happens to show excellent performance, the customer (UNHCR/34) will be able to trace the seed lot (AM/1838/97) from the supplier (NTSP/IG), which in turn will be able to trace the seed lot AM/1838/97 back to seed source IN/281.

Suppliers records

Species	Seed lot	Provenance	Customer	Seed source
ACACMANG	AM/1838/97		UNHCR/34	IN/281
ACACMANG	AM/1839/97		UNHCH/34	IN/281
ACACMANG	AM/1840/97		SALI/02	IN/430
ACACMANG	AM/1841/97		SALI/02	IN/435

Customers records

Species	Plantation	Seed lot acc. no.	Provenance	Supplier code
ACACMANG	E.Rift/2	AM/1838/97		NTSP/IG
ACACMANG	E.Rift/3	AM/1838/97		NTSP/IG
ACACMANG	W.Rift/1	AL/0018		NTSP/IG
ACACMANG	W.Rift/3	AL/1112		NTSP/IG

14.6 Data Management Systems

Data relating to seed sources, seed lots, sales etc. can be handled entirely manually by filling in, filing, copying and submitting forms like the ones shown in appendix A14.1. More extensive use of personal computers and development of their storage and operation capacity, plus development of user-friendly database programmes are nowadays replacing much of the laborious copying work and manual filing system. Databases can be used for statistical analysis and searching, and data storage requires much less space than manual systems. Establishment of computer networks allows quick and efficient entry and reading of data from linked computer terminals. It is therefore especially suitable for handling large numbers of seed lots. Computer systems do not, however, entirely replace manual systems, since forms must often be filled in the field or in the laboratory without access to terminals. An entirely electronic data management system is sensitive to power cuts and computer breakdown and a combination of manual and electronic systems are often preferred.

14.6.1 Manual systems

Most manual data management systems use a set of standard formula like the ones presented in this chapter. Permanent records such as seed source or plantation data were previously almost exclusively filed in ledgers, a system that is still much used. The ledger system is robust and simple but quite time consuming.

The viscard system is a loose-leaf filing system using hard index cards arranged according to a pre-decided system e.g. species or seed lots. The cards are filed in boxes, arranged so that one or two data lines containing key information is visible without taking out the whole card. Because the cards are relatively small (and expensive), the viscard system is used mainly for particular information like address catalogues, seed-stock register, seed-testing register or the like (Lauridsen 1994).

Loose-leaf systems are more flexible than the ledger system and more practical for filing. Original or photocopy of standard forms is normally filed in ring binders according to different systems e.g. species, seed lot, customer, supplier or other system.

14.6.2 Computer database systems

Documentation may be filed in standard forms in ordinary word processing systems, but with much data, the system becomes awkward. Simple databases may be set up in DOS based programmes like Dbase and Paradox. Newer databases based on Windows systems, e.g. Access, allow statistical analysis and combination of data from different registers. A complete seed documentation database system has been established for an Indonesian tree seed project (Purmono 1996). Examples of data entry screens for seed source and seed lot respectively are shown in figure appendix A14.3.

A database consists of a number of registers, each containing one or more key fields plus a number of data fields. Key fields are used for searching information and combing information from different registers. Examples of key fields are species codes, seed source accession code, seed lot accession code, supplier code etc. The same key fields may appear in different registers. Data fields contain specific information related to each register. For example, a seed source register contains data fields of provenance name, locality, ownership etc. Data fields are unique for each register. For a full seed procurement programme, the registers shown in table 14.3 may be needed.

Table 14.3. Seed registers for a database system on seed procurement including key fields and data fields for each register. Slightly modified from Lauridsen (1994). Note that the data fields contain the same type of information as listed in the forms in appendix A14.1.

Register	Key fields	Data fields
Species register	Species code	Species name (latin) incl. author, synonyms, common name
Seed source register	Seed source code/reference no., species code	Provenance name, locality details, owner, collection regulation, collection fee etc.
Seed supplier register	Supplier code	Name of supplier, address, phone etc.
Seed collection data register	Seed lot identity no., seed source code/reference number, species code, (supplier code)	All data obtained through collection and processing (except data contained in the seed source register)
Seed quality register	Seed lot identity no.	Main results from the seed test data forms
Seed stock register	Seed lot identity no.	Seed quantity received, seed customer code, date reserved/ordered, date dispatched, weight reserved/dispatched
Seed customer register	Seed customer code	Name of customer, address, phone etc.

REFERENCES

- Albrecht, J. 1993. Forest seed handling. In: Tropical Forestry Handbook, Vol. 1. (Pancel, L., ed.). Springer Verlag. pp. 381-462.
- ATSC 1995. Operations manual. Australian Tree Seed Centre.
- Barner, H. and Willan, R.L. 1985. Seed collection units: 1. Seed zones. Technical Notes No. 16. Danida Forest Seed Centre. Denmark.
- Bowen, M.R. 1980. Recording procedures for the collection, cleaning and drying (extraction), testing (certification), storage and issue of forest tree seeds. Working Paper No. 4, FAO/UNDP/MAL/78/009. Forest Research Centre (Sepilok), Malaysia.
- For. Com. 1994 Seed manual. Procedures for seed collection, handling, storage, purchase and accounting. Forestry Commission, Hobart, Tasmania.
- Lauridsen, E.B. 1994. Seed documentation. Lecture Note C-13. Danida Forest Seed Centre. Denmark.
- Poulsen, K. and Thomsen, K. 1999. Seed Handling Manual. Technical Note 54. Danida Forest Seed Centre. Denmark.
- Purnomo, H. 1996. Seed information system (SIS). Ministry of Forestry, Directorate General of Reforestation and Land Rehabilitation, Indonesia Tree Seed Source Development Project. Danagro Adviser A/S.
- Rasmussen, P. 1992. A guide to the NTSP seed documentation system, forms, routines, files, source and accession numbers. National Tree Seed Project, Morogoro, Tanzania.
- Willan, R.L. 1985. A guide to forest seed handling. FAO Forestry Paper 20/2. FAO, Rome.

APPENDICES

- Appendix A14.1 Forms and registers for seed documentation
- A14.1A: Seed source information form
- A14.1B: List of seed sources
- A14.1C: Country map of seed sources
- A14.1D: Collection and handling records
- A14.1E: Seed procurement accounting form
- A14.1F: Seed sampling form
- A14.1G: Seed test forms
- A14.1H: Seed stock and dispatch form
- A14.1I: Seed lot summary data sheet
- A14.1J: Seed labels
- Appendix A14.2. Examples of seed documentation forms used by seed centres
- A14.2A: OECD certificate of seed source
- A14.2B: Collection and handling form, Kenya Forestry Seed Centre
- A14.2C: Seed stock register, Australian Tree Seed Centre
- A14.2D: Germination test record form, Australian Tree Seed Centre
- A14.2E: Summary seed record form, Australian Tree Seed Centre
- Appendix A14.3. Examples of screen pictures and data entry forms in a computerised seed documentation system

Appendix A14.1A

SEED SOURCE INFORMATION FORM

Species information

Seed source reference No. _____	Seed zone _____
Provenance name _____	
Species (botanical): _____	Species code: _____
Common name: _____	

Seed source classification:

- Unclassified,
 Seed zone,
 Identified stand,
 Selected stand,
 Seed production area,
 Provenance seed stand,
 Seed orchard

Location description

Seed source location: _____	District/county _____
Region/state _____	Country _____
Geographical coordinates: Latitude: ° 'N/S , longitude: ° 'E/W	
Altitude: m.a.s.l	
Koeppen climatic code*: <input type="checkbox"/> Af, <input type="checkbox"/> Am , <input type="checkbox"/> Aw, <input type="checkbox"/> Aw1, <input type="checkbox"/> Bsh, <input type="checkbox"/> Bsk, <input type="checkbox"/> Bwh, <input type="checkbox"/> Bwk, <input type="checkbox"/> Cfa, <input type="checkbox"/> Cfb, <input type="checkbox"/> Cw, <input type="checkbox"/> Cw1, <input type="checkbox"/> Cs, <input type="checkbox"/> D	
Rainfall regime: <input type="checkbox"/> Summer, <input type="checkbox"/> Uniform, <input type="checkbox"/> Winter, <input type="checkbox"/> Bimodal	
Mean annual rainfall (mm): _____	
Length of dry season (<60mm) (indicate months): _____	
Mean annual temperature (°C): _____	
Mean daily min. temp coldest month (°C): _____, Mean daily max. temp hottest month (°C): _____	
Absolute min. temperature (°C): _____	
Other information: _____	

Site description

Terrain : <input type="checkbox"/> Flat, <input type="checkbox"/> Hilly, <input type="checkbox"/> Mountainous, <input type="checkbox"/> Ridge top	
Slope: <input type="checkbox"/> Flat or gentle (<5%), <input type="checkbox"/> Intermediate (5-10%), <input type="checkbox"/> Steep(11-45%) , <input type="checkbox"/> Sheer	
Aspect: <input type="checkbox"/> North, <input type="checkbox"/> East, <input type="checkbox"/> South, <input type="checkbox"/> West, <input type="checkbox"/> Level	
Soil type: _____	pH: _____

Stand description

Total area: hectares
Type of stand: <input type="checkbox"/> Unknown <input type="checkbox"/> Natural stand <input type="checkbox"/> Plantation, Planted year _____
<input type="checkbox"/> One species <input type="checkbox"/> Mixed species, Associated species: _____

Accessibility

Distance to nearest forest station:

Accessibility road, 2WD, 4WD,

Walking distance from nearest road accessible by 4WD _____

Collection permit: Required Not required

Seed production

Flowering period _____

Fruiting period _____

Harvestable fruit production (estimated): _____ Kg, or seed production (estimated): _____ Kg

Labour availability

Name(s) of nearest village: _____

Distance from seed source to nearest village: _____

Available labourers: _____

Other Information:

* Koeppen climatic codes

Af: Permanent humid

Am: Monsoonal, short dry season

Aw: Subhumid, drier than Am

Aw1: As Aw but bimodal rainfall

Bsh: Semi arid, hot "steppe", Sahel"

Bsk: Semi arid, warm to cold

Bwh: Arid, hot desert

Bwk: Arid, warm to cold

Cfa: Humid subtropics, east side of continents, incl. montane.

Cfb: Temperate, maritime

Cw: Highland, subhumid

Cw1: As Cw but bimodal rainfall

Cs: Mediterranean.

D: Temperate continental, also tropical & subtropical montane



Example from Nicaragua. 1 *Albizia guachapele*; 2 *Bombacopsis quinata*; 3 *Cedrela odorata*; 4 *Gliciridia sepium*; 5 *Hymenaea courbaril*; 6 *Pinus maximinii*; 7 *Swietenia humilis*; 8 *Swietenia macrophylla*.

COLLECTION AND HANDLING RECORDS

Species information

Provisional seed lot no. _____ Suppliers seed lot no. _____
 Species name (botanical): _____ Species code: _____
 Provenance name: _____
 Common name: _____
 Seed source ref. no. _____ Country _____

Fruit collection

Collection date: _____
 Collection methods: From the tree, From ground after shaking, From ground after natural fall
 Other, indicate details _____
 Genetic representation: Number of parent trees collected from: _____
 Average spacing between parent trees: _____
 Phenotypic selection of seed trees: Yes No
 if yes, indicate criteria: Height, Straightness, Branching habit, Health, Others, pls. indicate: _____

Quantity of fruits collected _____
 Container used: _____ Number of containers filled: _____
 Total weight of containers including fruits: _____
 Weight of empty containers: _____
 Weight of fruits before processing: _____

Field handling and transport

Condition of fruits during collection: _____
 No. of days from collection to field handling: _____
 Field handling: None, After-ripening, Drying, Extraction, pls. indicate method: _____
 If extracted, please indicate volume or weight of extracted seed fraction: _____
 Moisture content (moisture meter) after field handling: _____ %
 Temporary field storage: Duration _____ days, Mode and condition of transport _____

Date _____

Signature of collection officer _____

Date arrived:	Quantity (kg) :	Fruits,	Seeds
Apparent condition at arrival: _____			
Moisture content at arrival (moisture meter):		%	
After ripening: Date:		Method/conditions:	
Seed extraction:	<input type="checkbox"/> Sun drying, temperature	Duration	days
	<input type="checkbox"/> Kiln drying, temperature	Duration	days
	<input type="checkbox"/> Depulping, pls. indicate method: _____		
	<input type="checkbox"/> Other method, pls. indicate _____		
Other processing: <input type="checkbox"/> Dewinging, pls. indicate method _____			
<input type="checkbox"/> Cleaning, pls. indicate method _____			
<input type="checkbox"/> Other, pls. indicate _____			
Pest and diseases: <input type="checkbox"/> Infected or diseased seeds noted in seed lot, pls. indicate type: _____			
<input type="checkbox"/> Seed treatment, pls. indicate type _____			

Seed storage information

Date of entry into storage:	Quantity (kg) :	Moisture content	%
Suspected storage physiology: <input type="checkbox"/> Unknown, <input type="checkbox"/> Orthodox, <input type="checkbox"/> Recalcitrant, <input type="checkbox"/> Intermediate			
Storage unit: <input type="checkbox"/> Seed, <input type="checkbox"/> Dewinged seed, <input type="checkbox"/> Entire fruit, <input type="checkbox"/> Dewinged fruit, <input type="checkbox"/> Depulped fruit,			
<input type="checkbox"/> Other, pls indicate: _____			
Storage conditions:			
Packing and containers: <input type="checkbox"/> Open plastic bag, <input type="checkbox"/> Sealed plastic bag, <input type="checkbox"/> Sealed glass, <input type="checkbox"/> Tins,			
<input type="checkbox"/> Other, pls. indicate: _____			
Storage atmosphere: <input type="checkbox"/> Normal atmosphere, <input type="checkbox"/> CO ₂ , <input type="checkbox"/> Other, pls. indicate: _____			
Light conditions: <input type="checkbox"/> Light, <input type="checkbox"/> Dark			
Humidity:		<input type="checkbox"/> Ambient, <input type="checkbox"/> Controlled, approx. percentage	%
Temperature:		<input type="checkbox"/> Ambient <input type="checkbox"/> Controlled, degrees:	°C

_____	_____
Date	Signature

SEED PROCUREMENT ACCOUNTING FORM

Species information

Provisional seed lot no. _____	Suppliers seed lot no. _____
Species name (botanical): _____	Species code: _____
Provenance name: _____	
Common name: _____	
Seed source ref. no. _____	Country _____

Collection costs

Work type	basic unit	number	number x units	price per unit	total price
Crop survey					
Transport to, within and from site					
Labour cost :					
Supervisors					
Prof. collectors					
Casual workers					
Field accomodation					

Processing cost

Work type	basic unit	number	number x units	price per unit	total price

Total procurement cost

Collection + processing costs	
Quantity of clean seed	kg
Procurement cost per kg	

SEED SAMPLING FORM

Species information

Seed lot no. _____	Seed source ref. no. _____	Species code _____
Species name (botanical): _____	Provenance name: _____	
Common name: _____		
Country _____		

Fruit collection

Date of sampling: _____

Purpose of sampling: Moisture content
 Purity
 Seed weight , 1000 grain
 Germination
 Vigour
 Other, indicate details _____

Type of samples: Composite of primary
 Submitted of composite
 Working sample

Number of primary samples _____ from _____ number of bags or containers, or
 Number of primary samples _____ from _____ kilograms of seed

Weight of sample _____ g

Notes on sampling method:

_____	_____
Date	Signature

SEED TEST FORM I: Weight, purity and moisture content sheet

Seed lot no. _____ Seed source name : _____ Seed source ref. no: _____ Analysis no: _____
 Species _____

Date of completed germination analysis:

1000-grain weight (8x100) seed		Purity						Date of completed germination analysis:	
Replicate	X	Replicate	Weight of total sample	Weight of impurities	Weight of pure seed	Percentage impurities	Percentage pure seed	Difference between A and B:	
1		A	g	g	g	%	%	Max difference according to ISTA:	
2		B	g	g	g	%	%	New analysis yes no	
3		Average							
4									
		Moisture content							
Replicate		Replicate	Weight of empty container	Weight of fresh sample	Weight of oven-dry sample	Difference = weight of water	Moisture content	Difference between A and B:	
5		A	g	g	g	g	g	Moisture content = _____ %	
6		B	g	g	g	g	g	Max difference according to ISTA _____	
7		C	g	g	g	g	g	New analysis yes no	
8		D	g	g	g	g	g		
Sum									
Average X =	(n=8)								
Weight of 1000 seed =	gram	Average							
Range (=Largest - smallest) =									
Estimated standard deviation:									
Range / 2.85* =									
Var. Coeff: $\frac{100 \times \text{std.}}{X}$									
Not to exceed 4.0									
New analysis yes no									
* Table value for n=8									
Remarks (extra analysis, analysis errors, mech. seed damage etc.)									

SEED TEST FORM II: Germination test sheet

Seed lot no: _____ Seed source name: _____ Seed source ref. no: _____ Analysis no: _____
 Species: _____

Summary		Purity (%)	1000 seed wgt (g)	No. of seed/kg	Moisture content (%)	Germ. Cap. (%)	Rate of germ. (%)				Date of completed germination analysis			
Replication	Normal germination after days			Total norm. Germ. (a)	Fresh not germ. (b)	Abnormal germ. (c)	Dead seed		Total (a-e)	Rotten (mouldy seed) (f)	Polyem bryony (g)	Insect damaged seed (h)	Remarks (unlined field)	
							Empty (d)	Full (e)					Calculated germ. % (i)	Living seed (ex. Emp. Seed) (j)
A														
B														
C														
D														
Total														
Av. %														

Germination table / Cabinet type _____
 Germination substrate: _____
 Germination temp. ° C _____
 Pretreatment: _____
 Method _____
 Time _____
 Temp. (°C) _____

Damage: None Small Average Large

Fungi Insects Other

Difference in germination % A-B-C-D (%) _____
 Max. Difference (ISTA) (%) _____
 New analysis yes no
 New analysis start _____
 New analysis number _____
 Germination test done by _____

SEED LOT SUMMARY DATA SHEET

Seed lot no. _____ Supplier _____
 Species name (botanical): _____ Provenance name: _____
 Common name : _____
 Country _____

Seed source information

Seed source location: _____ Region/state _____ Country _____
 Geographical coordinates: Latitude: _____ ° 'N/S, longitude: _____ ° 'E/W
 Altitude: _____ m.a.s.l
 Mean annual rainfall (mm): _____ Rainfall regime: Summer Uniform Winter Bimodal
 Soil type: _____ pH _____
 Stand type: Natural stands Plantation
 Seed source type: Unclassified, Seed zone, Identified stand, Selected stand,
 Seed production area, Provenance seed stand, Seed orchard
 Other information: _____

Collection data

Collection date: _____
 Genetic representation: Number of parent trees collected from: _____
 Average spacing between parent trees: _____
 Phenotypic selection of seed trees: Yes No
 Selection criteria: Height, Straightness, Branching habit, Health, Others, _____

Test results

Date of (latest) test _____	Germination percentage _____
Purity: _____ %	Viability:
Moisture content : _____ %	Measured by : <input type="checkbox"/> TTZ
1000 grain seed weight: _____	<input type="checkbox"/> Cutting
	<input type="checkbox"/> X-ray
	<input type="checkbox"/> Other: _____
	No. of viable seeds per gram: _____

Seed treatment

Seeds treated with: _____
 Date of treatment _____

Recommended seed handling before sowing

Pretreatment:	<input type="checkbox"/> Scarification, method and duration: _____
	<input type="checkbox"/> Stratification, method and duration: _____
	<input type="checkbox"/> Soaking in water, duration: _____
	<input type="checkbox"/> Leaching, duration _____
	<input type="checkbox"/> Manual extraction, method: _____
	<input type="checkbox"/> Other, _____
Inoculation:	<input type="checkbox"/> Mycorrhiza, species / type: _____
	<input type="checkbox"/> Rhizobium, species/ type: _____
	<input type="checkbox"/> Frankia, species/ type : _____

_____	_____
Date	Signature

SEED LABELS

SEEDLOT PROVISIONAL NO.: _____

SPECIES: _____

SEED SOURCE (REF.): _____

SACK NO.: _____ OF TOTAL _____ NOS. OF BAGS

QUANTITY IN SACK: _____ kg/g _____ litres

COLLECTED BY: _____

COLLECTED DATE: _____

Appendix A14.2B
Example of collection and handling form used by Kenya Forest Seed Centre

Kenya Forest Seed Centre

SEED COLLECTION REPORT

Species Collection date
 Collector Collection No.

LOCALITY

District Location
 Forest Station Block/Compartment
 Longitude Latitude Elevation
 Detailed location (sketchmap with directions and distances a MUST on back of page):
 Persons to contact (in case)

SITE/SOIL

Topography: flat/mountainous Slope: gentle/medium/steep
 Soil type Depth
 Soil colour Drainage
 Soil texture Stoniness

STAND

Land use: Forest/agriculture/pasture/open range/roadside/plantation/town
 Density: Open/group/dense
 Associated species
 Natural stand/plantation Age if natural: Young/middle aged/old
 If plantation: Year of planting Original sources of trees
 Form: Boles: Single/forked/multistem Length of clear bole.....m
 Crown: Narrow/average/wide Flat/pointed

COLLECTION

Type of collection: Seed orchard/selected seed stand/selected single trees/general/established seed stand
 If selected trees/seed orchard/established stand number of trees/clones
 collected from
 If general/seed stand: how many trees collected from
 Minimum distance between collection trees:
 Method of collection: ground/crown Others (specify)
 Weight of collected fruit (kg)..... Viability (cutting test)
 Remarks

SEED HANDLING

Temporary storage method before extractionDuration
 Extraction method Duration
 Drying method after extraction Duration
 Temporary storage method before delivery Duration
 Weight of fruits/extracted seedskg
 Weight of fruits/extracted seeds received to Muguga kg
 Weight of fruits/extracted seeds received at Muguga kg
 Date of delivery to Muguga
 Remarks

Signature: Officer in Charge

Appendix A14.2C

Example of a page in a computerized seed stock register. Danida forest Seed Centre.

ACC_NO	DATEREC	QUAN_ORIG	QUAN_BAL	LASTUPDATE	Stock_Check	Stock_diff
01012/81	04-10-83	7500	2716	14-02-95	2730	-14
01013/81	04-10-83	11000	5722	06-03-97	5735	-13
01014/82	06-07-83	750	388	14-02-95	403	-15
01015/82	06-07-83	700	323	14-02-95	336	-13
01016/82	06-07-83	600	393	14-02-95	393	-18
01017/82	06-07-83	850	585	14-02-95	504	81
01018/82	06-07-83	800	446	14-02-95	464	-18
01022/82	13-05-82	42455	18624	27-09-96	18764	-140
01027/82	08-07-82	6600	3992	14-02-95	3930	62
01028/82	08-07-82	6600	4143	14-02-95	4143	98
01034/82	10-06-84	1000	562	14-02-95	575	-13
01035/82	11-05-84	2217	547	14-02-95	505	42
01036/82	11-05-84	1633	564	14-02-95	439	125
01037/82	12-12-85	9064	2990	10-01-96	3010	-20
01038/82	12-12-85	4425	1130	10-01-96	1099	31
01039/82	12-12-85	8418	4720	06-03-97	4740	-20
01040/82	12-12-85	12447	5463	06-03-97	5355	108
01041/82	12-12-85	1859	605	10-01-96	615	-10
01042/82	11-08-83	3887	1422	02-11-99	1428	4
01043/82	11-08-83	2539	800	10-01-96	805	-5
01044/82	11-08-83	2668	825	10-01-96	835	-10
01045/82	10-02-83	7000	4054	06-03-97	4065	-11
01046/82	10-02-83	3000	2810	10-01-96	2830	-20
01047/82	10-02-83	2000	920	02-02-98	955	-35
01051/82	01-09-82	8000	2325	19-03-98	2374	-49
01052/82	10-02-83	49600	39000	20-12-95	39320	-320
01053/82	10-02-83	39700	35450	06-03-97	35845	-395
01054/82	10-02-83	10000	4811	30-01-97	4790	21
01057/82	14-10-82	30000	24177	17-01-97	24123	54
01058/82	14-10-82	53000	32388	02-02-98	32440	-52
01059/82	14-10-82	30597	24438	10-01-96	24617	-179
01060/82	14-10-82	30080	2716	17-01-97	2796	-80
01061/82	14-10-82	20000	1637	10-01-96	1645	-8
01062/82	14-04-83	2000	375	10-01-96	383	-8
01065/82	14-04-83	1000	335	10-01-96	343	-8
01066/82	14-09-83	7000	4050	10-01-96	4070	-20
01067/82	25-01-83	4000	561	10-01-96	560	1
01068/82	02-08-84	10000	7000	02-11-99	7050	-40
01069/82	03-06-84	5500	1791	10-01-96	1795	-4
01070/82	03-06-84	8500	4400	06-03-97	4425	-25
01071/82	03-06-84	7000	2288	10-01-96	2295	-7
01072/82	03-06-84	11000	7820	10-01-96	7845	-25
01073/82	03-06-84	11000	8785	10-01-96	8825	-40
01074/82	03-06-84	10800	9360	10-01-96	9395	-35
01075/82	04-06-83	11000	8605	10-01-96	8579	26
01076/82	03-06-84	11000	5400	10-01-96	5420	-20
01077/82	03-06-84	11000	6692	10-01-96	6730	-38
01078/82	03-06-84	11000	8206	06-03-97	8235	-29
01079/82	03-06-84	11000	7776	10-01-96	7855	-79
01080/82	03-06-84	7500	4162	06-03-97	4170	-8
01081/82	03-06-84	4350	2596	02-11-99	2590	16
01082/82	03-06-84	11000	4505	10-01-96	4520	-15

Appendix A14.2E
 Example of summary seed record form used by Australian Tree Seed Centre

◀ **SEED RECORD** ▶



Seedlot No. 16009	Species code ACM	BOTANICAL NAME Eucalyptus acmenoides	Store code	Cost code 5														
EXACT LOCALITY OF COLLECTION		PARENT TREE(S)	SEED															
Dbase S11 3 km N.N.W. of Minichilli QLD Barakula State Forest		No. In bulk of parents 1 tree 1 D.B.H. 54 (cm) Total Height 15 (m) Form Fair	Collector C. Gardiner, D. Crawford Collectors No. CG230 Collection date 3 November 1986 Project ALCOA Identified by J. Turner Condition Good Storage Tin Room Temp Storage date 30 July 1987 Quantity 376g															
Forest Type		Remarks	GERMINATION															
Associate Trees E. punctata ssp. longirostrata, E. melliodora, Angophora costata, Callitris glauca																		
Latitude (°S) 26°22'	Longitude (°E) 150°27'																	
Altitude (m) 330	Aspect	Slope Flat																
Geology and Soil Orange-brown loamy sand		Fumigation method: CO₂ date: Nov '86																
pH: 4.5-5.0		Computer database updated																
			<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="2">Method</th> <th colspan="2">Date</th> <th rowspan="2">Viability/10 g or (%)</th> </tr> <tr> <th>From</th> <th>To</th> </tr> </thead> <tbody> <tr> <td>TVP 30°C</td> <td>6.10.87</td> <td>26.10.87</td> <td>37.5</td> </tr> <tr> <td>TVP 25°C</td> <td>16.8.95</td> <td>14.9.95</td> <td>27.5</td> </tr> </tbody> </table>		Method	Date		Viability/10 g or (%)	From	To	TVP 30°C	6.10.87	26.10.87	37.5	TVP 25°C	16.8.95	14.9.95	27.5
Method	Date		Viability/10 g or (%)															
	From	To																
TVP 30°C	6.10.87	26.10.87	37.5															
TVP 25°C	16.8.95	14.9.95	27.5															

Appendix A14.3

Example of screen data entry forms used in a computerized seed documentation system in Indonesia (form Purnomo 1996, plus SIS database)

Seed Source Description Data Entry

Species	<input type="text" value="Acacia mangium"/>	Province	<input type="text" value="1"/>	
Producer	<input type="text"/>	Seed Source No	<input type="text" value="01-1190"/>	
Seed Zone No	<input type="text"/>	<input type="button" value="STAND"/>		
District	<input type="text"/>	<input type="button" value="SITE"/>		
Division	<input type="text"/>	<input type="button" value="MAP-INFO"/>		
Name of site	<input type="text"/>	Seed Source Class <input checked="" type="radio"/> Identified Stand <input type="radio"/> Selected Stand <input type="radio"/> Seed Production Area <input type="radio"/> Provenance Seed Stand <input type="radio"/> Seedling Seed Orchard <input type="radio"/> Clonal Seed Orchard		
Latitude	<input type="text"/>			
Longitude	<input type="text"/>	Demarcation of boundary <input type="text"/>		
Altitude (m.a.s.l)	<input type="text"/>	Detailed location <input type="text"/>		
Total area (ha)	<input type="text" value="100"/>			
Productive area (ha)	<input type="text" value="8000"/>			
Date of latest assessme	<input type="text" value="0"/>			
Date of registration	<input type="text" value="0"/>			

Seed Lot Record Entry form

Select Seed Source
Seed lot accession number

Flowering intensit <input type="radio"/> heavy <input checked="" type="radio"/> medium <input type="radio"/> light	Fruiting intensit <input type="radio"/> heavy <input checked="" type="radio"/> medium <input type="radio"/> light	Fruit distributio <input checked="" type="radio"/> uniform <input type="radio"/> scattered <input type="radio"/> border trees	Insect Damage <input type="text" value="0"/>
Total harvest, kg fruit	<input type="text"/>		
Total yield, kg seed	<input type="text" value="40"/>	Effective collection area (% of total are	<input type="text" value="1.3"/>
Number of trees collected fro	<input type="text" value="0"/>	Spacing of trees collected from	<input type="text" value="0"/>
<input type="button" value="Add Seed Lot"/>	<input type="button" value="Tests of Seed Lot"/>	<input type="button" value="Close"/>	

Appendix A14.3 (cont.)

Example of screen data entry forms used in a computerized seed documentation system in Indonesia (form Purnomo 1996, plus SIS database)

SEED QUALITY DATA			
Seed lot accession number: 07-newest-12			
SPECIES:			
Botanical name:	<i>Abies procera</i>	English name	noble fir
Common name:	don't know	Local name:	-
Pretreatment:			
Day of germination (days)	0	Insects (X-ray) (%)	0
Germination (%)	0	No. germs/gram	0
1000 seed weight (gr)	0	T.Z. pure red (%)	0
Purity (%)	0	Cutting. Fresh (%)	0
Moisture (%)	0	Viable seeds/kg	0
Remarks:			
Date of finished test:		Seed test no.: 0	
Test carried out by:			
Signed:			