



The Hagedorn temperature of AdS(5)/CFT4 at finite coupling via the Quantum Spectral Curve

Harmark, Troels; Wilhelm, Matthias

Published in:
Physics Letters B

DOI:
[10.1016/j.physletb.2018.09.033](https://doi.org/10.1016/j.physletb.2018.09.033)

Publication date:
2018

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Harmark, T., & Wilhelm, M. (2018). The Hagedorn temperature of AdS(5)/CFT4 at finite coupling via the Quantum Spectral Curve. *Physics Letters B*, 786, 53-58. <https://doi.org/10.1016/j.physletb.2018.09.033>



The Hagedorn temperature of $\text{AdS}_5/\text{CFT}_4$ at finite coupling via the Quantum Spectral Curve

Troels Harmark*, Matthias Wilhelm

Niels Bohr Institute, Copenhagen University, Blegdamsvej 17, 2100 Copenhagen Ø, Denmark

ARTICLE INFO

Article history:

Received 4 July 2018

Accepted 16 September 2018

Available online 18 September 2018

Editor: N. Lambert

ABSTRACT

Building on the recently established connection between the Hagedorn temperature and integrability [1], we show how the Quantum Spectral Curve formalism can be used to calculate the Hagedorn temperature of $\text{AdS}_5/\text{CFT}_4$ for any value of the 't Hooft coupling. We solve this finite system of finite-difference equations perturbatively at weak coupling and numerically at finite coupling. We confirm previous results at weak coupling and obtain the previously unknown three-loop Hagedorn temperature. Our finite-coupling results interpolate between weak and strong coupling and allow us to extract the first perturbative order at strong coupling. Our results indicate that the Hagedorn temperature for large 't Hooft coupling approaches that of type IIB string theory in ten-dimensional Minkowski space.

© 2018 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The $\text{AdS}_5/\text{CFT}_4$ correspondence [2] provides an exact duality between two seemingly very different theories. On the one side, one has a four-dimensional gauge theory in the form of $\mathcal{N} = 4$ super-Yang–Mills (SYM) theory on $\mathbb{R} \times S^3$ with 't Hooft coupling λ . On the other side, one has type IIB string theory on the ten-dimensional target-space $\text{AdS}_5 \times S^5$. This makes the $\text{AdS}_5/\text{CFT}_4$ correspondence an important theoretical laboratory for understanding various interesting problems in physics. One such problem is the nature of the Hagedorn temperature in string theory. Tree-level string theory has an exponentially growing density of states at large energies, which leads to a singularity in the thermodynamic partition function defining the Hagedorn temperature. Since the $\text{AdS}_5/\text{CFT}_4$ correspondence provides a non-perturbative definition of string theory, it should enable one to study the Hagedorn temperature and all its related phenomena.

In the $\text{AdS}_5/\text{CFT}_4$ correspondence, the Hagedorn singularity is connected to the Hawking–Page transition that occurs at a lower temperature where the black hole phase becomes thermodynamically favorable over a gas of closed strings. On the gauge-theory side, this corresponds to the confinement–deconfinement transition, where the confined phase occurs due to the confinement of

the color degrees of freedom on a three-sphere [3–6]. However, in the limit of zero string coupling, or the strict limit of infinite colors on the gauge-theory side, this transition requires arbitrarily high energy to realize, and one is left with the Hagedorn temperature as a maximal possible temperature on both sides of the correspondence. In this somewhat simpler setting, a starting point for further exploration of finite-temperature physics is to establish a quantitative interpolation of the Hagedorn temperature between the gauge-theory and string-theory sides.

However, it is only for certain precious cases that one has exact methods available to make a quantitative interpolation from weak to strong 't Hooft coupling. One such method is integrability, see Refs. [7,8] for reviews. Recently, we proposed a framework for calculating the Hagedorn temperature of $\text{AdS}_5/\text{CFT}_4$ using integrability [1]. In this Letter, we take this a step further by exploiting this connection to compute the Hagedorn temperature at finite 't Hooft coupling. This enables us to interpolate all the way from zero 't Hooft coupling to large 't Hooft coupling where we find the Hagedorn temperature of type IIB string theory in flat space, in both cases matching a previous computation of Sundborg [5,9].

The proposal [1] for calculating the Hagedorn temperature of $\text{AdS}_5/\text{CFT}_4$ via integrability is as follows. Define $F(T)$ to be the free energy per unit classical scaling dimension in the limit of large classical scaling dimension of the spin chain associated with planar $\mathcal{N} = 4$ SYM theory. Then the Hagedorn temperature T_H in units of the S^3 radius is determined by

$$F(T_H) = -1, \quad (1)$$

* Corresponding author.

E-mail addresses: harmark@nbi.ku.dk (T. Harmark), matthias.wilhelm@nbi.ku.dk (M. Wilhelm).

for zero chemical potentials. This can be seen from the fact that Eq. (1) determines the temperature beyond which the planar partition function of $\mathcal{N} = 4$ SYM theory is singular. The free energy is computed from so-called Thermodynamic Bethe Ansatz (TBA) equations [1], which are an infinite system of integral equations. In Ref. [1], we have solved these equations perturbatively at weak coupling, reproducing the known tree-level result [5] and one-loop correction [10] as well as finding the previously unknown two-loop correction. In principle, one can employ the TBA equations to compute higher order corrections to T_H and to find T_H numerically at finite coupling as well. In practice, however, the nature of these equations massively complicates perturbative calculations and severely limits the numeric accuracy one can achieve at finite coupling.

The TBA equations for the spectral problem of $\mathcal{N} = 4$ SYM theory were recast into the form of the Quantum Spectral Curve (QSC) [11–13], see Refs. [14,15] for reviews. It consists of a finite system of finite-difference equations, which allows for a very efficient evaluation both perturbatively at weak coupling [16–18] and numerically at finite coupling [19,20]. The QSC was since used for the pomeron [21,22], cusped Wilson lines and the quark–antiquark potential [23–25] as well as integrable deformations of $\mathcal{N} = 4$ SYM theory [26–28].

In this Letter, we recast our TBA equations for the Hagedorn temperature T_H of AdS₅/CFT₄ into the form of the QSC. Moreover, we solve these equations perturbatively at weak coupling and numerically at finite coupling.

2. QSC equations for the Hagedorn temperature

QSC equations The TBA equations are an infinite system of integral equations given in terms of Y-functions. They can be recast into the form of the so-called Y-system and T-system, which are infinite systems of finite-difference equations, and subsequently into the form of the so-called Q-system, which is a finite system of finite-difference equations also known as the Quantum Spectral Curve (QSC). Since we are setting all chemical potentials to zero, we are in a situation with so-called left-right symmetry, which is a symmetry between the two $\mathfrak{su}(2|2)$ subalgebras of the superconformal symmetry algebra $\mathfrak{psu}(2,2|4)$. The QSC is then formulated in terms of the functions $\mathbf{P}_a(u)$, $\mathbf{Q}_i(u)$ and $Q_{a|i}(u)$, where $a, i = 1, 2, 3, 4$ and u is the spectral parameter. They satisfy the finite-difference equations

$$Q_{a|i}^+ - Q_{a|i}^- = \mathbf{P}_a \mathbf{Q}_i, \quad (2)$$

$$\mathbf{P}_a = -\mathbf{Q}^i Q_{a|i}^+, \quad (3)$$

where $f^\pm(u) = f(u \pm \frac{i}{2})$. The functions $Q_{a|i}$ are orthonormal in the sense that

$$Q_{a|i} Q^{b|i} = \delta_a^b, \quad Q_{a|i} Q^{a|j} = \delta_i^j. \quad (4)$$

Here the functions with upper indices are defined as

$$\mathbf{P}^a = \chi^{ab} \mathbf{P}_b, \quad \mathbf{Q}^i = \chi^{ij} \mathbf{Q}_j, \quad Q^{a|i} = \chi^{ab} \chi^{ij} Q_{b|j}, \quad (5)$$

where the non-zero entries of χ are $\chi^{14} = \chi^{32} = -1$, $\chi^{23} = \chi^{41} = 1$. The Eqs. (2)–(4) reflect the $\mathfrak{psu}(2,2|4)$ symmetry of $\mathcal{N} = 4$ SYM theory, where $\mathbf{P}_a(u)$ ($\mathbf{Q}_i(u)$) is associated to the conformal symmetry $\mathfrak{su}(2,2)$ (R-symmetry $\mathfrak{su}(4)$). They are universal in the sense that they do not depend on the specific physical observable that one is computing but are common to all cases so far investigated. In order to specify a particular physical observable, these universal equations have to be supplemented by the asymptotic behavior of the functions at large spectral parameter u , by the location of the branch cuts and by the discontinuities across these branch cuts.

otic behavior of the functions at large spectral parameter u , by the location of the branch cuts and by the discontinuities across these branch cuts.

Asymptotic behavior We can infer the asymptotic behavior of $\mathbf{P}_a(u)$ and $\mathbf{Q}_i(u)$ at large spectral parameter u from the asymptotic behavior of the Y-functions found in Ref. [1]. At large u , the Y-functions asymptote to constants determined from a one-parameter family of constant T-systems with parameter z , see Eqs. (26)–(27) in Ref. [1]. Using the TBA equations, we show in Ref. [29] that $F(T) = -4T \operatorname{arctanh} z$. Since the Hagedorn temperature T_H satisfies Eq. (1), this fixes $z = \tanh \frac{1}{4T_H}$. The asymptotic Y-functions are reproduced via¹

$$\begin{aligned} \mathbf{P}_1(u) &= A_1 \left(-e^{-\frac{1}{2T_H}} \right)^{-iu} \left(1 + \mathcal{O}(u^{-1}) \right), \\ \mathbf{P}_2(u) &= A_2 \left(-e^{-\frac{1}{2T_H}} \right)^{-iu} \left(u + \mathcal{O}(u^0) \right), \\ \mathbf{P}_3(u) &= A_3 \left(-e^{-\frac{1}{2T_H}} \right)^{+iu} \left(1 + \mathcal{O}(u^{-1}) \right), \\ \mathbf{P}_4(u) &= A_4 \left(-e^{-\frac{1}{2T_H}} \right)^{+iu} \left(u + \mathcal{O}(u^0) \right) \end{aligned} \quad (6)$$

and

$$\begin{aligned} \mathbf{Q}_1(u) &= B_1 \left(1 + \mathcal{O}(u^{-1}) \right), \quad \mathbf{Q}_2(u) = B_2 \left(u + \mathcal{O}(u^0) \right), \\ \mathbf{Q}_3(u) &= B_3 \left(u^2 + \mathcal{O}(u^1) \right), \quad \mathbf{Q}_4(u) = B_4 \left(u^3 + \mathcal{O}(u^2) \right), \end{aligned} \quad (7)$$

with $A_1 A_4 = A_2 A_3 = \frac{i}{\tanh^2 \frac{1}{4T_H}}$ and $3B_1 B_4 = B_2 B_3 = -8i \cosh^4 \frac{1}{4T_H}$. The asymptotic behavior of $Q_{a|i}(u)$ then follows from Eq. (2).

Branch cut structure and ansatz We consider the so-called direct theory rather than the mirror theory; hence, we use the Zhukowski variable

$$x(u) = \frac{u}{2g} \left(1 + \sqrt{1 - \frac{4g^2}{u^2}} \right), \quad (8)$$

which has a ‘short’ branch cut at the interval $(-2g, 2g)$, where $g^2 = \frac{\lambda}{16\pi^2}$ is the effective planar loop coupling. We work in a Riemann sheet in which the four functions $\mathbf{Q}_i(u)$ have one short cut at the interval $(-2g, 2g)$, while $\mathbf{P}_a(u)$ and its analytic continuation $\tilde{\mathbf{P}}_a(u)$ have an infinite set of short cuts at $(-2g, 2g) - in$ and $(-2g, 2g) + in$ with $n \in \mathbb{N}_{\geq 0}$, respectively. Since $\mathbf{Q}_i(u)$ has only a single short cut, we can make the ansatz

$$\mathbf{Q}_i(u) = B_i (gx(u))^{i-1} \left(1 + \sum_{n=1}^{\infty} \frac{c_{i,n}(g)}{(gx(u))^{2n}} \right). \quad (9)$$

For convenience, we choose the gauge $c_{3,1} = 0$ and $B_1 = B_2 = 1$.

The previous applications of the QSC formalism have all been in the mirror theory rather than in the direct theory that we consider here. In the mirror theory, it is the four functions $\mathbf{P}_a(u)$ for which one can choose a Riemann sheet where they have only one short cut at the real axis. This means one makes an ansatz for the functions $\mathbf{P}_a(u)$ instead of the functions $\mathbf{Q}_i(u)$ as we do in our case.

¹ Interestingly, the asymptotics (6)–(7) are similar to the ones of the spectral problem of twisted $\mathcal{N} = 4$ SYM theory [26]. Concretely, excluding the prefactors they are formally the same if we set the twists $1/x_1 = 1/x_2 = x_3 = x_4 = -e^{-\frac{1}{2T_H}}$, $y_1 = y_2 = y_3 = y_4 = 1$ and insert the Cartan charges $\Delta = S_1 = S_2 = J_1 = J_2 = J_3 = 0$. This is related to how the constant T-system [1] is obtained from the general $\mathfrak{psu}(2,2|4)$ character solution [43].

Gluing conditions To close the system of QSC equations, one needs to impose so-called gluing conditions [14]. They relate the analytic continuation $\tilde{\mathbf{P}}_a(u)$ through the short cut at the real axis to a linear combination of the complex conjugates of the $\mathbf{P}_b(u)$ functions. In our case, using the gauge choice $A_1 = iA_2 = -A_3 = -iA_4 = (\tanh \frac{1}{4T_H})^{-1}$ for the asymptotics (6), the gluing conditions are

$$\tilde{\mathbf{P}}_a(u) = (-1)^{1+a} \overline{\mathbf{P}_a(u)}. \quad (10)$$

Together with the QSC equations (2)–(4), the ansatz (9) for the functions $\mathbf{Q}_i(u)$ and the large- u asymptotics (6)–(7), the gluing conditions (10) determine the Hagedorn temperature T_H for any given value of g – which is one of the main results of this Letter. Another main result is that we will explicitly solve these equations perturbatively at weak coupling and numerically at finite coupling, as explained below.

3. Perturbative solution

To solve the QSC equations perturbatively at weak coupling, we start with the tree-level solution for $g = 0$. It can be obtained from Eqs. (6)–(7) by setting T_H to the tree-level Hagedorn temperature $T_H^{(0)} = 1/(2 \log(2 + \sqrt{3}))$ [5], which determines the leading coefficients. The only non-vanishing subleading coefficient at tree level is $c_{4,1}(0) = -1$. Using the Eqs. (2)–(4), one finds the corresponding functions $Q_{aj_i}(u)$ at tree level, which we denote below as $Q_{aj_i}^{(0)}(u)$.

Knowing the tree-level solution, we can now solve the QSC equations (2)–(4) perturbatively following a slightly modified version of the approach in Ref. [22]. Write the solution of Eq. (2) as

$$Q_{aj_i} = Q_{aj_i}^{(0)} + (b_a^c)^+ Q_{c_{ji}}^{(0)}. \quad (11)$$

Then $b_a^c(u)$ satisfies the first order finite-difference equation

$$(b_a^c)^{++} - b_a^c = dS_{aj_i}(Q^{(0)c_{ji}})^- + (b_a^b)^{++} dS_{bj_i}(Q^{(0)c_{ji}})^-, \quad (12)$$

where dS_{aj_i} is defined as

$$dS_{aj_i} \equiv Q_{aj_i}^{(0)+} - Q_{aj_i}^{(0)-} + \mathbf{Q}_i \mathbf{Q}_j Q_{aj_i}^{(0)+}. \quad (13)$$

Here, we use the ansatz (9) for $\mathbf{Q}_i(u)$, in which the sum truncates for any given loop order ℓ . Assume one has already determined the coefficients of $\mathbf{Q}_i(u)$ in Eq. (9) at $(\ell - 1)$ -loop order. Solving Eq. (12), we find $b_a^c(u)$ and hence $Q_{aj_i}(u)$ at ℓ -loop order in terms of the as yet undetermined parameters in the ansatz (9) as well as additional undetermined constant parameters from the homogeneous solution to Eq. (12). Because of the phases in Eq. (6), we encounter finite-difference equations of the type

$$z_i f^{++}(u) - f(u) = h(u), \quad z_i \in \{1, (2 + \sqrt{3})^{\pm 2}\}. \quad (14)$$

The solution can be written in terms of the generalized η functions [23,26]

$$\eta_{s_1, \dots, s_k}^{z_1, \dots, z_k}(u) \equiv \sum_{n_1 > n_2 > \dots > n_k \geq 0} \frac{z_1^{n_1} \dots z_k^{n_k}}{(u + in_1)^{s_1} \dots (u + in_k)^{s_k}}. \quad (15)$$

Since in some cases $z_i = (2 + \sqrt{3})^2$, we have to use analytic continuation as a regularization. Note that this is a worse divergence than in the twisted QSC [26] since in that case z_i is on the unit circle. Evaluated at $u = i$, the generalized η functions are proportional to multiple polylogarithms

$$\text{Li}_{s_1, \dots, s_k}(z_1, \dots, z_k) \equiv \sum_{n_1 > n_2 > \dots > n_k > 0} \frac{z_1^{n_1} \dots z_k^{n_k}}{n_1^{s_1} \dots n_k^{s_k}}, \quad (16)$$

in terms of which the result for the Hagedorn temperature is naturally expressed. Note that the multiple polylogarithms have branch cuts on the real axis; for instance, classical polylogarithms have branch cuts between 1 and ∞ . The ambiguity in evaluating these polylogarithms on the branch cut can be resolved by an $i\epsilon$ prescription.

The next step is to impose conditions to determine the un-fixed parameters. To begin with, we impose Eq. (4) which fixes half of the coefficients from the homogeneous solution in $Q_{aj_i}(u)$. By Eq. (3), we now find $\tilde{\mathbf{P}}_a(u)$. The gluing conditions (10) enter by imposing that $\mathbf{P}_a(u) + \tilde{\mathbf{P}}_a(u)$ and $(\mathbf{P}_a(u) - \tilde{\mathbf{P}}_a(u))/\sqrt{u^2 - 4g^2}$ are regular at $u = 0$. Finally, we have to impose the asymptotic behavior (6), which we implement by requiring that

$$\frac{\mathbf{P}_2(u)}{\mathbf{P}_1(u)} = -iu + \mathcal{O}(u^0). \quad (17)$$

Up to a gauge choice, this fixes all parameters of the ansatz (9) at ℓ -loop order, including the Hagedorn temperature T_H .²

We find, up to three-loop order,

$$\begin{aligned} T_H(g) &= \frac{1}{2 \log(2 + \sqrt{3})} + g^2 \frac{1}{\log(2 + \sqrt{3})} \\ &+ g^4 \left(48 - \frac{86}{\sqrt{3}} - \frac{48 \text{Li}_1\left(\frac{1}{(2+\sqrt{3})^2}\right)}{\log(2 + \sqrt{3})} \right) \\ &+ g^6 \left(624 \text{Li}_2\left(\frac{1}{(2+\sqrt{3})^2}\right) \right. \\ &\quad \left. + \frac{432 \text{Li}_1\left(\frac{1}{(2+\sqrt{3})^2}\right)^2}{\log(2 + \sqrt{3})} + \frac{312 \text{Li}_3\left(\frac{1}{(2+\sqrt{3})^2}\right)}{\log(2 + \sqrt{3})} \right) \\ &\quad \left. + \left(384\sqrt{3} - 864 + 416 \log(2 + \sqrt{3}) \right) \text{Li}_1\left(\frac{1}{(2+\sqrt{3})^2}\right) \right. \\ &\quad \left. - \frac{20}{\sqrt{3}} + \left(\frac{1900}{3} - 384\sqrt{3} \right) \log(2 + \sqrt{3}) \right) + \mathcal{O}(g^8) \\ &\approx (0.3796628588\dots) + (0.7593257175\dots)g^2 \\ &\quad + (-4.367638556\dots)g^4 + (37.22529358\dots)g^6 \\ &\quad + \mathcal{O}(g^8). \end{aligned} \quad (18)$$

This agrees with the previously known results at tree level [5], one-loop order [10] and two-loop order [1]. In an upcoming publication [29], we will also present the result at four-loop order and beyond.

4. Numerical solution

The QSC can also be solved numerically at finite values of g . Concretely, it can be reduced to a minimization problem that can be solved iteratively. We use a modified version of the approach in Ref. [19], see also Refs. [14,30].

Each iteration starts at some values for the Hagedorn temperature T_H and the coefficients $c_{1,n}$, $c_{2,n}$ for $n = 1, \dots, K$, $c_{3,n}$ for $n = 2, \dots, K$ and $c_{4,n}$ for $n = 3, \dots, K$ in the ansatz (9), which is truncated at some finite K . We make an ansatz for $Q_{aj_i}(u)$ at large u :

$$Q_{aj_i}(u) = \left(-e^{-\frac{1}{2T_H}} \right)^{-s_a i u} u^{p_{aj_i}} \sum_{n=0}^N \frac{B_{aj_i, n}}{u^n}, \quad (19)$$

² In contrast, the ℓ -loop anomalous dimension in the spectral problem is only fixed at the $(\ell + 1)$ th order.




Fig. 1. Numeric results and weak coupling approximation at various loop orders for the Hagedorn temperature as a function of g^2 .

which is truncated at some finite order depending on N . Here, we have used $s_a = 1$ and $p_{a|i} = a + i - 2$ for $a = 1, 2$ as well as $s_a = -1$ and $p_{a|i} = a + i - 4$ for $a = 3, 4$. We solve for the remaining coefficients $B_{a|i,n}$ by imposing Eqs. (4) and (2) where $\mathbf{P}_a(u)$ is eliminated using Eq. (3). In particular, this also fixes $c_{4,1}$ and $c_{4,2}$.

Starting at some finite but large imaginary value of u , we can shift $Q_{a|i}$ towards the real axis in steps of i using

$$Q_{a|i}^- = Q_{a|i}^+ + \mathbf{Q}_i \mathbf{Q}_i^j Q_{a|i}^+, \quad (20)$$

which follows from Eqs. (2)–(3). We can now reconstruct \mathbf{P}_a and its analytic continuation

$$\tilde{\mathbf{P}}_a = -\tilde{\mathbf{Q}}_i^j Q_{a|i}^+ \quad (21)$$

on the real axis, where $\tilde{\mathbf{Q}}_i$ is obtained from the ansatz (9) via $\tilde{x} = 1/x$. Note that this is however only possible for $a = 1, 2$, as in this case $Q_{a|i}^+$ is exponentially small for large imaginary u , while it is exponentially large for $a = 3, 4$.

Now we can define a function F that vanishes for an exact solution of the gluing conditions (10):

$$F(T_H, \{c_{i,n}\}) = \sum_{a=1}^2 \sum_{j=1}^P \left| \frac{\overline{\mathbf{P}_a(p_i)}}{\tilde{\mathbf{P}}_a(p_i)} + (-1)^a \right|^2, \quad (22)$$

where p_i are P points in the interval $(-2g, 2g)$. We can find an approximate solution for T_H and the coefficients $c_{i,n}$ by minimizing F iteratively, using for instance Newton's method or the Levenberg–Marquardt algorithm.

We have plotted our numeric results for the Hagedorn temperature T_H as a function of g^2 for $0 \leq g^2 \leq 0.1$ in Fig. 1. In addition, Fig. 1 contains the perturbative approximation to the non-perturbative results up to three-loop order. As expected, the perturbative series converges towards the exact results for sufficiently small values of g^2 .

Fig. 2 shows our numeric data for T_H as a function of \sqrt{g} for $0 \leq \sqrt{g} \leq 1.8$. In particular, we see that T_H tends towards a linear function in \sqrt{g} at strong coupling. Using a sixth-order fit in $1/\sqrt{g}$, we find the following approximate result for the leading coefficient:

$$T_H(g) = (0.399\dots)\sqrt{g} + \mathcal{O}(g^0), \quad (23)$$




Fig. 2. Numeric results and leading strong coupling approximation for the Hagedorn temperature as a function of \sqrt{g} .

where the uncertainty is in the last digit.³ At the given accuracy, this agrees with the expectation that the Hagedorn temperature measured in units of the S^3 radius R approaches the behavior

$$T_H(g) \simeq \sqrt{\frac{g}{2\pi}} \approx (0.3989422804\dots)\sqrt{g} \quad (24)$$

for large 't Hooft coupling. This corresponds to the Hagedorn temperature of tree-level type IIB string theory on ten-dimensional Minkowski space [9]. One can see this explicitly by reinstating $1/R$ using the $\text{AdS}_5/\text{CFT}_4$ dictionary. In terms of this, the S^3 radius R corresponds on the string-theory side to the radius $R = \lambda^{1/4} l_s$ of AdS_5 and S^5 , where l_s is the string length. Hence, Eq. (24) becomes

$$T_H(g) \simeq \frac{1}{\sqrt{8\pi} l_s}. \quad (25)$$

The reason that one expects this to be the Hagedorn temperature for large 't Hooft coupling is as follows. Consider a string with energy $l_s E$ in string units. If this energy is sufficiently high, also compared to the angular momenta on $\text{AdS}_5 \times S^5$, a particle mode of the string is probing distances shorter than the radius R of AdS_5 and S^5 . Moreover, if $\sqrt{l_s E} \ll R/l_s$, the extension of the string is much smaller than the radius R . Thus, there is an intermediate regime in which the spectrum of a string behaves as in flat space. As $\lambda \rightarrow \infty$, this means that the Hagedorn temperature approaches that of flat space.

5. Conclusion and outlook

In this Letter, we have derived integrability-based QSC equations that determine the Hagedorn temperature of planar $\mathcal{N} = 4$ SYM theory – and equally of type IIB string theory on $\text{AdS}_5 \times S^5$ – at any value of the 't Hooft coupling. We have solved these equations perturbatively at weak coupling, reproducing known results up to two-loop order and obtaining the previously unknown three-loop result (18). The same algorithm can also be used at higher orders, as we will demonstrate in an upcoming publication [29]. Moreover, we have solved the QSC numerically at finite coupling,

³ In a certain exactly solvable pp-wave limit of string theory, the leading contribution to the Hagedorn temperature is of order $\lambda^{1/3}$ with corrections in $1/\lambda^{1/3}$ [35]. In the present case, the leading contribution is of order $\lambda^{1/4}$ and we thus expect the corrections to be in $1/\lambda^{1/4}$.

allowing for an interpolation between weak and strong coupling. From our numeric results, we have read off the first coefficient in the strong coupling expansion. It would be interesting to increase the numeric precision even further using a C⁺⁺ implementation following Ref. [20] to obtain further coefficients in the strong coupling expansion.

We have found evidence that the Hagedorn temperature, which marks the temperature beyond which the planar partition function is singular, asymptotes to the Hagedorn temperature of type IIB string theory in ten-dimensional Minkowski space for large g . This is in line with the expectation that for large g the spectrum should approach that of type IIB string theory on flat space, as explained above. To test further that the spectrum approaches flat space, one could possibly use the techniques of this Letter to study the critical behavior of the partition function close to the Hagedorn singularity for large g . If the critical behavior matches the one of flat-space string theory, it would confirm that there is a regime of strongly coupled $\mathcal{N} = 4$ SYM theory in which the spectrum is that of tree-level type IIB string theory in ten-dimensional Minkowski space. Thus, this Letter opens up an interesting new regime in which one can explore the AdS₅/CFT₄ correspondence.

Note finally that while we have restricted ourselves to vanishing chemical potentials, the case of non-vanishing chemical potentials can be treated in a similar way [29]. This could provide a connection to the cases of the Hagedorn temperature in the pp-wave or spin-matrix-theory limits [31–40]. Moreover, it would be interesting to consider the Hagedorn temperature for integrable deformations of $\mathcal{N} = 4$ SYM theory (see [41] for one-loop results) and for the three-dimensional $\mathcal{N} = 6$ superconformal Chern–Simons theory, for which a QSC formulation for the spectral problem exists as well [42]. In particular, it would be intriguing to study what happens at strong coupling in these cases.

Acknowledgements

We thank Simon Caron-Huot, Marius de Leeuw, Claude Duhr, Nikolay Gromov, Sebastien Leurent, Fedor Levkovich-Maslyuk, Christian Marboe, Andrew McLeod, Stijn van Tongeren, Matt von Hippel and Konstantin Zarembo for very useful discussions. We thank Nikolay Gromov for sharing his Mathematica implementation of the algorithm in Ref. [19]. M.W. thanks the Institute for Advanced Study in Princeton for kind hospitality. T.H. acknowledges support from FNU grant number DFF-6108-00340. M.W. was supported in part by FNU through grant number DFF-4002-00037, the ERC starting grant number 757978, the Danish National Research Foundation (grant number DNRF91) and the Villum Fonden.

References

- [1] T. Harmark, M. Wilhelm, The Hagedorn temperature of AdS₅/CFT₄ via integrability, *Phys. Rev. Lett.* 120 (2018) 071605, arXiv:1706.03074 [hep-th].
- [2] J.M. Maldacena, The large N limit of superconformal field theories and supergravity, *Int. J. Theor. Phys.* 38 (1999) 1113–1133, arXiv:hep-th/9711200 [hep-th], *Adv. Theor. Math. Phys.* 2 (1998) 231.
- [3] J.J. Atick, E. Witten, The Hagedorn transition and the number of degrees of freedom of string theory, *Nucl. Phys. B* 310 (1988) 291–334.
- [4] E. Witten, Anti-de Sitter space, thermal phase transition, and confinement in gauge theories, *Adv. Theor. Math. Phys.* 2 (1998) 505–532, arXiv:hep-th/9803131 [hep-th].
- [5] B. Sundborg, The Hagedorn transition, deconfinement and $\mathcal{N} = 4$ SYM theory, *Nucl. Phys. B* 573 (2000) 349–363, arXiv:hep-th/9908001 [hep-th].
- [6] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas, M. Van Raamsdonk, The Hagedorn – deconfinement phase transition in weakly coupled large N gauge theories, *Adv. Theor. Math. Phys.* 8 (2004) 603–696, arXiv:hep-th/0310285 [hep-th].
- [7] N. Beisert, et al., Review of AdS/CFT integrability: an overview, *Lett. Math. Phys.* 99 (2012) 3–32, arXiv:1012.3982 [hep-th].
- [8] D. Bombardelli, A. Cagnazzo, R. Frassek, F. Levkovich-Maslyuk, F. Loebbert, S. Negro, I.M. Szécsényi, A. Sfondrini, S.J. van Tongeren, A. Torrielli, An integrability primer for the gauge-gravity correspondence: an introduction, *J. Phys. A* 49 (2016) 320301, arXiv:1606.02945 [hep-th].
- [9] B. Sundborg, Thermodynamics of superstrings at high-energy densities, *Nucl. Phys. B* 254 (1985) 583–592.
- [10] M. Spradlin, A. Volovich, A pendant for Polya: the one-loop partition function of $\mathcal{N} = 4$ SYM on $\mathbb{R} \times S^3$, *Nucl. Phys. B* 711 (2005) 199–230, arXiv:hep-th/0408178 [hep-th].
- [11] N. Gromov, V. Kazakov, S. Leurent, D. Volin, Quantum Spectral Curve for planar $\mathcal{N} = 4$ super-Yang–Mills theory, *Phys. Rev. Lett.* 112 (2014) 011602, arXiv:1305.1939 [hep-th].
- [12] N. Gromov, F. Levkovich-Maslyuk, G. Sizov, S. Valatka, Quantum spectral curve at work: from small spin to strong coupling in $\mathcal{N} = 4$ SYM, *J. High Energy Phys.* 07 (2014) 156, arXiv:1402.0871 [hep-th].
- [13] N. Gromov, V. Kazakov, S. Leurent, D. Volin, Quantum spectral curve for arbitrary state/operator in AdS₅/CFT₄, *J. High Energy Phys.* 09 (2015) 187, arXiv:1405.4857 [hep-th].
- [14] N. Gromov, Introduction to the spectrum of $\mathcal{N} = 4$ SYM and the Quantum Spectral Curve, arXiv:1708.03648 [hep-th], 2017.
- [15] V. Kazakov, Quantum Spectral Curve of γ -twisted $\mathcal{N} = 4$ SYM theory and fishnet CFT, arXiv:1802.02160 [hep-th], 2018.
- [16] C. Marboe, D. Volin, Quantum spectral curve as a tool for a perturbative quantum field theory, *Nucl. Phys. B* 899 (2015) 810–847, arXiv:1411.4758 [hep-th].
- [17] C. Marboe, V. Velizhanin, D. Volin, Six-loop anomalous dimension of twist-two operators in planar $\mathcal{N} = 4$ SYM theory, *J. High Energy Phys.* 07 (2015) 084, arXiv:1412.4762 [hep-th].
- [18] C. Marboe, D. Volin, The full spectrum of AdS₅/CFT₄ I: representation theory and one-loop Q-system, arXiv:1701.03704 [hep-th], 2017.
- [19] N. Gromov, F. Levkovich-Maslyuk, G. Sizov, Quantum Spectral Curve and the numerical solution of the spectral problem in AdS₅/CFT₄, *J. High Energy Phys.* 06 (2016) 036, arXiv:1504.06640 [hep-th].
- [20] Á. Hegedűs, J. Konczer, Strong coupling results in the AdS₅/CFT₄ correspondence from the numerical solution of the quantum spectral curve, *J. High Energy Phys.* 08 (2016) 061, arXiv:1604.02346 [hep-th].
- [21] M. Alfimov, N. Gromov, V. Kazakov, QCD pomeron from AdS/CFT Quantum Spectral Curve, *J. High Energy Phys.* 07 (2015) 164, arXiv:1408.2530 [hep-th].
- [22] N. Gromov, F. Levkovich-Maslyuk, G. Sizov, Pomeron eigenvalue at three loops in $\mathcal{N} = 4$ supersymmetric Yang–Mills theory, *Phys. Rev. Lett.* 115 (2015) 251601, arXiv:1507.04010 [hep-th].
- [23] N. Gromov, F. Levkovich-Maslyuk, Quantum Spectral Curve for a cusped Wilson line in $\mathcal{N} = 4$ SYM, *J. High Energy Phys.* 04 (2016) 134, arXiv:1510.02098 [hep-th].
- [24] N. Gromov, F. Levkovich-Maslyuk, Quark–anti-quark potential in $\mathcal{N} = 4$ SYM, *J. High Energy Phys.* 12 (2016) 122, arXiv:1601.05679 [hep-th].
- [25] A. Cavaglià, N. Gromov, F. Levkovich-Maslyuk, Quantum Spectral Curve and structure constants in $\mathcal{N} = 4$ SYM: cusps in the ladder limit, arXiv:1802.04237 [hep-th], 2018.
- [26] V. Kazakov, S. Leurent, D. Volin, T-system on T-hook: grassmannian solution and twisted Quantum Spectral Curve, *J. High Energy Phys.* 12 (2016) 044, arXiv:1510.02100 [hep-th].
- [27] R. Klabbbers, S.J. van Tongeren, Quantum Spectral Curve for the eta-deformed AdS₅ × S⁵ superstring, *Nucl. Phys. B* 925 (2017) 252–318, arXiv:1708.02894 [hep-th].
- [28] N. Gromov, V. Kazakov, G. Korchemsky, S. Negro, G. Sizov, Integrability of conformal fishnet theory, *J. High Energy Phys.* 01 (2018) 095, arXiv:1706.04167 [hep-th].
- [29] T. Harmark, M. Wilhelm, to appear.
- [30] M. Alfimov, N. Gromov, G. Sizov, BFKL spectrum of $\mathcal{N} = 4$ SYM: non-zero conformal spin, arXiv:1802.06908 [hep-th], 2018.
- [31] L.A. Pando Zayas, D. Vaman, Strings in RR plane wave background at finite temperature, *Phys. Rev. D* 67 (2003) 106006, arXiv:hep-th/0208066 [hep-th].
- [32] B.R. Greene, K. Schalm, G. Shiu, On the Hagedorn behaviour of PP wave strings and $\mathcal{N} = 4$ SYM theory at finite R charge density, *Nucl. Phys. B* 652 (2003) 105–126, arXiv:hep-th/0208163 [hep-th].
- [33] R.C. Brower, D.A. Lowe, C.-I. Tan, Hagedorn transition for strings on pp waves and tori with chemical potentials, *Nucl. Phys. B* 652 (2003) 127–141, arXiv:hep-th/0211201 [hep-th].
- [34] G. Grignani, M. Orselli, G.W. Semenoff, D. Trancanelli, The superstring Hagedorn temperature in a pp wave background, *J. High Energy Phys.* 06 (2003) 006, arXiv:hep-th/0301186 [hep-th].
- [35] T. Harmark, M. Orselli, Matching the Hagedorn temperature in AdS/CFT, *Phys. Rev. D* 74 (2006) 126009, arXiv:hep-th/0608115 [hep-th].
- [36] T. Harmark, M. Orselli, Spin matrix theory: a quantum mechanical model of the AdS/CFT correspondence, *J. High Energy Phys.* 11 (2014) 134, arXiv:1409.4417 [hep-th].
- [37] D. Yamada, L.G. Yaffe, Phase diagram of $\mathcal{N} = 4$ super-Yang–Mills theory with R-symmetry chemical potentials, *J. High Energy Phys.* 09 (2006) 027, arXiv:hep-th/0602074 [hep-th].

- [38] T. Harnack, M. Orselli, Quantum mechanical sectors in thermal $\mathcal{N} = 4$ super Yang–Mills on $\mathbb{R} \times S^3$, Nucl. Phys. B 757 (2006) 117–145, arXiv:hep-th/0605234 [hep-th].
- [39] R. Suzuki, Refined counting of necklaces in one-loop $\mathcal{N} = 4$ SYM, arXiv:1703.05798 [hep-th], 2017.
- [40] M. Gomez-Reino, S.G. Naculich, H.J. Schnitzer, More pendants for Polya: two loops in the SU(2) sector, J. High Energy Phys. 07 (2005) 055, arXiv:hep-th/0504222 [hep-th].
- [41] J. Fokken, M. Wilhelm, One-loop partition functions in deformed $\mathcal{N} = 4$ SYM theory, J. High Energy Phys. 03 (2015) 018, arXiv:1411.7695 [hep-th].
- [42] D. Bombardelli, A. Cavaglià, D. Fioravanti, N. Gromov, R. Tateo, The full Quantum Spectral Curve for AdS_4/CFT_3 , J. High Energy Phys. 09 (2017) 140, arXiv:1701.00473 [hep-th].
- [43] N. Gromov, V. Kazakov, Z. Tsuboi, PSU(2, 2|4) character of quasiclassical AdS/CFT, J. High Energy Phys. 07 (2010) 097, arXiv:1002.3981 [hep-th].