



## Complete Genome Sequence of *Vibrio anguillarum* Nontailed Bacteriophage NO16

Kalatzis, Panos G.; Carstens, Alexander B.; Katharios, Pantelis; Castillo, Daniel; Hansen, Lars H.; Middelboe, Mathias

*Published in:*  
Microbiology resource announcements

*DOI:*  
[10.1128/MRA.00020-19](https://doi.org/10.1128/MRA.00020-19)

*Publication date:*  
2019

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY](#)

*Citation for published version (APA):*  
Kalatzis, P. G., Carstens, A. B., Katharios, P., Castillo, D., Hansen, L. H., & Middelboe, M. (2019). Complete Genome Sequence of *Vibrio anguillarum* Nontailed Bacteriophage NO16. *Microbiology resource announcements*, 8(15), 1-2. [e00020-19]. <https://doi.org/10.1128/MRA.00020-19>



# Complete Genome Sequence of *Vibrio anguillarum* Nontailed Bacteriophage NO16

Panos G. Kalatzis,<sup>a</sup> Alexander B. Carstens,<sup>b</sup> Pantelis Katharios,<sup>c</sup> Daniel Castillo,<sup>a</sup> Lars H. Hansen,<sup>b</sup> Mathias Middelboe<sup>a</sup>

<sup>a</sup>Marine Biological Section, University of Copenhagen, Helsingør, Denmark

<sup>b</sup>Department of Environmental Science, Aarhus University, Roskilde, Denmark

<sup>c</sup>Institute of Marine Biology, Biotechnology, and Aquaculture, Hellenic Centre for Marine Research, Heraklion, Crete, Greece

**ABSTRACT** A rare nontailed virus designated NO16 was isolated against *Vibrio anguillarum*, a major aquaculture pathogen for both fish and shellfish. Here, we announce the 10,594-bp genome sequence of *Vibrio* phage NO16 with a 23-gene content.

*Vibrio anguillarum* is a pathogenic bacterium of both cultured fish and shellfish (1, 2). The host of bacteriophage NO16 is *V. anguillarum* strain A023 (GenBank accession numbers CP010036 and CP010037), isolated from turbot in Spain (3).

Bacterial cells were cultured in liquid medium containing 0.5% tryptone (Difco), 0.1% yeast extract (Difco), and 2% sea salts (Sigma-Aldrich), and PFU were picked from the bacterial lawn produced by the double-agar-layer method (4). Phage DNA was extracted following the protocol listed in reference 5 and was sequenced from two single plaques using the MiSeq platform (Illumina, San Diego, CA, USA) as 2 × 250-bp paired-end reads according to the direct plaque sequencing (DPS) protocol (5) using 0.1% SDS instead of 1%. *De novo* assembly of the 548,174 reads (coverage, 1,014.74×) was performed using Genomic Workbench 9.5.3 (CLC Bio, Aarhus, Denmark). Briefly, reads were trimmed using the trim sequences tool (default settings), and overlapping reads were merged using the overlapping pairs tool (mismatch cost, 2; minimum score, 8; gap cost, 3; maximum unaligned end mismatches, 0) (6). The bacterial host's reads were removed using the map to reference tool. Circularity of the genome was confirmed by two independent approaches, (i) restriction enzyme digestion of the genome using different digestion sites followed by analysis of the size of the resulting segments and (ii) PCR amplification of specific sequences in the genome representing the site circularization (i.e., a sequence that would be amplified only if the genome was circular). Restriction enzyme *Cl*I (restriction site ATCGAT) produced two DNA bands on an agarose gel, whereas PCR with specific primers (forward, TGCCGGACAGAATCGAA CTC; reverse, ATGCGGAGGACACGACATGA) amplified, as expected, a 625-bp DNA fragment between the phage's genomic ends. Hence, bacteriophage NO16 has a 10,594-bp circular double-stranded DNA (dsDNA) genome with a GC content of 47.4%.

The 23 genes of the bacteriophage were predicted by Glimmer 3 (7), and they were then annotated with Rapid Annotations using Subsystems Technology (RAST) (8) and protein fold recognition server Phyre2 (9). DNA-binding protein (gene 6), *S*-adenosylhomocysteine hydrolase (gene 7), phage protein (gene 11), double jelly roll (DJR) capsid protein (gene 19), and ATPase (gene 21) are the 5 genes with some attributed function, whereas the remaining 18 open reading frames (ORFs) are hypothetical proteins. The presence of gene 19 classifies NO16 in the lineage of DJR viruses, of which very few marine members have been characterized so far (10, 11). In NCBI, the genome of NO16 was found in chromosome II of *V. anguillarum* strains 87-9-116, NB10, and VIB18 (query coverages of 98%, 98%, and 52%, percent

**Citation** Kalatzis PG, Carstens AB, Katharios P, Castillo D, Hansen LH, Middelboe M. 2019. Complete genome sequence of *Vibrio anguillarum* nontailed bacteriophage NO16. *Microbiol Resour Announc* 8:e00020-19. <https://doi.org/10.1128/MRA.00020-19>.

**Editor** Catherine Putonti, Loyola University Chicago

**Copyright** © 2019 Kalatzis et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mathias Middelboe, [mmiddelboe@bio.ku.dk](mailto:mmiddelboe@bio.ku.dk).

**Received** 10 February 2019

**Accepted** 15 March 2019

**Published** 11 April 2019

identities of 99.65%, 99.65%, and 99.91%, and GenBank accession numbers CP010045, LK021129, and CP011437, respectively), suggesting that it follows a temperate life cycle, although no integrase gene has been identified in its genome. With the signature gene of the DJR viral lineage (gene 19) as a query, the most closely related phage genomes are those of PM2 (12–14) and Cr39582 (11) (query coverages of 94% and 94%; percent identities of 35.80% and 35.16%, and GenBank accession numbers NC\_000867 and MG966533, respectively). Additionally, the DJR genes of several *Vibrio* phages recently published by Kauffman and colleagues (10) have similarities of 33% and below at query coverages of 96 to 97%.

**Data availability.** The genome sequence of bacteriophage NO16 was submitted to GenBank under the accession number MH730557, whereas the raw reads were uploaded to the European Nucleotide Archive (ENA) under the accession number PRJEB30917.

## ACKNOWLEDGMENTS

This work was supported by grants from The Danish Directorate for Food, Fisheries, and Agri Business (ProAqua, project number 09-072829), the Danish Research Council for Independent Research (project number DFF-7014-00080), and the Greek National Strategic Reference Framework 2007–2013 (cofunded by the European Social Fund and Greek National Funds [FISHPHAGE] project 131).

## REFERENCES

- Rønneseth A, Castillo D, D'Alvise P, Tønnesen Ø, Haugland G, Grotkjaer T, Engell-Sørensen K, Nørremark L, Bergh Ø, Wergeland HI, Gram L. 2017. Comparative assessment of *Vibrio* virulence in marine fish larvae. *J Fish Dis* 40:1373–1385. <https://doi.org/10.1111/jfd.12612>.
- Frans I, Michiels CW, Bossier P, Willems KA, Lievens B, Rediers H. 2011. *Vibrio anguillarum* as a fish pathogen: virulence factors, diagnosis and prevention. *J Fish Dis* 34:643–661. <https://doi.org/10.1111/j.1365-2761.2011.01279.x>.
- Castillo D, Alvise PD, Xu R, Zhang F, Middelboe M, Gram L. 2017. Comparative genome analyses of *Vibrio anguillarum* strains reveal a link with pathogenicity traits. *mSystems* 2:e00001-17. <https://doi.org/10.1128/mSystems.00001-17>.
- Kutter E. 2009. Phage host range and efficiency of plating. *Methods Mol Biol* 501:141–149. [https://doi.org/10.1007/978-1-60327-164-6\\_14](https://doi.org/10.1007/978-1-60327-164-6_14).
- Kot W, Vogensen FK, Sørensen SJ, Hansen LH. 2014. DPS: a rapid method for genome sequencing of DNA-containing bacteriophages directly from a single plaque. *J Virol Methods* 196:152–156. <https://doi.org/10.1016/j.jviromet.2013.10.040>.
- Carstens AB, Kot W, Lametsch R, Neve H, Hansen LH. 2016. Characterisation of a novel enterobacteria phage, CAjan, isolated from rat faeces. *Arch Virol* 161:2219–2226. <https://doi.org/10.1007/s00705-016-2901-0>.
- Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23: 673–679. <https://doi.org/10.1093/bioinformatics/btm009>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Pyre2 Web portal for protein modelling, prediction, and analysis. *Nat Protoc* 10:845–858. <https://doi.org/10.1038/nprot.2015.053>.
- Kauffman KM, Hussain FA, Yang J, Arevalo P, Brown JM, Chang WK, Vaninsberghe D, Elsherbini J, Sharma RS, Cutler MB, Kelly L, Polz MF. 2018. A major lineage of non-tailed dsDNA viruses as unrecognized killers of marine bacteria. *Nature* 554:118–122. <https://doi.org/10.1038/nature25474>.
- Leigh BA, Breitbart M, Oksanen HM, Bamford DH, Dishaw LJ. 2018. Genome sequence of PM2-like phage Cr39582, induced from a *Pseudoalteromonas* sp. isolated from the gut of *Ciona robusta*. *Genome Announc* 6:e00368-18. <https://doi.org/10.1128/genomeA.00368-18>.
- Espejo RT, Canelo ES. 1968. Properties of bacteriophage PM2: a lipid-containing bacterial virus. *Virology* 34:738–747. [https://doi.org/10.1016/0042-6822\(68\)90094-9](https://doi.org/10.1016/0042-6822(68)90094-9).
- Männistö RH, Kivelä HM, Paulin L, Bamford DH, Bamford JKH. 1999. The complete genome sequence of PM2, the first lipid-containing bacterial virus to be isolated. *Virology* 262:355–363. <https://doi.org/10.1006/viro.1999.9837>.
- Espejo RT, Canelo ES, Sinsheimer RL. 1969. DNA of bacteriophage PM2: a closed circular double-stranded molecule. *Proc Natl Acad Sci U S A* 63:1164–1168. <https://doi.org/10.1073/pnas.63.4.1164>.