



Dietary risk factors for inflammatory bowel diseases in a high-risk population
Results from the Faroese IBD study

Hammer, T; Lophaven, S Nymand; Nielsen, K Rubek; Petersen, M Skaalum; Munkholm, P; Weihe, P; Burisch, J; Lyng, E

Published in:
United European Gastroenterology Journal

DOI:
[10.1177/2050640619852244](https://doi.org/10.1177/2050640619852244)

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/)

Citation for published version (APA):
Hammer, T., Lophaven, S. N., Nielsen, K. R., Petersen, M. S., Munkholm, P., Weihe, P., ... Lyng, E. (2019). Dietary risk factors for inflammatory bowel diseases in a high-risk population: Results from the Faroese IBD study. *United European Gastroenterology Journal*, 7(7), 924-932. <https://doi.org/10.1177/2050640619852244>

Dietary risk factors for inflammatory bowel diseases in a high-risk population: Results from the Faroese IBD study

T Hammer^{1,2} , S Nymand Lophaven³, K Rubek Nielsen⁴,
M Skaalum Petersen^{1,5}, P Munkholm⁶, P Weihe^{1,5}, J Burisch⁶ and E Lynge²

United European Gastroenterology Journal
2019, Vol. 7(7) 924–932
© Author(s) 2019



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2050640619852244
journals.sagepub.com/home/ueg



Abstract

Background: The Faroe Islands currently have the highest recorded inflammatory bowel disease (IBD) incidence in the world.

Objective: This study investigated environmental risk factors for IBD in the Faroese population.

Methods: Environmental exposure data including lifestyle risk factors and neurotoxicants collected for over 30 years were retrieved from the Children's Health and the Environment in the Faroes (CHEF) cohorts including mainly mother-child pairs, with exposure data collected from pregnant mothers. For lifestyle risk factors, the incidence of IBD and ulcerative colitis (UC) was calculated as the rate ratio (RR) with 95% confidence intervals (CI) in exposed versus non-exposed persons. For neurotoxicants RR was calculated for persons with high versus low exposure.

Results: Six cohorts included 5698 persons with complete follow-up data and at least one exposure, and 37 were diagnosed with IBD. For pilot whale/blubber, the RR was 1.02 (95% CI, 0.48–2.18); RR of 1.01 for fish (95% CI, 0.35–2.91); and of the pollutants studied, a statistical significantly increased risk was found for 1,1,1,-trichloro-2,2-bis-(p-chlorophenyl) ethane (p,p'-DDT); RR 3.04 (95% CI, 1.12–8.30). RRs were 1.96 (95% CI, 1.03–3.73) for smoking and 1.10 (95% CI, 0.55–2.19) for alcohol intake.

Conclusion: The high IBD incidence is unlikely to be caused by special dietary habits or by environmental pollutants.

Keywords

Inflammatory bowel diseases, Faroe Islands, environmental risk factors

Received: 17 March 2019; accepted: 28 April 2019

Key summary

- The incidence of IBD is increasing worldwide
- The Faroe Islands currently have the highest recorded IBD incidence in the world
- Environmental factors play a key role
- We have previously reported on the importance of environmental factors in the development of IBD and especially of UC in Faroese migrants to Denmark

Main findings of our study

- To our knowledge, this is the first study to assess associations between several environmental toxicants and IBD

¹Department of Occupational Medicine and Public Health, The Faroese Hospital System, Tórshavn, Faroe Islands

²Nykøbing Falster Hospital, University of Copenhagen, Copenhagen, Denmark

³Larix A/S, Herlev, Denmark

⁴Medical Centre, National Hospital, Tórshavn, Faroe Islands

⁵Centre of Health Sciences, University of the Faroe Islands, Tórshavn, Faroe Islands

⁶Department of Gastroenterology, North Zealand Hospital, Frederikssund, Denmark

Corresponding author:

Turid Hammer, Department of Occupational Medicine and Public Health, Sigmundargøta 5, FO-100 Tórshavn, Faroe Islands.

Email: tuha@sund.ku.dk

- The special Faroese dietary habits and the environmental pollutants in this food is unlikely to have caused the record high IBD incidence
- Statistically significant associations were found between smoking and IBD and for one environmental toxicant (p,p'-DDT)

Introduction

Inflammatory bowel disease (IBD) is a chronic inflammatory and incurable condition in the gastrointestinal tract of unknown aetiology. There are two main subgroups of IBD; ulcerative colitis (UC) and Crohn's disease (CD). Currently, IBD incidence is increasing worldwide, and IBD is now considered to be a global disease.¹ The Faroe Islands in the North Atlantic is no exception. This small country with 50,000 inhabitants was in 2010 identified in the European Epi-IBD inception cohort study to have the highest IBD incidence in the world with an age standardised (European Standard Population, ESP) incidence rate of 81.5 per 100,000,² and the incidence has increased over time.³

The risk of IBD is known to have both genetic and environmental components, as it is an abnormal and dysregulated immune response occurring in genetically susceptible individuals. Although multiple susceptibility genes have been identified, genetics does not explain the drastic increase in the incidence of IBD.⁴ A hereditary predisposition for IBD may exist in the genetically homogeneous and isolated Faroese population, where genetically inherited diseases such as glycogen storage disease type IIIA,⁵ carnitine transporter deficiency⁶ and mitochondrial encephalomyopathy⁷ are frequent.

On environmental factors, it is known that children of parents, who migrate from a country with a low IBD prevalence to one with a high prevalence, adopt the IBD risk of their new home country.⁸ The reverse situation is present when Faroese people migrate to Denmark, where the IBD incidence is lower than in the Faroe Islands. First-generation Faroese immigrants to Denmark had a higher IBD incidence than Danes and the excess risk derived solely from UC and was restricted to the first 10 years of residence in Denmark, indicating the importance of environmental factors in the development of IBD and especially of UC.⁹

The evidence on modifiable environmental factors is inconclusive. Therefore it is difficult to undertake preventive measures and no clear guidance is available from the gastroenterology societies.¹⁰ Substantial data on environmental risk factors have been collected for more than 30 years within the Children's Health and the Environment in the Faroes (CHEF) project, established to assess impact on health of pollutants in the marine diet.^{11,12} On this background, the CHEF project enabled a unique search for

environmental risk factors for IBD in this remotely located, high-risk population.

Materials and methods

Children's Health and the Environment in the Faroes (CHEF) project

In 1985, Grandjean and Weihe investigated women aged 20 to 50 years in a small Faroese fishing settlement. The aim was to measure mercury exposure from marine food in blood samples. Due to the findings of increased mercury concentrations in these women, a larger nationwide study of mothers and their newborn children was initiated, marking the beginning of the CHEF project.¹³ In total, four mother-child cohorts have been established since 1986 and one cohort of pregnant women. The project has been carried out at the Department of Occupational Medicine and Public Health within the Faroese Hospital System, and the project currently includes over 2,500 Faroese mother-child pairs. In the present observational and prospective study, the study population consisted of the four mother-child cohorts, and grandmothers and fathers¹⁴ that were recruited within the last mother-child cohort. Furthermore, the cohort of only pregnant women¹² was included alongside one additional cohort investigating semen quality in young Faroese men, born between 1981 and 1984, and their mothers (Table 1).¹⁵

In the Faroe Islands, hunting of pilot whale for human consumption is a tradition going far back in time. The pilot whale is a toothed whale and thus a carnivorous marine mammal eating squid and fish, and in modern times contaminated with high levels of pollutants accumulated in the marine food chain.¹⁶ The measured environmental toxicants include methylmercury (MeHg), persistent organic pollutants (POPs) including polychlorinated biphenyls (PCBs) and perfluorinated compounds (PFCs).

The first mother-child pairs were recruited in 1986–87, the last pairs, and the young men and their mothers, in 2007–09. The cohorts represented up to 75% of all Faroese births in the recruitment periods (Table 1). Number of participants by gender and cohort is provided in supplementary Tables 1 (any registration) and 2 (first registration only). The CHEF project has gathered extensive exposure information from questionnaires answered by the mothers at recruitment during pregnancy or shortly after delivery and

Table 1. Overview of the Faroese cohorts included in this study.

Cohort name	Recruitment period	Percentage of all births	Entry date	Children	Adults	Total
Cohort 1 ^a	March 1986–1987 (end of)	75%	Child's birthdate	1001	980	1981
Cohort 2 ^a	March 1994–March 1995	64%	Child's birthdate	189	189	378
Cohort 3 ^b	Nov. 1997–March 2000	60%	Child's birthdate	654	638	1292
Cohort 4 ^c	Oct. 2000–Sept 2001	23% ^e	1 April 2001	-	166	166
Cohort 5 ^a	Oct. 2007–April 2009	73%	Child's birthdate	488	504	992
- Fathers from cohort 5 ^d	Feb. 2009–Feb. 2010		Child's birthdate	-	282	282
- Maternal grandmothers from cohort 5 ^d	Oct. 2007–April 2009		Child's birthdate	-	337	337
- Paternal grandmothers from cohort 5 ^d	Oct. 2007–April 2009		Child's birthdate	-	201	201
Young men	Feb. 2007–Feb. 2009		1 Feb. 2008	263	207	470
Total ^f				2595	3504	6099
Unique records ^g						5773

^aCohort 1, 2 and 5 included singleton births only.

^bCohort 3 included twin births also.

^cCohort 4 recruited pregnant women only.

^dCohort 5 enrolled fathers alongside maternal and paternal grandmothers.

^eCohort 4 did not intend to include all births within the period, thus the lower percentage.

^fA given person could be included more than one time.

^gA given person included only first time.

biological samples including umbilical cord blood, whole blood, maternal hair and breast milk.

Data on environmental exposures in the Faroe Islands

In the present study, we included exposure data on smoking, alcohol consumption and dietary intake of pilot whale/blubber and fish retrieved from self-reported questionnaires that were filled in at recruitment, though the timing of the exposure status varied across cohorts. For mothers, their exposure status refers to exposure during pregnancy; for children their exposure in uterus; for fathers their exposure shortly before the pregnancy of their partners. For grandmothers their exposure for smoking and alcohol consumption refers to the time prior to their pregnancy with the now pregnant daughter/son's pregnant partner, while their intake of traditional Faroese foods refers to exposure during pregnancy. For the young men their exposure is at the time of recruitment; and for mothers of young men their exposure prior to their pregnancy for smoking and alcohol consumption, while intake of traditional Faroese foods refers to exposure during pregnancy. The phrasing of the exposure questions varied slightly across the six cohorts, and in order to standardize the data, all answers were dichotomized as yes/high or no/low to usage/intake (Table 2).

Furthermore, data were included on toxicants measured in the blood samples taken at the time of recruitment: POPs (PCBs: 1,1-dichloro-2,2-bis(p-chlorophenyl) ethane (p,p'-DDE); 1,1,1-trichloro-2,2-bis(p-chlorophenyl) ethane (p,p'-DDT); β -hexachlorocyclohexane (β -HCH)); perfluorooctanoic acid (PFOA); perfluorooctane sulfonate (PFOS); perfluorononanoic acid (PFNA); perfluorodecanoic acid (PFDA); perfluorohexane sulfonate (PFHxS); perfluorooctane sulphonamide (Total FOSA). MeHg was measured in hair and blood samples. In order to ensure that potential differences in the results were not due to the measurement method, both measurements were included. Because PCB is a mixture of several congeners, the Σ PCB was calculated as the sum of PCB-138, PCB-153 and PCB-180 multiplied by 2, as the sum of these represent close to 50% of the total PCB concentration.¹⁷

The cohorts were followed up via the Faroese National Population Register from the date of their first recruitment (entry date) until date of emigration, death, diagnosis of IBD or 31 May 2017 (end of follow-up). Incident cases of IBD among the cohort participants were identified within the nationwide IBD database established in the Faroese IBD study.³ All cases were clinically characterized according to the Montreal classification.¹⁸ For the purpose of the current study, data were updated until end of follow-up in

Table 2. Definition of exposure status for smoking, alcohol consumption and intake of pilot whale/blubber and fish.

Cohort members	Smoking		Alcohol		Pilot whale/blubber ^b		Fish	
	Exposed answered Yes to	Non-exposed answered No to	Exposed answered Yes to	Non-exposed answered No to	Exposed ate more than one	Non-exposed ate less than one	Exposed ate more than one	Non-exposed ate less than one
Mothers in cohort 1, ^a 2, ^a 3, ^a 4, ^a 5 ^a	Smoking during pregnancy	No to	Alcohol consumption during pregnancy	No to	Dinner per month during pregnancy	Dinner per month during pregnancy	Dinner per week during pregnancy	Dinner per week during pregnancy
Fathers in cohort 5	Smoking shortly before pregnancy of their partner	No to	Alcohol consumption shortly before pregnancy of their partner	No to	Dinner per year at recruitment	Dinner per year at recruitment	Dinner per week at recruitment	Dinner per week at recruitment
Maternal and paternal grandmothers in cohort 5 and mothers of young men	Smoking shortly before pregnancy	No to	Alcohol consumption shortly before pregnancy	No to	Dinner per year during pregnancy	Dinner per year during pregnancy	Dinner per week during pregnancy	Dinner per week during pregnancy
Young men	Current smoking	No to	Current alcohol consumption	No to	Dinner per year at recruitment	Dinner per year at recruitment	Dinner per week at recruitment	Dinner per week at recruitment

^aExposure status of the children in cohorts 1, 2, 3 and 5 equates to that of the mothers as questionnaires were answered during pregnancy.

^bFor the analysis the different time scales were standardized.

cooperation with the Medical Centre at the National Hospital of the Faroe Islands. Definition and classification of IBD cases in the Faroese IBD study have been described previously.³

Analysis

For a given exposure, we estimated the incidence of IBD among exposed persons as compared with the incidence of IBD among non-exposed. A given cohort member was included in the analysis from her/his date of first recruitment into CHEF, and followed up until date of IBD diagnosis, death, emigration or end of the study. A re-immigrated person was re-included when she/he returned to the Faroe Islands. The rate ratio (RR) and the 95% confidence interval (95% CI) of IBD incidence and of UC incidence between exposed and non-exposed cohort members were estimated with Poisson regression controlling for age (0–19, 20–39, 40–59, 60–79, 80+) and calendar period (before 1 January 2000 and after). Gender was not controlled for, as the IBD incidence is identical for men and women in the Faroe Islands.³ Data on the environmental toxicants were divided into tertiles with RR and 95% CI calculated as comparison of the incidence between the highest and the lowest tertiles. Geometric mean and interquartile range for each exposure group were calculated. SAS software version 9.4 and R version 3.4.3 were used for the statistical analysis.

Ethics

This is a register-based study approved by the Faroese Committee on Biomedical Research Ethics and the Faroese Data Inspection Agency, J. nr. 16/00037. No contact was made to the participating cohort members, IBD patients or relatives. All cohort members signed an informed consent in order to participate in any of the investigations conducted at the Department of Occupational Medicine and Public Health.

Results

In total, the six cohorts included 6099 records of which 5773 were unique persons with 5698 having complete follow-up information on migration and death and data on at least one type of exposure (Figure 1). During the follow-up period, 37 individuals within the age range of 11–67 years were diagnosed with IBD, of which 32 had UC and 5 had CD. None of the CD cases had perianal disease. The Montreal classifications of the IBD cases are shown in supplementary Table 3.

The six cohorts contributed a total of 96,507.6 person-years (PY). The RRs and 95% CIs for the associations between the environmental risk factors and the




Figure 1. Flow chart of number of participants in the cohorts with information on smoking, alcohol consumption and food intake.

incidence of IBD and UC are listed in Table 3. Only smoking was statistically significantly associated with IBD. Exposure data on smoking were complete for 5670 persons. Comparison of the 1935 smokers (33,132.2 PY) with 19 cases of IBD with the 3735 non-smokers (63,072.6 PY) with 18 cases of IBD resulted in a RR of 1.96 (95% CI, 1.03–3.73).

Table 4 illustrates the RRs and 95% CIs for the associations between the environmental toxicants and the incidence of IBD and UC. The only toxicant statistically significantly associated with IBD was p,p'-DDT. In total, 1721 persons had high exposure and 1712 had low exposure to this pollutant with a RR of 3.04 (95% CI, 1.12–8.30). We tested also the association for p,p'-DDT and UC; RR of 3.60 (95% CI, 1.20–10.82). No statistically significant association was found for other compounds on the risk of UC (data not shown). Not all environmental toxicants were measured in every cohort.

Discussion

The incidence of IBD in the Faroe Islands is the highest in the world, and a strong influence of environmental factors on disease risk has been demonstrated in

Faroese migrants to Denmark. Therefore, further studies on environmental exposures in Faroese inhabitants prior to the emergence of IBD are important. The present study did not indicate the high risk of IBD in the Faroe Islands to derive from the intake of traditional Faroese food in the form of pilot whale/blubber and fish. Neither did we find clear associations between the measured toxicants in seafood and IBD. Only one compound (p,p'-DDT) was statistically significantly associated with an increased risk of IBD and UC, but with the many measured associations this could be a chance finding. For toxicants, most risk estimates were actually below 1, and one could speculate that this could be an immunosuppressive effect, in line with the prior finding of a reduced antibody response to childhood vaccines in children exposed to PCBs.¹⁹

Alcohol consumption was not associated with the risk of IBD and/or UC. The risk of IBD was, however, statistically significantly elevated in smokers, a tendency that remained for UC only, though based on small numbers. Data were too scarce for separate analysis of CD.

Although the Faroe Islands are located in Northern Europe, this country differs from the rest of Europe in terms of its traditional food of pilot whale and blubber.

Table 3. Number of persons (N), person-years (PY), IBD and UC (cases) by exposure status; rate ratios (RRs) and 95% confidence intervals (95% CIs) for exposed persons as compared with non-exposed persons.

Exposure	Exposed persons			Non-exposed persons			Total			RR (95% CI)
	N	PY	Cases	N	PY	Cases	N	PY	Cases	
Smoking and IBD	1935	33,132.2	19	3735	63,072.6	18	5670 ^a	96,204.8	37	1.96 (1.03–3.73)
Smoking and UC	1935	33,138.2	16	3735	63,084.7	16	5670 ^a	96,222.9	32	1.85 (0.92–3.70)
Alcohol and IBD	1759	27,287.8	12	3902	68,804.0	25	5661 ^b	96,091.8	37	1.10 (0.55–2.19)
Alcohol and UC	1759	27,287.8	12	3902	68,821.9	20	5661 ^b	96,109.7	32	1.37 (0.67–2.80)
Pilot whale/blubber and IBD	3456	63,253.5	27	1564	21,801.2	9	5020 ^c	85,054.7	36	1.02 (0.48–2.18)
Pilot whale/blubber and UC	3456	63,268.4	23	1564	21,801.2	9	5020 ^c	85,069.6	32	0.86 (0.39–1.87)
Fish and IBD	4195	76,652.1	32	804	8115.5	4	4999 ^d	84,767.6	36	1.01 (0.35–2.91)
Fish and UC	4195	76,658.1	29	804	8124.5	3	4999 ^d	84,782.6	32	1.24 (0.37–4.14)

Missing data for:

^a28 persons;

^b37 persons;

^c678 persons;

^d699 persons.

Table 4. Geometric mean (GM), interquartile range (IQR), number of persons (N), person-years (PY) and IBD (cases) by toxicological compound; rate ratios (RRs) and 95% confidence intervals (95% CIs) for persons with high exposure vs. low.

Exposure	GM		IQR (25%–75%)		N		PY		Cases		RR (95% CI)
	Low	High	Low	High	Low	High	Low	High	Low	High	
ΣPCB ^a	0.270	2.063	0.20–0.42	1.45–2.61	1715	1561	27,991.9	28,145.2	9	11	1.27 (0.52–3.07)
p,p'-DDE ^a	0.093	1.062	0.07–0.16	0.71–1.41	1706	1584	27,165.8	29,671.1	9	9	0.97 (0.38–2.44)
p,p'-DDT ^a	0.002	0.064	0.002–0.003	0.036–0.096	1712	1721	27,365.6	38,146.7	5	18	3.04 (1.12–8.30)
β-HCH ^a	0.004	0.045	0.003–0.006	0.03–0.07	1691	1680	34,649.8	31,043.8	13	10	0.88 (0.38–1.99)
PFOA ^b	0.948	4.422	0.76–1.34	3.55–4.98	1457	1386	26,048.6	26,979.7	11	7	0.60 (0.23–1.56)
PFOS ^b	2.332	26.88	1.93–2.90	21.90–32.24	1494	1304	37,074.7	20,369.9	15	3	0.30 (0.08–1.07)
PFNA ^b	0.209	1.099	0.21–0.21	0.79–1.38	1815	1212	45,552.9	14,396.5	18	5	0.62 (0.22–1.76)
PFDA ^b	0.196	0.453	0.21–0.21	0.34–0.53	2373	1222	52,524.5	15,634.7	19	5	0.70 (0.25–1.95)
PFHxS ^b	0.184	0.722	0.20–0.21	0.49–0.94	1819	845	37,274.4	14,277.2	16	6	0.80 (0.31–2.07)
Total FOSA ^b	0.297	1.832	0.23–0.41	1.22–2.20	606	638	15,380.5	15,947.1	5	4	0.76 (0.20–2.84)
MeHg (blood) ^b	2.142	39.59	1.62–3.34	28.15–53.05	1038	1275	9755.7	31,927.3	6	10	0.84 (0.29–2.40)
MeHg (hair) ^c	0.636	6.931	0.47–1.00	4.87–9.05	1829	1769	19,887.7	40,029.7	11	15	0.81 (0.36–1.79)

Toxicants were measured in:

^aμg/g lipid;

^bμg/L;

^cμg/g hair.

Evidence suggests that a diet high in fruits, vegetables, fibre and omega-3 fatty acids is associated with a lower IBD risk.¹⁰ It is noteworthy to mention that pilot whale, blubber and fish to various extents contain omega-3-fatty acids, which may in part explain why these foods did not increase the risk of IBD.¹¹ The northern latitude of the Faroe Islands and thus the low exposure to sunlight is believed to result in lower

levels of vitamin D, although similar levels have been found in Denmark, Norway and Finland.²⁰ Vitamin D deficiency is common in IBD patients and considered an important component in the development of IBD.²¹ In the present study we did not include data on vitamin D status, previously reported from cohort 3 and 5.²² The high intake of seafood in the studied cohorts may compensate for the low sunlight exposure.

Nevertheless, since January 2018, Faroese milk with 0.5% fat has been enriched with vitamin D.

To our knowledge, this is the first study assessing the associations between environmental toxicants and IBD. An association between PFOA and UC has been reported from an occupationally exposed cohort,^{23,24} but we did not see this association. A possible explanation for the discrepancy between our study and the US study is the lower exposure in the Faroese cohorts. Another explanation is differences in assessment of exposure: Steenland et al.²³ combined occupational and residential serum estimates, while exposure in the Faroese cohorts was measured directly from each individual, showing current exposure, not cumulative or estimated exposure. In our study, only p,p'-DDT was associated with IBD and UC. As no other study has investigated p,p'-DDT and IBD, this positive finding needs replication. However, the lack of associations between environmental toxicants and IBD was not surprising, as we found no association between intake of seafood and IBD.

It was surprising that intake of traditional food was not associated with an increased risk of IBD. The Faroese immigrants to Denmark had an excess IBD incidence during their first 10 years in Denmark, but thereafter their incidence resembled that of Danes.⁹ One could have expected this change to coincide with a decreased intake of the traditional Faroese food. However, the IBD incidence in the Faroe Islands increased from the 1960s to the 2010s,³ which is a period where the Faroese diet became increasingly westernized. Taking the combined evidence into account, it seems reasonable to hypothesize that the high incidence of IBD in the Faroe Islands derives from recent dietary or other habits not found in Denmark, but at present we are unable to pinpoint specific factors.

The lack of an association between alcohol consumption and the risk of IBD was confirmed in the recently reported European prospective cohort study (EPIC).²⁵ A meta-analysis on beverage consumption also found no association between alcohol intake and the risk of UC,²⁶ but the role of alcohol in the aetiology of IBD is inconclusive and widely debated, as some studies found an increased IBD risk in relation to alcohol consumption.²⁷

The association between smoking and IBD is complex. CD is four-fold more frequent in current smokers than in never smokers, and former smokers have an increased prevalence of CD. Smoking cessation is recommended to CD patients, as continued smoking often complicates disease course. Smoking has, on the other hand, a protective effect on the development of UC, and smoking cessation in UC will induce a more severe disease course.¹⁰ The majority of the studies on smoking and IBD are case-control studies, where

recall- and reporting biases cannot be excluded. The evidence on the long-term effect of smoking on CD and UC is limited, but one prospective study from the Nurses Health Study found an increased risk of UC among former smokers with a hazard ratio of 1.56 (95% CI, 1.26–1.93).²⁸ When we combined their hazard ratios for current and former smokers, the estimate was 1.32, well within the 95% confidence interval of our result for smoking; RR 1.85 (95% CI, 0.92–3.70). Further prospective cohort data on smoking and risk of UC are highly warranted. The Faroe Islands continue to have the highest percentage of daily smokers in the Nordic countries at 20% for men and 24% for women.²⁹ No free of charge tobacco cessation courses are available in the country.

Strengths and limitations

We investigated risk factors for IBD in the largest data set on human environmental exposures available from the Faroe Islands with individual follow-up of cohort members for death and emigrations from the population register, and with individual linkage to the nationwide Faroese IBD database.³ From previous work we had established that Faroese re-immigrate repeatedly, hence the need for a precise measurement of person-years.⁹ The participants in this study can be regarded as representative of the general population as they were unselected and resided throughout the country, though individuals from the capital region were overrepresented. The only exception was cohort 2 where participants resided away from the capital area of Tórshavn.¹²

Data on lifestyle risk factors were derived from self-reported questionnaires. For 49% of the cohorts (mothers and fathers), exposure data referred to exposure at time of recruitment, but for 40% (children) the data referred to exposure before birth; and for the remaining 11% (grandmothers/fathers; mothers of young men) to time of pregnancy with their now adult daughter/son. While these time differences probably do not affect the data on dietary habits, they could have affected the smoking and alcohol data. The use of the mother's intake of traditional food during pregnancy as a proxy for the child's exposure is also a limitation of the study. Furthermore, it was not possible to control for various other environmental factors that the cohort members might have been exposed to in the Faroe Islands or abroad. A final limitation was the limited number of IBD cases due to the still relatively young age of the cohort members recruited as newborns as these had not yet reached the age at which IBD may develop. Exposure to environmental toxicants is commonly found in the Faroese population why it can be expected that no one is truly unexposed.

We therefore compared individuals of the lowest exposure tertile with those in the highest (Table 4).

Conclusion

This study combined comprehensive data on marine environmental exposures with that of IBD incidence in the Faroe Islands. The study was undertaken within an ideal setting for investigating the impact of the environment on IBD risk in a location that has seen a dramatic increase in the incidence of IBD within the same decades as those in which the studied cohorts were recruited. The study showed that neither the special dietary habits in the form of pilot whale and blubber nor the environmental pollutants in this food were likely to explain the high incidence of IBD in the Faroe Islands.

Acknowledgements

TH would like to thank colleagues at the Department of Occupational Medicine and Public Health for their help and highly valued encouragements.

Author contributions

All authors contributed to this work. PW and MSP collected the data. TH and EL designed the study, obtained funding and drafted the manuscript. SNL performed the statistical analysis and contributed to the interpretation. TH, SNL, KRN, MSP, PM, PW, EL and JB critically revised and approved the final manuscript.

Conflict of interest

JB reports personal fees from AbbVie, personal fees from Janssen-Cilag, personal fees from Celgene, personal fees from MSD, personal fees from Pfizer, grants and personal fees from Takeda, outside the submitted work.

Ethics approval

This is a register-based study approved by the Faroese Committee on Biomedical Research Ethics and the Faroese Data Inspection Agency, J. nr. 16/00037. No contact was made to the participating cohort members, IBD patients or relatives.

Funding

This work was supported by the Faroese Research Council, the European Crohn's and Colitis Organisation (ECCO), Beckett Foundation, the Danish Colitis-Crohn Patients Organisation (CCF) and Aage & Johanne Louis-Hansens Foundation (to TH).

Informed consent

All cohort members signed an informed consent in order to participate in any of the investigations conducted at the Department of Occupational Medicine and Public Health.

ORCID iD

T Hammer  <https://orcid.org/0000-0002-5242-955X>

References

1. Ng SC, Shi HY, Hamidi N, et al. Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet* 2017; 390: 2769–2778.
2. Burisch J, Pedersen N, Čuković-Čavka S, et al. East-West gradient in the incidence of inflammatory bowel disease in Europe: the ECCO-EpiCom inception cohort. *Gut* 2014; 63: 588–597.
3. Hammer T, Nielsen KR, Munkholm P, et al. The Faroese IBD study: incidence of inflammatory bowel disease across 54 years of population-based data. *J Crohns Colitis* 2016; 10: 934–942.
4. Molodecky NA and Kaplan GG. Environmental risk factors for inflammatory bowel disease. *Gastroenterol Hepatol* 2010; 6: 339–346.
5. Santer R, Kinner M, Steuerwald U, et al. Molecular genetic basis and prevalence of glycogen storage disease type IIIA in the Faroe Islands. *Eur J Hum Genet* 2001; 9: 388–391.
6. Poulsen SD, Lund AM, Christensen E, et al. [Carnitine transporter deficiency is a hereditary disease with a high incidence in the Faroe Islands]. *Ugeskr Laeger* 2012; 174: 1217–1219.
7. Ostergaard E, Hansen FJ, Sorensen N, et al. Mitochondrial encephalomyopathy with elevated methylmalonic acid is caused by SUCLA2 mutations. *Brain* 2007; 130: 853–861.
8. Kaplan GG. The global burden of IBD: from 2015 to 2025. *Nat Rev Gastroenterol Hepatol* 2015; 12: 720–727.
9. Hammer T, Lophaven SN, Nielsen KR, et al. Inflammatory bowel diseases in Faroese-born Danish residents and their offspring: further evidence of the dominant role of environmental factors in IBD development. *Aliment Pharmacol Ther* 2017; 45: 1107–1114.
10. Maaser C, Langholz E, Gordon H, et al. European Crohn's and Colitis organisation topical review on environmental factors in IBD. *J Crohns Colitis* 2017; 11: 905–920.
11. Grandjean P, Weihe P, White RF, et al. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicol Teratol* 1997; 19: 417–428.
12. Weihe P and Grandjean P. Cohort studies of Faroese children concerning potential adverse health effects after the mothers' exposure to marine contaminants during pregnancy. *Acta Vet Scand* 2012; 54: S7.
13. Grandjean P, Weihe P, Jørgensen PJ, et al. Impact of maternal seafood diet on fetal exposure to mercury, selenium, and lead. *Arch Environ Health* 1992; 47: 185–195.
14. Petersen MS, Halling J, Weihe P, et al. Spermatogenic capacity in fertile men with elevated exposure to polychlorinated biphenyls. *Environ Res* 2015; 138: 345–351.
15. Halling J, Petersen MS, Jørgensen N, et al. Semen quality and reproductive hormones in Faroese men: a

- cross-sectional population-based study of 481 men. *BMJ Open* 2013; 3: e001946.
16. Julshamn K, Andersen A, Ringdal O, et al. Trace elements intake in the Faroe Islands. I. Element levels in edible parts of pilot whales (*Globicephalus meleanus*). *Sci Total Environ* 1987; 65: 53–62.
 17. Grandjean P, Weihe P, Needham LL, et al. Relation of a seafood diet to mercury, selenium, arsenic, and polychlorinated biphenyl and other organochlorine concentrations in human milk. *Environ Res* 1995; 71: 29–38.
 18. Silverberg MS, Satsangi J, Ahmad T, et al. Toward an integrated clinical, molecular and serological classification of inflammatory bowel disease: report of a Working Party of the 2005 Montreal World Congress of Gastroenterology. *Can J Gastroenterol* 2005; 19(Suppl A): 5A–36A.
 19. Heilmann C, Grandjean P, Weihe P, et al. Reduced antibody responses to vaccinations in children exposed to polychlorinated biphenyls. *PLoS Med* 2006; 3: e311.
 20. Dalgård C, Petersen MS, Schmedes AV, et al. High latitude and marine diet: vitamin D status in elderly Faroese. *Br J Nutr* 2010; 104: 914–918.
 21. Chetcuti Zammit S, Ellul P, Girardin G, et al. Vitamin D deficiency in a European inflammatory disease inception cohort: an Epi-IBD study. *Eur J Gastroenterol Hepatol* 2018; 30: 1297–1303.
 22. Dalgård C, Petersen MS, Steuerwald U, et al. Umbilical cord serum 25-hydroxyvitamin D concentrations and relation to birthweight, head circumference and infant length at age 14 days. *Paediatr Perinat Epidemiol* 2016; 30: 238–245.
 23. Steenland K, Zhao L, Winquist A, et al. Ulcerative colitis and perfluorooctanoic acid (PFOA) in a highly exposed population of community residents and workers in the mid-Ohio valley. *Environ Health Perspect* 2013; 121: 900–905.
 24. Steenland K, Zhao L and Winquist A. A cohort incidence study of workers exposed to perfluorooctanoic acid (PFOA). *Occup Environ Med* 2015; 72: 373–380.
 25. Bergmann MM, Hernandez V, Bernigau W, et al. No association of alcohol use and the risk of ulcerative colitis or Crohn's disease: data from a European Prospective cohort study (EPIC). *Eur J Clin Nutr* 2017; 71: 512–518.
 26. Nie JY and Zhao Q. Beverage consumption and the risk of ulcerative colitis: Systematic review and meta-analysis of epidemiological studies. *Medicine* 2017; 96: e9070.
 27. Yang XY, Wei WB, Zeng LR, et al. Controversial role of alcohol in the development of inflammatory bowel diseases. *Eur J Clin Nutr* 2018; 72: 304.
 28. Higuchi LM, Khalili H, Chan AT, et al. A prospective study of cigarette smoking and the risk of inflammatory bowel disease in women. *Am J Gastroenterol* 2012; 107: 1399–1406.
 29. Nomesco. Health Statistics in the Nordic countries 2017. 2017. <http://norden.diva-portal.org/smash/get/diva2:1148509/FULLTEXT05.pdf> (accessed 10 October 2018).