



The antiresorptive effect of GIP, but not GLP-2, is preserved in patients with hypoparathyroidism - a randomized crossover study

Skov-Jepesen, Kirsa; Hepp, Nicola; Oeke, Jannika; Hansen, Morten Steen; Jafari, Abbas; Svane, Maria Saur; Balenga, Nariman; Olson, John A; Frost, Morten; Kassem, Moustapha; Madsbad, Sten; Beck Jensen, Jens-Erik; Holst, Jens Juul; Rosenkilde, Mette Marie; Hartmann, Bolette

Published in:

Journal of Bone and Mineral Research

DOI:

[10.1002/jbmr.4308](https://doi.org/10.1002/jbmr.4308)

Publication date:

2021

Document version

Publisher's PDF, also known as Version of record

Document license:

[CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Citation for published version (APA):

Skov-Jepesen, K., Hepp, N., Oeke, J., Hansen, M. S., Jafari, A., Svane, M. S., Balenga, N., Olson, J. A., Frost, M., Kassem, M., Madsbad, S., Beck Jensen, J-E., Holst, J. J., Rosenkilde, M. M., & Hartmann, B. (2021). The antiresorptive effect of GIP, but not GLP-2, is preserved in patients with hypoparathyroidism - a randomized crossover study. *Journal of Bone and Mineral Research*, 36(8), 1448-1458. <https://doi.org/10.1002/jbmr.4308>

The Antiresorptive Effect of GIP, But Not GLP-2, Is Preserved in Patients With Hypoparathyroidism—A Randomized Crossover Study

Kirsa Skov-Jepesen,^{1,2} Nicola Hepp,³ Jannika Oeke,¹ Morten Steen Hansen,⁴ Abbas Jafari,⁵ Maria Saur Svane,³ Nariman Balenga,⁶ John A Olson Jr,⁶ Morten Frost,⁴ Moustapha Kassem,^{4,5} Sten Madsbad,³ Jens-Erik Beck Jensen,^{3,7} Jens Juul Holst,^{1,2} Mette Marie Rosenkilde,¹ and Bolette Hartmann^{1,2}

¹Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark

²Novo Nordisk Foundation Center for Basic Metabolic Research, University of Copenhagen, Copenhagen, Denmark

³Department of Endocrinology, Hvidovre University Hospital, Hvidovre, Denmark

⁴Molecular Endocrinology Unit (KMEB), Department of Endocrinology, Odense University Hospital, Odense, Denmark

⁵Department of Cellular and Molecular Medicine, Novo Nordisk Foundation Center for Stem Cell Biology (Danstem), University of Copenhagen, Copenhagen, Denmark

⁶Division of General and Oncologic Surgery, Department of Surgery, Marlene and Stewart Greenebaum Comprehensive Cancer Center, University of Maryland School of Medicine, Baltimore, MD, USA

⁷Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

ABSTRACT

Glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-2 (GLP-2) are gut hormones secreted postprandially. In healthy humans, both hormones decrease bone resorption accompanied by a rapid reduction in parathyroid hormone (PTH). The aim of this study was to investigate whether the changes in bone turnover after meal intake and after GIP- and GLP-2 injections, respectively, are mediated via a reduction in PTH secretion. This was tested in female patients with hypoparathyroidism given a standardized liquid mixed-meal test ($n = 7$) followed by a peptide injection test ($n = 4$) using a randomized crossover design. We observed that the meal- and GIP- but not the GLP-2-induced changes in bone turnover markers were preserved in the patients with hypoparathyroidism. To understand the underlying mechanisms, we examined the expression of the GIP receptor (GIPR) and the GLP-2 receptor (GLP-2R) in human osteoblasts and osteoclasts as well as in parathyroid tissue. The GIPR was expressed in both human osteoclasts and osteoblasts, whereas the GLP-2R was absent or only weakly expressed in osteoclasts. Furthermore, both GIPR and GLP-2R were expressed in parathyroid tissue. Our findings suggest that the GIP-induced effect on bone turnover may be mediated directly via GIPR expressed in osteoblasts and osteoclasts and that this may occur independent of PTH. In contrast, the effect of GLP-2 on bone turnover seems to depend on changes in PTH and may be mediated through GLP-2R in the parathyroid gland. © 2021 The Authors. *Journal of Bone and Mineral Research* published by Wiley Periodicals LLC on behalf of American Society for Bone and Mineral Research (ASBMR).

KEY WORDS: GIP; GLP-2; BONE TURNOVER; OSTEOCLASTS; OSTEOBLASTS; BIOCHEMICAL MARKERS OF BONE TURNOVER

Introduction

Gastrointestinal hormones secreted after meal ingestion have been suggested as mediators in the gut–bone axis.^(1–8) In particular, glucose-dependent insulinotropic

polypeptide (GIP) and glucagon-like peptide 2 (GLP-2) seem to be potent inhibitors of bone resorption in humans.^(2–6,9–12) Thus, exogenously administered GIP as well as GLP-2 reduce bone resorption, as estimated from serum concentrations of collagen type 1 C-terminal telopeptide (CTX), to 60% to 50% of baseline with a maximal effect reached after approximately 90 minutes

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Received in original form November 3, 2020; revised form March 17, 2021; accepted April 8, 2021; Accepted manuscript online April 14, 2021.

Address correspondence to: Bolette Hartmann, PhD, Department of Biomedical Sciences, The Panum Institute, University of Copenhagen, Building 12.2, Blegdamsvej 3, DK-2200 Copenhagen, Denmark. E-mail: bhartmann@sund.ku.dk

Additional Supporting Information may be found in the online version of this article.

Journal of Bone and Mineral Research, Vol. 36, No. 8, August 2021, pp 1448–1458.

DOI: 10.1002/jbmr.4308

© 2021 The Authors. *Journal of Bone and Mineral Research* published by Wiley Periodicals LLC on behalf of American Society for Bone and Mineral Research (ASBMR).

for GIP and 180 minutes for GLP-2.⁽¹²⁾ Importantly, the bone effects of both GIP and GLP-2 are of clinical relevance because individuals with altered GIP receptor (GIPR) activity due to a specific mutation (Glu354Gln[rs1800437]) that desensitizes the receptor⁽¹³⁾ have decreased bone mineral density (BMD) and a more than 50% increased risk of non-vertebral fractures,⁽¹⁴⁾ whereas exogenous GLP-2 has been found to increase BMD dose-dependently in postmenopausal women with low BMD.⁽⁵⁾ In addition to its antiresorptive effect, GIP seems to induce a small increase in bone formation as reflected by increased serum levels of procollagen type 1 N-terminal propeptide (P1NP).^(11,12,15) Thus, an important feature of GIP is its ability to uncouple bone resorption and bone formation, which are normally tightly coupled.⁽¹⁶⁾

At the cellular level, GIPR expression has been described in a murine osteoclast-like cell line⁽¹⁷⁾ and on human osteoblast-like cells,⁽¹⁸⁾ supporting the notion of a direct effect of GIP on bone cells. Expression of the GLP-2 receptor (GLP-2R) on bone cells has, so far, only been reported in a single study, where it was found on osteoblast-like cell lines (MG-63 and TE-85) derived from osteosarcomas displaying features of immature osteoblasts.⁽¹⁹⁾ Whether the GLP-2R is expressed on mature human osteoblasts or in osteoclasts is therefore an open question and the mechanism behind the effect of GLP-2 on bone is unexplained. The modulation of bone turnover in response to exogenous injection of GIP and GLP-2 is accompanied by a sudden and drastic decrease in parathyroid hormone (PTH); a similar decrease is observed after meal ingestion.^(11,12,20–23) The extent to which the effects of GIP or GLP-2 on bone turnover are mediated via a reduction in PTH is unknown.

In this study, we first conducted a meal test to investigate the acute responses to meal intake on the markers of bone turnover (CTX and P1NP), GIP, and GLP-2 in patients with hypoparathyroidism. Second, we gave injections of GIP and GLP-2 to the patients after a fasting period to investigate whether the acute effects of exogenous GIP and GLP-2 on bone turnover were dependent on changes in PTH. We included patients with surgical hypoparathyroidism (due to complete thyroidectomy) with low or undetectable levels of PTH to eliminate PTH-mediated effects on bone turnover. We hypothesized that the effect of GIP and GLP-2 on bone resorption (measured by CTX) would be lost in the patients if it was dependent on an inhibition of PTH.

Furthermore, we examined human parathyroid tissue for expression of GIPR and GLP-2R to evaluate the possibility of direct effects of GIP and GLP-2 on PTH secretion as an explanatory factor for their effects on bone. This was combined with an examination of the expression of GIPR and GLP-2R in *in vitro* cultured human osteoblasts and osteoclasts.

Materials and Methods

Human study participants and procedures

Participants with chronic hypoparathyroidism due to total thyroidectomy were recruited from Hvidovre Hospital, Denmark. Both men and women were eligible for inclusion. Exclusion criteria were diabetes, use of recombinant PTH replacement, anti-osteoporotic medication, or corticosteroid therapy within the last year, pregnancy, prior history of malabsorptive disease (eg, celiac, inflammatory bowel disease, or bariatric surgery), allergy to the interventional substances, or Hgb <7 mmol/L. Participants were studied during a meal test, and thereafter they were asked to participate in peptide test days. There was at least 1 week

between the meal test and the peptide tests. The peptide test consisted of three separate test days (in a randomized crossover design) with subcutaneous injections of exogenous GIP, GLP-2, or placebo, respectively (no meals were served during the peptide test days) (Fig. 2A). Apart from the applied stimuli, the procedures for the test days were similar. Experiments were conducted at Hvidovre Hospital, Denmark. Participants refrained from strenuous physical activities and ingestion of alcohol for 2 days before each study day. On study days, participants arrived at 8:00 a.m. after fasting at least 10 hours. Medication throughout the fasting period and during the test day was avoided if possible. Participants were resting in a hospital bed during the test day with a catheter inserted in a cubital vein for blood sampling. Blood samples were drawn at time points –10, –5, 7, 15, 30, 45, 60, 90, 120, 150, 180, 210, and 240 minutes relative to the ingestion of the Nutridrink (meal test) or the subcutaneous injection (peptide test).

Meal test

A 200 mL liquid mixed meal (Nutridrink, N.V. Nutricia, Zoetermeer, The Netherlands) containing 1260 kJ, carbohydrate 49 E%, protein 16 E%, fat 35 E%, and 91 mg calcium/100 mL was ingested at approximately 8:30 a.m. ($t = 0$ minutes).

Peptide test

We conducted a cross-over study with a balanced design (a Williams design). A list of the sequences was generated by a computer and participants were randomly assigned to the sequences. All participants were studied on the three study days: GIP, GLP-2, and placebo, which were separated by at least 1 week. Participants were blinded to the order of the injections. At approximately 8:30 a.m. ($t = 0$ minutes), GIP 100 μ g, GLP-2 400 μ g, or placebo was subcutaneously injected in the umbilical region.

Synthetic human GIP(1–42) was from PolyPeptide (Strasbourg, France) and synthetic human GLP-2(1–33) was from Bachem (Bubendorf, Switzerland), both with a purity above 97%. Peptides were dissolved in sodium hydrogen carbonate buffer with 0.5% human serum albumin to a final peptide concentration of 100 μ g/mL for GIP and 400 μ g/mL for GLP-2. One-milliliter vials were prepared by The Capital Region Pharmacy (Herlev, Denmark) and stored at -20°C . Placebo injection was 1 mL of saline (NaCl 9 mg/mL).

Blood samples

Blood for plasma preparation was collected in chilled EDTA tubes with added dipeptidyl peptidase-4 inhibitor (valine pyrrolidide) to a final concentration of 0.01 mmol/L and kept on ice. Blood for serum was collected in clot activator tubes and kept at room temperature for 30 minutes for coagulation. Tubes were centrifuged 10 minutes at 1200g at 4°C . Plasma and serum samples were stored at -20°C until measurements.

Measurements

Blood pressure and heart rate were measured using a standard blood pressure monitor (Omron M6, Intelli Sense, Omron Healthcare Europe B.V., Hoofddorp, The Netherlands). Glucose was measured with the glucose oxidase method using a YSI (YSI model 2300D STAT plus analyzer, YSI Inc., Yellow Springs, OH, USA). Serum insulin and C-peptide were measured by sandwich immunoassay (Advia Centaur XP, Siemens, Munich, Germany).

Plasma concentrations of GIP and GLP-2 were measured by in-house radio-immunoassays. Total GIP was measured using an antibody directed toward the C-terminal (code no. 80867), which reacts fully with intact and N-terminally truncated GIP. Intact GIP was measured using an antibody that reacts with the N-terminal of intact GIP (code no. 98171). The standard used was human GIP (Bachem, catalog no. H-5645) and the tracer was ¹²⁵I-labeled human GIP (Perkin Elmer, Waltham, MA, USA; catalog no. Nex402). Intact GLP-2 was measured using an antibody specific for the intact N-terminus of GLP-2 (code no. 92160). The standard was recombinant human GLP-2 and the tracer was ¹²⁵I-labeled rat GLP-2 with an Asp33 to Tyr33 substitution. Sensitivity for all assays was below 5 pmol/L and intra-assay coefficient of variation was below 10%. Serum CTX, P1NP, and intact PTH were measured using the chemiluminescence technology on an IDS-iSYS Multi-Discipline Automated System (Immundiagnostic Systems, Frankfurt am Main, Germany). The reportable range for CTX (code IS-3000) is 0.33 to 6.000 ng/mL and the expected values for women is 0.0034 to 1.037 ng/mL. The reportable range for P1NP (code IS-4000) is 2 to 230 ng/mL with expected values for normal adults 27.7 to 127.6 ng/mL. The reportable range for PTH (code IS-3600) is 5 to 5000 pg/mL with expected values for normal adults 11.5 to 78.4 pg/mL.

Expression of the GIP and GLP-2 receptor in parathyroid tissue

The data regarding the relative expression of GLP-1R, GIPR, and GLP-2R among normal human parathyroid glands were derived from our previous comparative transcriptome analysis (PMID: 27760455, 21393447), deposited at Gene Expression Omnibus (GEO accession: GSE83421).

Gene expression analyses in osteoblast and osteoclast cultures

To determine the expression of GIPR and GLP-2R in osteoblasts, a well-characterized human bone marrow stromal (mesenchymal) stem cell line (hMSC-TERT) was employed to obtain osteoblast cultures.^(24,25) hMSC were cultured in a standard growth medium containing minimal essential medium (MEM) (Gibco Invitrogen, Carlsbad, CA, USA) supplemented with 10% FBS (Biocrom, Cambridge, UK) and 1% penicillin/streptomycin (Gibco Invitrogen) at 37°C in a humidified atmosphere containing 5% CO₂. The medium was changed every third day until the cells became 90% confluent. For osteoblast differentiation, cells were seeded at 20,000 cells/cm² and the day after, osteoblast differentiation was induced by supplementing the hMSC culture media with 10 mM β-glycerophosphate (Calbiochem-Merck, Darmstadt, Germany), 50 μg/mL L-ascorbic acid-2-phosphate (Wako Chemicals GmbH, Neuss, Germany), 10 nM dexamethasone (Sigma-Aldrich, Soborg, Denmark), and 10 nM calcitriol (1.25-dihydroxy vitamin-D₃ kindly provided by Leo Pharma, Ballerup, Denmark).⁽²⁶⁾ Gene expression analyses were done using Affymetrix (Santa Clara, CA, USA) Human GeneChip U133 Plus 2.0 arrays in hMSC cultures at the baseline (day 0) and 6 and 12 days after induction of osteoblast differentiation. To determine the expression of GIPR and GLP-2R in osteoclasts, a publicly available data set (GSE107295) was employed, in which gene expression analysis was done using Affymetrix Human GeneChip U133 Plus 2.0 Array in osteoclast cultures established by CSF 1 (33 ng/mL) and RANKL (66 ng/mL) treatment of osteoclast precursors.⁽²⁷⁾ We also determined expression of GIPR and GLP-2R mRNA in

mature osteoclasts, where human CD14⁺ monocytes were isolated from buffy coats obtained from blood donations from healthy women aged 18 to 49 years (*n* = 8). Monocytes were seeded in T25 cell culture flasks (1.67 × 10⁶ cells/flask) stored in an incubator (37°C, 5% CO₂) and differentiated into mature osteoclasts (*t* = 10 days) using minimal essential medium α (Thermo Fisher Scientific, Waltham, MA, USA) with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (Sigma-Aldrich, Seelze, Germany) and stimulated with macrophage colony-stimulating factor (MCSF) (R&D Systems, Minneapolis, MN, USA), 25 ng/mL. Media that contained MCSF and receptor activator of NF-κB ligand (RANKL) (R&D Systems), 25 ng/mL each, were changed on days 3, 6, and 8.^(28,29)

Analyses of gene expression by RNA-sequencing

Primary mature human osteoclasts were harvested in TRIZOL (Thermo Fisher Scientific). RNA was purified using Econo Spin columns (Epoch Life Science, Missouri City, TX, USA). RNA-sequencing was performed according to manufacturer's instructions (TruSeq 2, Illumina, San Diego, CA, USA) using 2 μg RNA for preparation of cDNA libraries. Sequencing reads were mapped to the human genome (hg19) using STAR,⁽³⁰⁾ and tag counts were summarized at the gene level using HOMER.⁽³¹⁾ Differential gene expression was analyzed using DESeq2.⁽³²⁾ Gene ontology analysis was performed using goseq.⁽³³⁾

Statistics

Primary outcome of the human study was bone resorption measured as CTX. CTX and P1NP levels were presented as percentage of baseline (mean of *t* = -10 minutes and *t* = -5 minutes). The changes after peptide administrations were analyzed using a related-samples Friedman's two-way analysis of variance by ranks (non-parametric test) and if relevant, followed by post hoc pairwise comparison between the groups to evaluate differences at specific time points. Differences in baseline values between test days were analyzed by Friedman test (non-parametric test). Participants' baseline characteristics are presented as median and range. Results are presented as mean ± SEM. When measurements were below the detection limit, the detection limit was entered. Two-sided *p* values <0.05 were considered statistically significant. All statistical analyses and graphs were made in SPSS Version 26 for Mac (IBM, Armonk, NY, USA) and GraphPad Prism Version 7.00 for Windows (GraphPad Software, La Jolla, CA, USA). The sample size was determined based on previous studies.^(2,6,9,12)

Study approval

The study was conducted according to Declaration of Helsinki principles. The Scientific Ethical Committee of the Capital Region of Denmark approved the study (protocol no. H-16047626), and it was registered at the Danish Data Protection Agency (journal no. SUND-2017-21) and at ClinicalTrials.gov (NCT03728959). All participants gave written informed consent before inclusion in the study.

Results

Seven female participants were included in the meal test, and 4 of these also completed the subsequent peptide test, where GIP, GLP-2, or placebo, respectively, was administered

subcutaneously on separate study days. Patients were recruited and tested from August 2019 to February 2020. Only women volunteered to participate. Blood samples were drawn in the fasting state and regularly for 240 minutes after the intervention (Fig. 2A). Baseline characteristics of participants are listed in Table 1. In the meal test, 5 of the 7 participants were postmenopausal. In the peptide test, 3 of the 4 participants were postmenopausal. Four participants were not able to skip all their medication and took necessary medication (judged by their physician and the participant) the night before the test day and/or in the morning on the test day. For further information regarding medication, see Table 2 and Supplemental Table S1.

During the meal test, a normal postprandial response with lowering of bone resorption was found in patients with hypoparathyroidism

Participants in the meal test had a median age of 54 (range 40 to 75) years and a median body mass index (BMI) of 23.6 (range 19.8 to 30.1) kg/m². All participants had low basal plasma levels of intact PTH with a median of 10 ± 3.8 (range <5 to 24.7) pg/mL (expected values for normal adults are 11.5 to 78.4 pg/mL). The clinical characteristics of participants are shown in Table 1.

Table 1. Baseline Characteristics of Study Participants in the Meal Test and the Peptide Test

	Meal test	Peptide test
Sex (female/male)	7/0	4/0
Postmenopausal/ premenopausal	5/2	3/1
Age (years)	54 (40–75)	58 (40–75)
Weight (kg)	71 (58–82)	71 (60–82)
Height (m)	1.70 (1.60–1.82)	1.65 (1.60–1.74)
BMI (kg/m ²)	23.6 (19.8–30.1)	27.7 (19.8–30.1)
PTH intact (pg/mL)	10.0 ± 3.80 (<5 to 24.7)	12.3 (<5–21.7)
Calcium total (mmol/L)	2.25 (2.18–2.38)	2.22 (2.22–2.27)
Calcium ion (mmol/L)	1.21 (1.08–2.21)	1.21 (1.08–2.21)
eGFR (mL/min/1.73 m ²)	77 (44–95)	75 (44–95)
Phosphate (mmol/L)	1.28 (0.95–1.48)	1.32 (0.95–1.48)
Potassium (mmol/L)	3.9 (3.6–4.8)	3.9 (3.8–4.8)
Creatinine (μmol/L)	71 (63–115)	68 (65–115)
Alkaline phosphatase (U/L)	65 (34–102)	60 (48–66)
25OH vitamin D (nmol/L)	130 (69–172)	113 (69–149)

BMI = body mass index; PTH = parathyroid hormone; eGFR = estimated glomerular filtration rate.

Baseline characteristics at screening of participants in the meal test (*n* = 7) and the peptide test (*n* = 4) given as median and range.

Table 2. Medication

	Medication	Test
Subject 2	Levothyroxine, alfacalcidol, calcium	Meal test + peptide test
Subject 4	Alfacalcidol, oxycodone, enalapril	Meal test + peptide test
Subject 6	Alfacalcidol, pregabalin, metoprolol succinate, rivaroxaban	Meal test
Subject 7	Metoprolol succinate	Meal test

Four of the 7 participants had to take some of their medication the night before the test day and/or in the morning on the test day.

Plasma concentrations of total GIP and intact GLP-2 increased after the meal as expected. GIP plasma level increased from a basal level 17 ± 3.5 pmol/L (mean ± SEM) to a maximum of 69 ± 10 pmol/L reached at *t* = 45 minutes and then decreased toward the basal level toward the end of the test (Fig. 1A). GLP-2 plasma level increased from 18 ± 2.5 pmol/L to a maximum of 30 ± 4.3 pmol/L reached at *t* = 15 minutes and returned to the baseline level within 1 hour (Fig. 1B).

Insulin and C-peptide increased after the meal with a peak value around *t* = 60 minutes (~250 pmol/L insulin and ~1500 pmol/L C-peptide) followed by a decrease to the baseline level (Fig. 1C, D). Glucose rose modestly from a basal concentration of 5.0 ± 0.1 mmol/L to a peak concentration of 6.2 ± 0.5 mmol/L at *t* = 60 minutes and then decreased to 4.6 ± 1.8 mmol/L (Fig. 1E).

Changes in blood pressure and heart rate were also observed. A decrease in systolic blood pressure from 143 ± 6.7 to 126 ± 6.1 mmHg and diastolic blood pressure from 85 ± 4.4 to 75 ± 6.8 mm Hg was observed after the meal, while the heart rate increased from 66 ± 4.6 to 71 ± 4.8 bpm (Table 3).

After the meal, a small and insignificant decrease in PTH to 7.5 ± 2.9 pg/mL was observed (Fig. 1F). Three of the participants had basal PTH levels below the detection limit of the assay (below 5.0 pg/mL).

Regarding the biochemical markers of bone resorption (CTX) and bone formation (P1NP), all participants responded to the meal with a marked decrease in CTX to a mean ± SEM of 55.5 ± 17.1% of baseline reached at *t* = 120 minutes and then returned toward the baseline level (Fig. 1G). Absolute basal value of CTX was 0.45 ± 0.15 ng/mL. P1NP increased to a maximum of 108 ± 8.9% of baseline reached at *t* = 30 minutes and hereafter P1NP stayed near the baseline level (Fig. 1H). Absolute basal concentration of P1NP was 60.9 ± 13.7 pg/mL.

In patients with hypoparathyroidism, exogenous GIP inhibited bone resorption, whereas the effect of GLP-2 appeared to be lost

Four participants from the meal test also volunteered to participate in the peptide test. Their median age was 58 (range 40 to 75) years and the median BMI was 27.7 (range 19.8 to 30.1) kg/m². For further characteristics of participants, see Tables 1 and 2.

Timeline for the study days is depicted in Fig. 2A. CTX decreased after GIP injection but not after injection of GLP-2. At baseline, CTX levels were comparable with absolute concentrations of 0.30 ± 0.10, 0.37 ± 0.06, and 0.32 ± 0.09 ng/mL in the GIP, GLP-2, and placebo group, respectively. After placebo, CTX stayed near the baseline level during the 4-hour study period. GIP injection significantly decreased CTX to 80.5 ± 5.8% of baseline after 90 minutes compared with 104.2 ± 5.5% of baseline after placebo (*p* = 0.034). GLP-2 did not significantly




Fig 1. Results from the meal test. Seven female participants with hypoparathyroidism were included in the meal test. After at least 10 hours of fasting, the participants met in the morning. After baseline blood samples, a liquid meal was ingested at time point $t = 0$ minute followed by blood sampling regularly for 240 minutes. (A) Plasma concentrations of total glucose-dependent insulinotropic polypeptide (GIP), (B) plasma concentrations of intact glucagon-like peptide-2 (GLP-2), (C) serum insulin, (D) serum C-peptide, (E) serum glucose, (F) serum parathyroid hormone (PTH), (G) serum collagen type 1 C-terminal telopeptide (CTX) presented as percentage of baseline, and (H) serum procollagen type 1 N-terminal propeptide (P1NP) presented as percentage of baseline. Data ($n = 7$) are shown as mean \pm SEM.

Table 3. Insulin, C-Peptide, Glucose, Blood Pressure, and Heart Rate at the Day of the Meal Test and the Peptide Test

	Meal test (n = 7)	Peptide test (n = 4)		
		GIP	GLP-2	Placebo
Insulin (pmol/L)				
Baseline	47 ± 6.3	53 ± 5.7	37 ± 3.3	32 ± 2.7
Peak	262 ± 82	77 ± 11	44 ± 6.8	32 ± 3.7
C-peptide (pmol/L)				
Baseline	501 ± 43	530 ± 72	451 ± 41	399 ± 29
Peak	1536 ± 325	639 ± 80	447 ± 64	404 ± 48
Glucose (mmol/L)				
Baseline	5.0 ± 0.1	5.2 ± 0.1	5.0 ± 0.1	5.0 ± 0.0
Peak/nadir	6.2 ± 0.5 peak	4.6 ± 0.3 nadir	4.9 ± 0.2 nadir	4.7 ± 0.1 nadir
Systolic blood pressure (mmHg)				
Baseline	143 ± 6.7	135 ± 9.1	130 ± 8.3	133 ± 5.9
Nadir	127 ± 6.1	121 ± 9.2	118 ± 12	128 ± 12
Diastolic blood pressure (mmHg)				
Baseline	85 ± 4.4	77 ± 4.3	70 ± 1.1	73 ± 2.6
Nadir	75 ± 6.8	75 ± 8.5	67 ± 6.6	68 ± 3.8
Heart rate (bpm)				
Baseline	66 ± 4.6	67 ± 2.7	66 ± 3.1	62 ± 2.2
Peak	71 ± 4.8	71 ± 3.5	70 ± 3.2	63 ± 4.4
CTX (ng/mL)				
Baseline	0.45 ± 0.15	0.30 ± 0.10	0.37 ± 0.06	0.32 ± 0.09
Nadir	0.21 ± 0.05	0.25 ± 0.16	0.27 ± 0.20	0.31 ± 0.19
P1NP (pg/mL)				
Baseline	60.9 ± 13.7	37.3 ± 12.2	38.2 ± 13.5	39.7 ± 12.2
Peak	65.8 ± 15.4	44.6 ± 21.7	38.9 ± 14.7	44.5 ± 17.0
Nadir	59.5 ± 14.0	26.2 ± 18.1	35.4 ± 10.6	38.9 ± 10.7

Insulin, C-peptide, glucose, blood pressure, and heart rate at the day of the liquid meal test and the peptide test. Data are presented as mean ± SEM baseline and peak/nadir values as relevant.

change CTX compared with placebo (except for a small decrease at t = 210 minutes) (Fig. 2B). There were no effects of GIP or GLP-2 on bone formation as measured by serum levels of P1NP (Fig. 2C), and basal values were comparable on the three study days (37.3 ± 12.2, 38.2 ± 13.4, and 39.7 ± 12.2 ng/mL on the GIP, GLP-2, and placebo day, respectively).

Baseline PTH concentrations were low and below the detection limit for 2 of the participants in the peptide test (<5, <5, 17.9, and 21.7 pg/mL, respectively, for the 4 participants). For the 2 participants with measurable PTH levels, PTH initially decreased slightly after all three interventions (GIP, GLP-2, and placebo) with no differences between the interventions (Fig. 2D).

Plasma concentrations of GIP and GLP-2 increased after the injections, as expected. After GIP administration, total GIP increased to a peak concentration of 163 ± 13 pmol/L reached at t = 60 minutes. Intact GIP increased to a peak of 81 ± 12 pmol/L at t = 15 minutes and dropped to baseline levels at t = 120 minutes. After administration of GLP-2, intact GLP-2 increased to a maximum of 2004 ± 609 pmol/L at t = 45 minutes and concentrations remained elevated until t = 240 minutes (Fig. 2E, F).

Insulin, C-peptide, and glucose were all affected by GIP but not by GLP-2. Compared with placebo, GIP significantly increased levels of insulin and C-peptide during the first 30 minutes after injection. Insulin increased from 53 ± 5.7 to 77 ± 11 pmol/L and C-peptide from 530 ± 72 to 639 ± 80 pmol/L followed by a decrease in glucose from 5.2 ± 0.1 to

4.6 ± 0.3 mmol/L. GLP-2 had no significant effect on insulin, C-peptide, or glucose levels compared with placebo (Table 3).

Regarding blood pressure and heart rate, both GIP and GLP-2 insignificantly decreased systolic blood pressure compared with placebo. There were no changes in diastolic blood pressure. Compared with placebo, the heart rate increased significantly from 67 ± 2.7 to 71 ± 3.5 bpm after GIP and from 66 ± 3.1 to 70 ± 3.2 bpm after GLP-2, similar to the changes observed in the meal test (Table 3).

Receptor expression in human parathyroid tissue

Analysis of our previously published transcriptome array data (PMID: 27760455, 21393447) showed that GLP-1R, GIPR, and GLP-2R are expressed in normal human parathyroid glands. Moreover, the relative expression of GLP-1R, GIPR, and GLP-2R revealed comparable expression levels (Fig. 3).

Receptor expression in human osteoblasts and osteoclasts

To determine the expression of GIPR and GLP-2R in osteoblasts, cultured human bone marrow stromal (mesenchymal) stem cell line (hMSC-TERT) was induced toward osteoblast differentiation, and gene expression analysis was performed at different stages of differentiation (days 0, 6, and 12). The analysis showed expression of GIPR at all time points, whereas GLP-2R expression was not detectable at any time point (Fig. 4A). A publicly available gene expression data set⁽²⁷⁾ was employed to determine the




Fig 2. Results from the peptide test. (A) Timeline for the study. Baseline blood samples were drawn from the fasting participants in the morning (after at least 10 hours of fasting) at time points $t = -10$ and -5 minutes. At $t = 0$ minute, glucose-dependent insulinotropic polypeptide (GIP), glucagon-like peptide-2 (GLP-2), or placebo (saline) was subcutaneously injected followed by blood sampling at $t = 7, 15, 30, 45, 60, 90, 120, 150, 180, 210,$ and 240 minutes. (B) Serum collagen type 1 C-terminal telopeptide (CTX) presented as percentage of baseline, (C) serum procollagen type 1 N-terminal propeptide (P1NP) presented as percentage of baseline, and (D) serum parathyroid hormone (PTH) in response to GIP (green circles), GLP-2 (blue squares), and placebo (gray triangles). (E) Plasma concentration of total GIP (white circles) and intact GIP (black circles) and (F) plasma concentrations of intact GLP-2 (black triangles). (B–D) The data were analyzed by a related-samples Friedman’s two-way analysis of variance by ranks, and if relevant, followed by post hoc pairwise comparisons between groups to evaluate differences between interventions (GIP or GLP-2) and placebo at individual time points. Actual p values (when $p < 0.10$ or $p > 0.001$) are indicated on the figure. Data ($n = 4$) are shown as mean \pm SEM.

expression of GIPR and GLP-2R in osteoclasts. Using this method, we found an increasing expression of GIPR during osteoclast differentiation, whereas GLP-2R was not detectable (Fig. 4B). We also assessed expressions of the GIPR and GLP-2R in osteoclasts differentiated from human CD14⁺ monocytes isolated from

buffy coats obtained from anonymous blood donors ($n = 8$). Gene expression analyses using RNA sequencing were performed on day 10, ie, in mature osteoclasts. These analyses confirmed expression of GIPR in primary mature human osteoclasts. However, contradictory to the publicly available gene expression

data set, we observed expression of GLP-2R in primary mature osteoclast in vitro in all donors. However, the expression of the GLP-2R was substantially weaker compared with that of the GIPR (Fig. 4C).

Discussion

The present study provides new insight into the gut–bone axis and demonstrates basic aspects of bone turnover in response to meal ingestion and subcutaneous injection of GIP and GLP-2. We conducted a liquid mixed meal test and injected




Fig 3. Receptor expression in normal parathyroid glands. Glucagon-like peptide-1 receptor (GLP-1R), glucose-dependent insulinotropic polypeptide receptor (GIPR), and glucagon-like peptide-2 receptor (GLP-2R) are expressed in normal human parathyroid tissue. Analysis of data from GEO series accession GSE83421 shows the relative expression of the GLP-1R, GIPR, and GLP-2R in normal parathyroid tissue. Grubbs' test identified a gland as an outlier in GLP-2R expression and data were excluded from further statistical analysis by one-way ANOVA. Data are shown as boxplot (with median, interquartile range, maximum, and minimum) showing all data points. No significant differences were found among presented genes.

exogenous GIP, GLP-2, and placebo to investigate the effects on bone turnover in patients with hypoparathyroidism (due to thyroidectomy) with no or very low levels of PTH. We did this to assess to what extent the changes in bone turnover markers (CTX and P1NP), as observed in healthy individuals,^(2–6,9–12) depend on inhibition of PTH secretion, which is normally acutely lowered after meal ingestion as well as after injections of both GIP and GLP-2.^(12,23,34) If PTH inhibition was essential, the changes in bone turnover markers would not be expected to occur in the patients with hypoparathyroidism.

During the mixed meal, total GIP and intact GLP-2 plasma concentrations increased to ~ 70 pmol/L and 30 pmol/L, respectively, which is similar to what is found in healthy persons.^(2,23,35) Concomitantly, CTX levels decreased to $55.5 \pm 17.1\%$ of baseline in the patients with hypoparathyroidism, similar to the meal-induced decrease to 50% to 60% of baseline observed in healthy individuals.^(1,2,7) The mixed meal also resulted in a transient increase in P1NP to $108 \pm 3.7\%$ of baseline. Regarding the effect of a meal or glucose intake on bone formation, the literature is inconsistent. Whereas some studies report decreases in markers of bone formation, others report minor increases.^(1,11,36,37) However, some studies have shown that GIP may induce a small acute increase in P1NP.^(12,15,34) Regarding PTH, we confirmed that the participants had low levels of PTH and 3 of the 7 participants in the meal test even had PTH levels below the measurable concentration (<5 pg/mL). Importantly, all participants responded during the meal test with a marked decrease in CTX, irrespective of their basal PTH levels, suggesting that the changes in bone turnover found after meal ingestion are probably not related to changes in PTH levels. During the meal test, insulin, C-peptide, and glucose increased as normally found in healthy subjects.^(35,38,39)

With the peptide injections, we studied the individual effects of exogenously administered GIP and GLP-2 in our patients with hypoparathyroidism. After GIP, we observed a clear reduction in CTX to $80.5 \pm 5.8\%$ of baseline at $t = 90$ minutes compared with a value of $104.2 \pm 5.5\%$ after placebo ($p = 0.034$). The reduction in CTX is less than previously observed after subcutaneous injection




Fig 4. Receptor expression in osteoblasts and osteoclasts. Expression of glucose-dependent insulinotropic polypeptide receptor (GIPR) and glucagon-like peptide-2 receptor (GLP-2R) in human osteoblast and osteoclasts. (A) Analyses of GIPR and GLP-2R expression at different time points (days 0, 6, 12) during osteoblast differentiation of human mesenchymal stem cells (hMSC) and (B) analyses of GIPR and GLP-2R expression at different time points (36 hours, 72 hours, 96 hours) during osteoclast differentiation of human osteoclast precursors treated with macrophage-colony stimulating factor MCSF-1 and receptor activator of NF- κ B ligand (RANKL). Actin beta (ACTB), lipoprotein lipase (LPL), alkaline phosphatase (ALPL), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), bone gamma-carboxyglutamate protein (BGLAP), and cathepsin K (CTSK) were used as housekeeping, negative, and positive controls, respectively. (C) mRNA expression of GLP-2R, GIPR, and cathepsin K (CTSK) (positive control) in primary mature human osteoclasts treated with MCSF and RANKL ($n = 8$). Each gene is normalized to gene length and the number of reads sequenced per sample (reads per kilobase million [RPKM]). (C) Data are shown as boxplot (with median, interquartile range, maximum, and minimum) showing all data points.

tion of GIP in healthy individuals, where a maximal inhibition to ~55% of baseline was observed after 90 minutes.⁽¹²⁾ However, it should be noted that in the previous study, a higher dose of GIP was used (200 µg versus 100 µg in the present study) and also a greater reduction in CTX after placebo was observed.⁽¹²⁾ In the present study, P1NP increased to 116% ± 15% of baseline after GIP administration, but this change did not reach significance. In healthy individuals and in patients with type 1 diabetes, P1NP has been found to significantly increase after GIP,^(11,12,15) indicating an acute uncoupling of bone formation from bone resorption. However, the present study was undoubtedly underpowered to detect a significant difference in P1NP since only 4 participants received the injections and the P1NP response is substantially smaller than the CTX response.⁽¹²⁾

After subcutaneous injection of GLP-2, CTX levels remained unchanged and similar to those observed with placebo, suggesting that the effect of GLP-2 is diminished in hypoparathyroidism and may depend on changes in PTH levels. In comparison, a significant reduction to ~60% of baseline CTX is found after GLP-2 in healthy young men.⁽¹²⁾ We have previously observed basal levels of CTX in the range of ~0.7 to 0.9 ng/mL and basal levels of P1NP of ~70 ng/mL in healthy young men.⁽¹²⁾ In the present study, the participants had basal levels of CTX ranging from 0.3 to 0.45 ng/mL and P1NP basal levels were 38 to 60 pg/mL, which was as expected because patients with hypoparathyroidism have low levels of bone turnover markers. Hypoparathyroidism is most often caused by accidental removal during thyroid surgery due to either cancer or hyperthyroidism,^(40,41) and the disease is characterized by low PTH, leading to a low bone turnover, low serum calcium, elevated levels of serum phosphorus, and low 1,25(OH)D₃. In most patients with hypoparathyroidism, BMD is increased compared with healthy controls due to the low bone turnover, although the microarchitecture of the bone is abnormal.^(40,41) Normally, BMD correlates positively with reduced fracture risk, but lower bone remodeling may result in impaired regeneration and accumulated old bone with fatigue microfractures. Whether the fracture risk is affected in hypoparathyroidism is not clear,^(41,42) but it may be unchanged or lower in some patients.⁽⁴³⁾

For the two participants with measurable PTH levels, we observed a small decrease in PTH after both GIP and GLP-2, but this was also observed after placebo (absolute changes were 2.9, 2.9, and 2.1 pg/mL, respectively, within 45 minutes). In healthy participants, the changes in PTH are more pronounced with more acute decreases around 9 pg/mL (from ~29 to ~20 pg/mL within 15 minutes) after GIP and GLP-2 and with no change after placebo.⁽¹²⁾ Thus, the antiresorptive effect of GIP injection seems to be independent of changes in PTH, whereas the effect of GLP-2 may depend on an inhibitory effect on PTH secretion.

Our finding that GIP decreases CTX in patients with hypoparathyroidism is consistent with our *in vitro* results demonstrating GIPR expression in human osteoblast and osteoclast cultures and support that GIP may act directly on bone cells in a PTH-independent fashion. By contrast, GLP-2R was not expressed in osteoblasts and absent or only weakly expressed in osteoclasts (depending on the method used), in agreement with the indiscernible effect of GLP-2 in patients with hypoparathyroidism. However, we did find expression of GLP-2R (as well as the GIPR and GLP-1R) in human parathyroid tissue, which could explain the effect of GLP-2 on bone as an indirect effect mediated by GLP-2R in the parathyroid gland. However, further studies are now required to investigate the possible direct effect of GLP-2 on the parathyroid glands.

In a previous study, investigating the effect of GLP-2 on bone turnover in patients with short bowel syndrome (SBS), a GLP-2 injection (1600 µg) only reduced PTH and CTX in patients with ileostomy and not in patients with jejunostomy.⁽²⁰⁾ This might suggest that the underlying mechanism whereby GLP-2 reduces CTX may involve reductions in PTH as well as a factor derived from the intestine, eg, a cytokine or a growth factor, or by increased calcium absorption that in turn may change the PTH level. Because the GLP-2R is expressed in enteric neurons,⁽⁴⁴⁾ the effect of GLP-2 may also be mediated through neuronal signaling.

A limitation of the present study is the small sample size and lack of a non-hypoparathyroid control group. However, regarding the GIP response, we achieved statistically significant results regarding CTX, and we know from previous studies in healthy controls (conducted using the same methods) that subcutaneous GLP-2 injections normally result in a response of similar magnitude as that obtained after subcutaneous GIP (with CTX being reduced to around 60% of baseline).^(12,45) Our results may also be confounded by the inclusion of participants using various types of medication. The main strength of the investigation is that it was possible to carry out these studies in patients with hypoparathyroidism and that all participants served as their own controls in a crossover design in the peptide test. Although our results strongly indicate that GIP modulates bone turnover independent of changes in PTH and that the GLP-2 effect appears to depend on changes in PTH, we recognize that the patients with hypoparathyroidism still had some residual PTH secretion and small changes in PTH levels therefore did occur during the tests. However, we assume that these small changes in PTH are biologically insignificant; in addition, the changes in PTH were of similar magnitude after all three interventions in the peptide test (ie, after GIP, GLP-2, and placebo). All patients had undergone total thyroidectomy and therefore also lacked the thyroid gland, which may have affected the calcitonin responses to GIP and GLP-2, if any. Thus, a further limitation is that we did not measure calcitonin, and we also did not measure total or ionized calcium during test days. Further studies are now needed to confirm our findings.

Our findings are compatible with the notion that GIP seems to act directly on the bone cells mediated by the GIPR in a non-PTH-dependent fashion, while the effect of GLP-2 may be mediated in a PTH-dependent fashion, which would explain the diminished GLP-2 effect in the patients with hypoparathyroidism in our study. The apparently different underlying mechanisms of action of GIP and GLP-2 on bone turnover may mean that a combination of the two hormones might have additive or even synergistic effects, which would be of interest from a pharmacological point of view.

Disclosures

JJH and MMR are shareholders of Antag Therapeutics. JJH, MMR, and BH are shareholders of Bainan Biotech. SM has served on advisory boards of AstraZeneca, Boehringer Ingelheim, Eli Lilly, Intarcia Therapeutics, Merck Sharp & Dohme, Novartis, Novo Nordisk, and Sanofi; has received lecture fees from AstraZeneca, Boehringer Ingelheim, Merck Sharp & Dohme, Novo Nordisk, and Sanofi; and is a research grant recipient from Novo Nordisk and Boehringer Ingelheim. JBJ has served on advisory boards of Amgen, Eli Lilly, UCB, and Gedion Richter, and has received lecture fees from Amgen, Eli Lilly, UCB, Gilead, and Otsuka. NH has

received research funding from Alexion Pharmaceuticals, Inc. All other authors have declared no conflicts of interest.

Acknowledgments

This work was supported by AP Møller Fonden, Novo Nordisk Foundation, and in part by NIH grant (1R01DK088188-01) and Novo Nordisk Foundation Center for Basic Metabolic Research. Novo Nordisk Foundation Center for Basic Metabolic Research is an independent research center, based at the University of Copenhagen, Denmark, and partially funded by an unconditional donation from the Novo Nordisk Foundation (www.cbmr.ku.dk) (grant no. NNF18CC0034900).

Authors' roles: KS, NH, SM, JBJ, JJH, MMR, and BH conceptualized and designed the human study. KS and NH conducted the human study. KS and BH measured bone markers, GIP, and GLP-2 levels in samples from the human study. JO and AJ conducted the *in vitro* study in osteoblasts and did the determination of the GIPR and GLP-2R expression in osteoclasts using the publicly available data set. MSH and MF conducted the *in vitro* study in mature human osteoclasts with measurements of mRNA expression of the GIPR and GLP-2R. NB and JAO provided the data regarding the GIPR and GLP-2R expression in parathyroid tissue. All contributed to the interpretation of data. KS drafted the manuscript, and all other authors critically revised the manuscript for important intellectual content. All authors have approved the final version of the manuscript. BH is responsible for the integrity of the work as a whole.

Author contributions: Kirska Skov-Jeppesen: Conceptualization; formal analysis; investigation; methodology; project administration; writing-original draft; writing-review & editing. Nicola Hepp: Conceptualization; investigation; methodology; resources; writing-review & editing. Jannika Oeke: Formal analysis; investigation; writing-review & editing. Morten Hansen: Formal analysis; investigation; methodology; writing-review & editing. Abbas Jafari Kerani: Formal analysis; investigation; methodology; writing-review & editing. Maria Svane: Writing-review & editing. Nariman Balenga: Formal analysis; methodology; writing-review & editing. John Olson: Formal analysis; methodology; writing-review & editing. Morten Frost: Formal analysis; investigation; methodology; writing-review & editing. Moustapha Kassem: Writing-review & editing. Sten Madsbad: Conceptualization; methodology; resources; writing-review & editing. Jens-Erik Beck Jensen: Conceptualization; methodology; resources; supervision; writing-review & editing. Jens Holst: Conceptualization; funding acquisition; methodology; resources; supervision; writing-review & editing. Mette Rosenkilde: Conceptualization; funding acquisition; methodology; supervision; writing-review & editing. Bolette Hartmann: Conceptualization; funding acquisition; methodology; project administration; resources; supervision; writing-review & editing.

Peer review

The peer review history for this article is available at <https://publons.com/publon/10.1002/jbmr.4308>.

Data availability statement

Data available on request from the authors.

References

1. Clowes JA, Allen HC, Prentis DM, Eastell R, Blumsohn A. Octreotide abolishes the acute decrease in bone turnover in response to oral glucose. *J Clin Endocrinol Metab.* 2003;88(10):4867-4873.
2. Henriksen DB, Alexandersen P, Bjarnason NH, et al. Role of gastrointestinal hormones in postprandial reduction of bone resorption. *J Bone Miner Res.* 2003;18(12):2180-2189.
3. Henriksen DB, Alexandersen P, Byrjalsen I, et al. Reduction of nocturnal rise in bone resorption by subcutaneous GLP-2. *Bone.* 2004;34(1):140-147.
4. Henriksen DB, Alexandersen P, Hartmann B, et al. Disassociation of bone resorption and formation by GLP-2: a 14-day study in healthy postmenopausal women. *Bone.* 2007;40(3):723-729.
5. Henriksen DB, Alexandersen P, Hartmann B, et al. Four-month treatment with GLP-2 significantly increases hip BMD: a randomized, placebo-controlled, dose-ranging study in postmenopausal women with low BMD. *Bone.* 2009;45(5):833-842.
6. Nissen A, Christensen M, Knop FK, Vilsboll T, Holst JJ, Hartmann B. Glucose-dependent insulinotropic polypeptide inhibits bone resorption in humans. *J Clin Endocrinol Metab.* 2014;99(11):E2325-E2329.
7. Bjarnason NH, Henriksen EE, Alexandersen P, Christgau S, Henriksen DB, Christiansen C. Mechanism of circadian variation in bone resorption. *Bone.* 2002;30(1):307-313.
8. Schiellerup SP, Skov-Jeppesen K, Windelov JA, et al. Gut hormones and their effect on bone metabolism. Potential drug therapies in future osteoporosis treatment. *Front Endocrinol (Lausanne).* 2019;10:75.
9. Askov-Hansen C, Jeppesen PB, Lund P, Hartmann B, Holst JJ, Henriksen DB. Effect of glucagon-like peptide-2 exposure on bone resorption: effectiveness of high concentration versus prolonged exposure. *Regul Pept.* 2013;181:4-8.
10. Gottschalck IB, Jeppesen PB, Holst JJ, Henriksen DB. Reduction in bone resorption by exogenous glucagon-like peptide-2 administration requires an intact gastrointestinal tract. *Scand J Gastroenterol.* 2008;43(8):929-937.
11. Christensen MB, Lund A, Calanna S, et al. Glucose-dependent Insulinotropic polypeptide (GIP) inhibits bone resorption independently of insulin and glycemia. *J Clin Endocrinol Metab.* 2017;103(1):288-294.
12. Skov-Jeppesen K, Svane MS, Martinussen C, et al. GLP-2 and GIP exert separate effects on bone turnover: a randomized, placebo-controlled, crossover study in healthy young men. *Bone.* 2019;125:178-185.
13. Gabe MBN, van der Velden WJC, Gadgaard S, et al. Enhanced agonist residence time, internalization rate and signalling of the GIP receptor variant [E354Q] facilitate receptor desensitization and long-term impairment of the GIP system. *Basic Clin Pharmacol Toxicol.* 2020;126(Suppl 6):122-132.
14. Torekov SS, Harslof T, Rejnmark L, et al. A functional amino acid substitution in the glucose-dependent insulinotropic polypeptide receptor (GIPR) gene is associated with lower bone mineral density and increased fracture risk. *J Clin Endocrinol Metab.* 2014;99(4):E729-E733.
15. Gasbjerg LS, Hartmann B, Christensen MB, et al. GIP's effect on bone metabolism is reduced by the selective GIP receptor antagonist GIP (3-30)NH₂. *Bone.* 2020;130:115079.
16. Hadjidakis DJ, Androulakis II. Bone remodeling. *Ann N Y Acad Sci.* 2006;1092:385-396.
17. Zhong Q, Itokawa T, Sridhar S, et al. Effects of glucose-dependent insulinotropic peptide on osteoclast function. *Am J Physiol Endocrinol Metab.* 2007;292(2):E543-E548.
18. Bollag RJ, Zhong Q, Phillips P, et al. Osteoblast-derived cells express functional glucose-dependent insulinotropic peptide receptors. *Endocrinology.* 2000;141(3):1228-1235.
19. Pacheco-Pantoja EL, Ranganath LR, Gallagher JA, Wilson PJ, Fraser WD. Receptors and effects of gut hormones in three osteoblastic cell lines. *BMC Physiol.* 2011;11:12.
20. Gottschalck IB, Jeppesen PB, Hartmann B, Holst JJ, Henriksen DB. Effects of treatment with glucagon-like peptide-2 on bone resorption

- in colectomized patients with distal ileostomy or jejunostomy and short-bowel syndrome. *Scand J Gastroenterol*. 2008;43(11):1304-1310.
21. Valderas JP, Padilla O, Solari S, Escalona M, Gonzalez G. Feeding and bone turnover in gastric bypass. *J Clin Endocrinol Metab*. 2014;99(2):491-497.
 22. Polymeris AD, Doumouchtsis KK, Giagourta I, Karga H. Effect of an oral glucose load on PTH, 25OHD3, calcium, and phosphorus homeostasis in postmenopausal women. *Endocr Res*. 2011;36(2):45-52.
 23. Fuglsang-Nielsen R, Rakvaag E, Vestergaard P, et al. Consumption of nutrients and insulin resistance suppress markers of bone turnover in subjects with abdominal obesity. *Bone*. 2020;133:115230.
 24. Simonsen JL, Rosada C, Serakinci N, et al. Telomerase expression extends lifespan and prevents senescence-associated impairment of osteoblast functions. *Nat Biotechnol*. 2002;20(6):592-596.
 25. Jafari A, Isa A, Chen L, et al. TAF2A2 induces skeletal (stromal) stem cell migration through activation of Rac1-p38 signaling. *Stem Cells*. 2019;37(3):407-416.
 26. Jafari A, Qanie D, Andersen TL, et al. Legumain regulates differentiation fate of human bone marrow stromal cells and is altered in postmenopausal osteoporosis. *Stem Cell Reports*. 2017;8(2):373-386.
 27. Carey HA, Hildreth BE 3rd, Geisler JA, et al. Enhancer variants reveal a conserved transcription factor network governed by PU.1 during osteoclast differentiation. *Bone Res*. 2018;6:8.
 28. Boissy P, Andersen TL, Abdallah BM, Kassem M, Plesner T, Delaisse JM. Resveratrol inhibits myeloma cell growth, prevents osteoclast formation, and promotes osteoblast differentiation. *Cancer Res*. 2005;65(21):9943-9952.
 29. Soe K, Delaisse JM. Glucocorticoids maintain human osteoclasts in the active mode of their resorption cycle. *J Bone Miner Res*. 2010;25(10):2184-2192.
 30. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29(1):15-21.
 31. Heinz S, Benner C, Spann N, et al. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell*. 2010;38(4):576-589.
 32. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550.
 33. Young MD, Wakefield MJ, Smyth GK, Oshlack A. Gene ontology analysis for RNA-seq accounting for selection bias. *Genome Biol*. 2010;11:R14.
 34. Christensen MB, Lund A, Calanna S, et al. Glucose-dependent insulinotropic polypeptide (GIP) inhibits bone resorption independently of insulin and glycemia. *J Clin Endocrinol Metab*. 2018;103(1):288-294.
 35. Vilsboll T, Krarup T, Sonne J, et al. Incretin secretion in relation to meal size and body weight in healthy subjects and people with type 1 and type 2 diabetes mellitus. *J Clin Endocrinol Metab*. 2003;88(6):2706-2713.
 36. Clowes JA, Robinson RT, Heller SR, Eastell R, Blumsohn A. Acute changes of bone turnover and PTH induced by insulin and glucose. Euglycemic and hypoglycemic hyperinsulinemic clamp studies. *J Clin Endocrinol Metab*. 2002;87(7):3324-3329.
 37. Bergmann NC, Lund A, Gasbjerg LS, et al. Separate and combined effects of GIP and GLP-1 infusions on bone metabolism in overweight men without diabetes. *J Clin Endocrinol Metab*. 2019;104(7):2953-2960.
 38. Gasbjerg LS, Helsted MM, Hartmann B, et al. GIP and GLP-1 receptor antagonism during a meal in healthy individuals. *J Clin Endocrinol Metab*. 2020;105(3):e725-e728.
 39. Alsalm W, Tura A, Pacini G, et al. Mixed meal ingestion diminishes glucose excursion in comparison with glucose ingestion via several adaptive mechanisms in people with and without type 2 diabetes. *Diabetes Obes Metab*. 2016;18(1):24-33.
 40. Rubin MR, Bilezikian JP. Hypoparathyroidism: clinical features, skeletal microstructure and parathyroid hormone replacement. *Arq Bras Endocrinol Metabol*. 2010;54(2):220-226.
 41. Clarke BL. Bone disease in hypoparathyroidism. *Arq Bras Endocrinol Metabol*. 2014;58(5):545-552.
 42. Silva BC, Rubin MR, Cusano NE, Bilezikian JP. Bone imaging in hypoparathyroidism. *Osteoporos Int*. 2017;28(2):463-471.
 43. Underbjerg L, Sikjaer T, Mosekilde L, Rejnmark L. Postsurgical hypoparathyroidism—risk of fractures, psychiatric diseases, cancer, cataract, and infections. *J Bone Miner Res*. 2014;29(11):2504-2510.
 44. Bjerknes M, Cheng H. Modulation of specific intestinal epithelial progenitors by enteric neurons. *Proc Natl Acad Sci U S A*. 2001;98(22):12497-12502.
 45. Gabe MBN, Skov-Jeppeesen K, Gasbjerg LS, et al. P12 Osteoporosis treatment with GIP and GLP-2 dual-agonists based on their synergistic actions in humans—a novel therapeutic principle in children and premenopausal women suffering from bone fragility 20. Presented at: ICCBH VIRTUAL FORUM, Bone Fragility Disorders in Children 18–20; November 2020.