



Genotypic and technological characterization of Lactic acid bacteria from African fermented foods

Adimpong, David Bichala; Nielsen, Dennis Sandris; Sørensen, K.I.; Derkx, P.M.F.; Jespersen, Lene


Published in:
22nd International ICFMH Symposium Food Micro 2010

Publication date:
2010


Document version
Early version, also known as pre-print

Citation for published version (APA):
Adimpong, D. B., Nielsen, D. S., Sørensen, K. I., Derkx, P. M. F., & Jespersen, L. (2010). Genotypic and technological characterization of *Lactic acid bacteria* from African fermented foods. In *22nd International ICFMH Symposium Food Micro 2010* Copenhagen.

22nd International ICFMH Symposium Food Micro 2010



Copenhagen 30th August - 3rd September



Final Programme & Abstract Book



www.foodmicro.dk

Author index

Abay S	PED2.60	Andreasen, AS	PSE1.03	Bahalul M	PEA1.66, PEA1.67	Bester Ingrid	
Abban S	PEA1.70	Andrews S	PEB2.19	Bak Christensen B	PED2.14	Bešvir V	F
Abban, Stephen	PSD1.03	Andritsos Nikolaos	PEC1.01, PEC1.72	Balla C	PED1.19, PED1.20	Beuchat Larry R.	F
Abbasadegan M	PEC2.45	Angelidis Apostolos	PEA1.60, PED2.51	Bang-Berthelsen I	PED1.03	Beumer R	F
Abdelgadir WS	PEA1.70	Anihouvi Victor Bienvenu	PEA1.38	Baranyi, J	PSC1.02, PSE2.04	Bianchi Daniela Man	
Abee T	PEC1.29, PEB2.29, PSD2.02, PSB2.03	Anita F	PEA2.22	Barbaro A	PEB1.29	Bianchi DM	
Abee Tjikko	PEB2.09	Anslow P	PEB2.44	Barber A	PEE2.02	Biavati, B	
Abildgaard L	PEB1.30	Anyogu A	PEA2.18, PEC2.28	Barbosa Matheus de Souza	PED2.48	Bidot, D	
Adam Katharine	PEB2.02	Anyogu Amara	PEB2.39	Barbuti S	PEC1.76	Bigwood, T	
Adebayo CO	PED2.26, PED2.27	Aota T	PEB1.02	Bardowski J.	PEA1.46	Binter C	
Adelantado C	PEB1.25	Aponte M	PEA1.43	Baril Eugenie	PEB2.15	Birk T	
Adelantado Carles	PEE2.19	Aponte Maria	PEA1.75	Barile M	PED2.06	Birk Tina	
Aderiye Jadesola B	PED2.26, PED2.27	Aquilanti Lucia	PEA1.47	Barker G	PEC2.06, PEC2.11, PEC2.37	Bisotti S	F
Adimpong David B	PEA1.40	Arabshahi M	PEB1.27	Barmark G	PEC1.12	Bizzarri F	
Adolphe Ysabelle	PEC1.40, PEC1.41	Aranda Medina E	PEE2.06	Barrucci F	PEC2.40	Bjerrum Lotte	
Aertsen A	PEB2.66, PSB2.06	Arapaki S	PEB2.50	Barrucci Federica	PEC2.39	Björkroth KJ	
Aguilhon C	PEC1.22	Arbault Patrice	PEC1.80, PEB2.55	Bartolomeoli I	PEC1.34, PEB2.62, PEB2.63	Björkroth KJ	
Akabanda F	PEA1.36	Arcangeli G	PEC1.35	Basaglia M	PEA2.08	Blaiotta G	
Akabanda Fortune	PEA1.37	Arendt EK	PEA2.27	Bassami MR	PEC2.45	Blaiotta Giuseppe	I
Akbarien H	PEB1.27	Argyri Anthoula	PEC1.55	Bassi D	PEC2.53	Blana V	I
Akissoe NH	PEA1.14	Argyri, A	PSD2.06	Bassi Daniela	PEA2.37	Blank L	
Alapont C	PEC2.19	Arinder P	PEC1.48	Batista Almeida e Silva J	PEA1.26	Bleicher A	
Alastrure A	PEC2.19	Arinder Pernilla	PED2.29	Battilani A	PEC2.54	Blessington T	
Albert I	PEC2.49	Arneborg N	PEA1.41, PED2.16, PED2.39	Battilani A	PEC2.54	Blohmke C	
Alegria A	PEA1.46	Arneborg Nils	PEA1.10	Bavan T	PEA1.19	Bodrossy L	
Alessandria V	PEA1.45, PEA1.65, PEC1.71, PSB1.05	Arosemena EL	PEB1.25	Baylis C	PED1.30	Body C	
Alessio G	PEA1.34	Arosemena EL	PEE2.19	Beal Catherine	PEA1.03	Boel Jeppe	
Alexandre, H	PSA2.01	Asakura M	PEC1.77, PEA2.38	Beck Hansen T	PED2.40	Bogicevic, Biljana	
Alftren J	PEE2.02	Atamer Zeynep	PEA1.16	Begum M	PEC1.56	Bolaños S	
Ali Y	PEA1.16	Atindehou MM	PEB1.24	Behera P	PEB1.17	Bolton Declan	
Allende A	PED2.42, PSD1.04	Aubert, X	PSD2.04	Belessi Charalambia-Irini	PEB2.50, PEC2.46	Bonetta S	I
Almeida E	PEA1.39, PEA1.58, PEA1.76	Audenaert W TM	PED1.25	Belfiore C	PEB2.42	Bonetta Silvia	I
Almeida MV	PEC1.13	Auvinen P	PEA2.28	Belk KE	PED1.27	Bonke R	
Alonso-Calleja C	PEB2.51	Avendano-Perez, Gaspar	PSE2.04	Belletti N	PEC1.32, PEC1.93, PED2.58	Bonnichsen L	
Alsady M	PEE2.56	Axelsson L	PEB2.52	Bello OO	PEA2.34	Borch E	PEC1.48,
Al-Soud WA	PEE2.14, PSE1.03	Ayaz ND	PEB1.01	Beltran, D	PSD1.04	Bord C	
Álvarez JC	PED1.05	Aydin F	PED2.60	Beny L	PED2.34	Bordignon Stefano	
Álvarez-Ordóñez Avelino	PEB2.51, PEB2.52	Ayernor GS	PEA1.38	Beny Laurent	PED2.35	Bordin P	
Alves de Ávila AR	PED2.61	Aymerich T	PEA1.35, PEC1.32, PEC1.45, PED1.18, PED2.33	Benito Bernáldez MJ	PEA1.13	Bordoni, A	
Alvseike O.	PEB2.52	Aymerich Teresa	PEB2.40	Bensoussan M	PEC1.73, PEC1.74, PEC1.75	Borilova G	
Ambrožič M	PEC2.32	Azadnia Parisa	PEA1.02	Berkvens D	PEC1.30	Borkovcova I	
Amrouche, Tahar	PSA2.06	Azevedo A	PEC2.22	Bermúdez E	PEA2.38	Boseret G	
An BK	PEB2.10	Azimi-rad M	PEB1.27	Bernard M	PEC2.62	Botsaris George	
Anastasio A	PED2.06	Azokpota Paulin	PEA1.42	Bernardi T	PED2.47	Bottari B	
Anastasiou R	PEA1.11	Baba-Moussa Lamine	PEB1.24	Berndtsson B	PEB1.06	Bottari Benedetta	
Ancelet-Enjalric Sophie	PEC2.49	Badirou EM	PEB1.24	Bernini V	PEA2.09, PEE2.09	Botteldoorn N	I
Andersson G	PEC1.07, PEC2.37	Baert L	PEC2.48	Bernini Valentina	PED2.18	Bouix M	
Andjelkovic M	PEC1.102, PSB1.04	Baert Leen	PEC2.15, PSC2.04	Berstad A	PEC2.08	Boumgha-Bourtcha	
Andrade MJ	PEA2.38	Baffone W	PEC1.101	Berthoud H	PEA1.19	Bourke P	
Andre Stéphane	PEB2.12	Baffoni, Loredana	PSE1.04	Berthoud, H	PSA1.03, PSC2.06	Bouvier M	
		Baggesen DL	PEC1.100, PED1.23, PSE1.02	Bessell, P		Bouw E	
						Bouwknegt M	
						Bover-Cid S	I

Nielsen (3), L Jespersen (3)

another depending on the used, the processing conditions certain tribes of the West and so little or no scientific knowledge of the microorganisms present. The survey revealed that traditional calabashes or plastic containers in the environment, processing and fermentation was followed to the end of the fermentation process from 4.00 to 9.00 log CFU/g diversity was evaluated using DGGE. The predominant lactic acid bacteria are *Lactobacillus helveticus*, *Lactobacillus delbrueckii*, *Weissella cibaria* and other species are involved in natural and controlled fermentations.

Production by starter cul-

(4)

of *Lanhouin* were tested to determine the cell pattern was enumeration and the role of individual components by GC-MS. For all fermentations, the cell count was 3.7×10^8 cfu/g after 48 h of fermentation respectively. The fermentation progressed. The histogram of 41 aroma compounds were identified. These compounds included alcohols, 8 amines, 3 aldehydes, 3 phenols, thiazoles and pyrroles.

PEA1.39 Bacterial community present in a fermented beverage produced from cotton seed with rice by Brazilian Amerindians

Cintia Ramos (1), E Almeida (1), AL Freire (1), D Dias (2), R Schwan (1)
(1) UFLA, Brazil
(2) UNILAVRAS, Brazil

Cauim is a fermented beverage prepared by Tapirapé Amerindians in Brazil from substrates such as cassava, rice, peanuts, pumpkin, banana, cotton seed and maize. The process is traditional and rudimentary. Here we study the microorganisms associated with seed cotton and rice fermentation using a combination of culture-dependent and independent methods. Samples of cotton seed cauim were collected every 8 h of fermentation for analysis of the microbiota present during 48 h of fermentation. The bacterial population reached values around 8.0 log CFU / mL. A total of 162 strains of bacteria consisting of 33 morphotypes were isolated and identified by polyphasic taxonomy. The bacteria were grouped into Gram-negative (12.96%), Gram-positive catalase positive (58.15%) and Gram-positive catalase negative (41.84). Among the Gram-positive species *Lactobacillus* sp (13.47%), *Bacillus* sp (12.76%), *Lactobacillus plantarum* (11.34%), *Corynebacterium amycolatum* (10.63%), and *Lactobacillus agilis* (9.92%) were the most frequent microorganisms. The genus *Lactobacillus*, comprised the largest number of isolates and 15 different species. It was observed a decrease in pH value from 6.92 (T0) to 4.76 (T48 h) and the presence of lactic acid were only detected from the sample T40 h, reaching higher values in sample T48 (23.96 g / L). DGGE analysis were performed in order to observe the dynamics of communities of bacteria, revealing that there was a predominance of *Lactobacillus* throughout the fermentation. It was possible to observe that after 32 hours, the population did not modified significantly. The results showed by DGGE technique were in agreement with the identification by the culture dependent method.

* PEA1.40 Genotypic and technological characterization of Lactic acid bacteria from African fermented foods

David B Adimpong (1), DS Nielsen (1), KI Sørensen (2), PMF Derckx (2), L Jespersen (1)
(1) University of Copenhagen, Denmark
(2) Chr. Hansen A/S, Hørsholm, Denmark

Fermented indigenous African foods consist of those groups of foods that are prepared in homes, villages and small scale cottage industries. The fermentation is done to improve flavour, texture and prolong shelf-life among others. The predominant genera in fermented indigenous African foods are mainly *Lactobacillus* (both homo- and hetero-fermentative), *Leuconostoc* and to a lesser extent *Pediococcus*, *Lactococcus* and *Weissella*. The aim of this study was to characterize LAB from fermented indigenous African foods by molecular techniques and explore their technological properties which could result in the selection of appropriate multifunctional starter cultures. 33 isolates from 3 different fermented indigenous African foods; namely fermenting cocoa beans, dolo/pito wort, and a cereal based dough, were grouped using rep-PCR amplification technique and then followed by sequencing of the 16S rRNA gene. Species-specific oligonucleotides based on the *recA* gene was used to identify strains belonging to *Lactobacillus plantarum* and *Lactobacillus paraplantarum* which shows over 98% percentage identity of the 16S rRNA sequence for both species. Similarly, *Pediococcus acidilactici* and *Pediococcus pentosaceus* strains were identified with species-specific oligonucleotides based on the 16S rRNA and *ldhD* genes. Technological characterization included growth speed, milk acidification, inhibition of pathogenic and spoilage microorganisms, carbohydrates fermentation patterns, minimum inhibitory concentration of bile salt, selection of bile salt mutant strains and enzymatic potentials using the API ZYM kit (bioMérieux, Inc). The isolates were able to ferment a wide range of complex carbohydrates. 50% of the isolates were able to inhibit growth of *Listeria monocytogenes* 4140, *Escherichia coli* and *Salmonella typhimurium*. 15 strains belonging to *Lb. plantarum*, *Lb. paraplantarum*, *Ped. pentosaceus*, and *Ped. acidilactici* were able to grow on MRS agar with bile salt concentration of 2%. 16 of the 19 substrates used to assay the enzymatic potentials were hydrolysed with concentrations ranging from 5nmoles to over 40nmoles. None of the strains was able to produce the following enzymes; β -glucuronidase, α -mannosidase and α -fucosidase. Some of the strains are selected for further technological characterization with the intention to develop new starter cultures and probiotic strains for both the European and African food and feed industry.