



The effect of long- and short term exposure to laser light at 1070 nm on growth of *Saccharomyces cerevisiae*

Aabo, Thomas Ask; Perch-Nielsen, Ivan R.; Dam, J. Seidelin; Palima, D.Z.; Siegumfeldt, Henrik; Glückstad, Jesper; Arneborg, Nils


Published in:
22nd International ICFMH Symposium Food Micro 2010

Publication date:
2010


Document version
Early version, also known as pre-print

Citation for published version (APA):
Aabo, T. A., Perch-Nielsen, I. R., Dam, J. S., Palima, D. Z., Siegumfeldt, H., Glückstad, J., & Arneborg, N. (2010). The effect of long- and short term exposure to laser light at 1070 nm on growth of *Saccharomyces cerevisiae*. In *22nd International ICFMH Symposium Food Micro 2010* (pp. 308)

22nd International ICFMH Symposium Food Micro 2010



Copenhagen 30th August - 3rd September



Final Programme & Abstract Book



www.foodmicro.dk

	PEC1.24		PEC1.54	Zurera G	PEC2.60
	PEB2.20		PEC2.20	Zwartkruis A	PEC2.05
	PEE2.08		PED2.32	Zwartkruis-Nahuis A	PEC1.03
e, M	PSA1.04		PED2.22	Zwietering M	PEC1.29
	PEC1.57	Wille C	PEA2.47		PEC2.27
	PEC2.39	Wilson, D	PSD2.01		PEA1.55
	PSD1.02	Wingstrand A	PEC2.51		PEC2.55
	PEA1.34	Witthuhn C	PEA2.07		PSD2.02
	PEC1.25		PEC1.06	Økland M	PEC1.56
	PED1.26		PEC1.23		PEC1.86
	PEE2.17		PED2.02	Østlie H	PEA1.17
	PEE2.23		PED2.03	Østlie Hilde	PEA1.72
	PED2.55	Wolkers-Rooijackers J	PEA1.55	Aabo S	PEC1.26
	PEE2.14	Wongkrajang K	PEA1.08		PEC1.68
	PEA1.17	Wouters Dorrit	PEA1.22		PED2.14
	PSE1.03	Xavier D	PEA1.33		PED2.40
	PEC1.06	Xu Y Zh	PEA2.18		PED2.54
	PEB2.44	Yadav AS	PED2.24	X Aabo Thomas	PED2.39
	PED1.30	Yakhchali Bagher	PEA1.01	Aarts HK	PED1.36
	PEA1.21	Yamasaki Shinji	PEC1.77	Aarts, H	PSC1.04
	PEB1.14	Yang J-Y	PEC2.03		
	PED1.15	Yasui C	PEE2.07		
	PED1.11	Yavarmanesh Masoud	PEC2.45		
	PEA1.12	Yazdankhah S	PEC2.08		
	PEC1.95	YOON H-s	PEE2.20		
	PEC1.98	Yoon H-s	PEE2.21		
	PEC1.99	Ýpek D	PEC2.63		
	PEB2.32	Ýpek D	PED2.62		
	PEC2.06	Yuste J	PEB2.28		
	PEB1.02	Yuste, J	PSB2.03,		
	PEC1.98	Z. Palima D	PED2.39		
	PEC1.95	Zakrzewska A	PEB2.04		
	PEB1.20	Zali M	PEB1.27		
?	PEB1.04	Zali N	PEB1.27		
ia	PSB1.03	Zamfir M	PEA1.22		
	PEE2.25	Zannini E	PEA1.47		
	PEB1.02	Zannini Emanuele	PEA2.27		
	PSA2.06	Zelenik K	PEC1.84		
	PEB1.34	Zeller-Peronnet V	PEE2.11		
HJ	PEC2.55	Zenisova K	PEA1.05		
	PEC2.55	Zenoni S	PEB2.37		
	PEC1.58	Zeppa G	PEA1.45		
	PEC1.91	Zeppa G	PEA2.42		
	PSB1.01	Zhou, K	PSC1.02		
	PSB2.06	Zhu X	PEB2.36		
	PED2.16	Zilberstein, G	PSD1.05		
	PED1.13	Zinno Paola	PED2.20		
	PED1.14	Zorba M	PEC2.57		
	PEB1.07	Zotta Teresa	PEB2.25		
	PED1.12		PEB2.26		
	PEB2.04	Zoumpopoulou G	PEA1.11		
	PEB1.08	Zounli P	PEB1.24		
	PED2.43	Zuliani Veronique	PED2.38		
	PED2.44	Zunabovic M	PEC1.88		
	PEC2.26	Zurera G	PEC2.02		

- * PED2.39 The effect of long- and short term exposure to laser light at 1070 nm on growth of *Saccharomyces cerevisiae*
 Thomas Aabo (1), I R. Perch-Nielsen (3), J Seidelin Dam (2), D Z. Palima (2), H Siegumfeldt (1), J Glückstad (2), N Arneborg (1)
 (1) Copenhagen University, Institute of Food Science, DK-1958 Frederiksberg C, Denmark
 (2) DTU Fotonik, Department of Photonics Engineering, Technical University of Denmark, DK-2800 Kongens Lyngby, Denmark
 (3) DTU Nanotech, Department of Micro- and Nanotechnology, Technical University of Denmark, DK-2800 Kongens Lyngby, Denmark

As a preliminary step, before using a biophotonics workstation for studying microbial interactions within food systems, we have investigated the effect of a 1070 nm continuous wave Ytterbium fiber laser on exponentially growing *Saccharomyces cerevisiae* yeast cells over a span of 4 hours. The cells were immobilized onto Concanavalin A covered microscope slides and the growth was measured using the area increase of the cells in 2D. Using a continuous dual beam plane wave with a uniform spatial intensity distribution, we found that a continuous radiant flux through a single cell as low as 0.5 mW in 1.5 hours significantly changed the growth and division rate of *S. cerevisiae*. With the dual beam setup used we were able to successfully manipulate single *S. cerevisiae* cells in 3 dimensions with a minimum flux thorough the cell of 3.5 mW. In the regime investigated from 0.7 mW to 2.6 mW we found no threshold for the photo damage, but rather a continuous response to the increased accumulated dose.

- PED2.40 Growth of *Salmonella* in minced meat after freezing
 Søren Aabo (1), T Beck Hansen (1), T Birk (1),
 (1) National Food Institute, Technical University of Denmark, Denmark

Adaptive response to environmental changes is fundamental for bacterial survival and differs between exponential and stationary phase cells, with stationary phase cells being generally more resistant. Bacterial survival under food environmental stress is typically performed with stationary phase cells, however, during slaughter; bacteria in the feces may appear in mixed growth phases. One of the most commonly used preservative techniques for meat is freezing, which arrests bacterial growth and reduces the number of bacteria. The potential for survival and growth of *Salmonella* frozen in exponential phase is unknown.

Aims: One aim of this study was to investigate the survival of exponential and stationary phase cells of *Salmonella* in minced meat after freezing and thawing. Secondly, the aim was to investigate the lag-phase and the growth rate of exponential and stationary phase *Salmonella* cultured at 25°C in the juice drip from the defrosted meat.

Methods: Minced meat was inoculated with exponential and stationary phase cells of *Salmonella* and frozen at -20°C. The meat was defrosted at 5°C after 0, 2, 8, 20 and 34 days and bacterial counts were performed for measuring of survival. The meat juice was collected and incubated at 25°C, and bacterial numbers were established after 0, 2, 4, 6, 8, 10 and 27 hours. Estimation of the lag phases during growth in meat juice was performed using the Baranyi and Roberts model from DMFit.

Results: Significant difference of survival between the exponential and stationary cells was found after freezing. No reduction of stationary cells was observed after 20 days whereas exponential cells were reduced up to 2 log₁₀ units. A difference in lag phase for the exponential and stationary cells was identified, during culturing in the meat juice but depended on the length of the freezing period prior to culturing. After freezing for 2 and 5 days at -20°C the average lag phase was estimated to 2.5 and 5.3 hours for the exponential and stationary cells, respectively. When the meat had been stored for 20 days at -20°C the differences in lag phases were less pronounced having lag phases of 5.1 and 6.4 hours for the exponential and stationary cells, respectively.

- PED2.41 Safety in
 Maria Da
 (1) Univer

In Burundi and in sub-presence of toxic cyan are processed followin tion is an important st natural fermentations cassava foods are subj HCN equivalent/kg dw, removal during proces wide worldwide area a government and interr economical, nutritional foods and to stimulate nologies in the original This paper reports the p and quality of cassava t We compared the cyan *cerevisiae* selected strai The cyanide removal st cyanides contents (fror cyanide content was det compounds were also t imeter methods respect The results showed tha 2) using selected strains 3) *Saccharomyces cerev*. These preliminary result quality and safety imprc

- PED2.42 Challenge:
 Maria I. Gil
 (1) CEBAS-1

Chlorine is the most wic toxicity of chlorination t fective and innocuous al able by-product residues by-product formation h: absence in the produce. be prohibited if it is not but it is important that and therefore achieving disinfection agent durin agents is chlorine dioxid information has been pu potential formation of tr (3 mgL⁻¹) did not negati preserved after 10 days c washing with sodium hy 700mgL⁻¹). Moreover, trik applied. On the other har water, avoiding cross-co recommended if adequat