



Once in a lifetime

strategies for preventing re-replication in prokaryotic and eukaryotic cells.

Nielsen, Olaf; Løbner-Olesen, Anders

Published in:
EMBO Reports

DOI:
[10.1038/sj.embor.2008.2](https://doi.org/10.1038/sj.embor.2008.2)

Publication date:
2008

Document version
Early version, also known as pre-print

Citation for published version (APA):
Nielsen, O., & Løbner-Olesen, A. (2008). Once in a lifetime: strategies for preventing re-replication in prokaryotic and eukaryotic cells. *EMBO Reports*, 9(2), 151-156. <https://doi.org/10.1038/sj.embor.2008.2>

Once in a lifetime: strategies for preventing re-replication in prokaryotic and eukaryotic cells

Olaf Nielsen¹⁺ & Anders Løbner-Olesen²⁺⁺

¹University of Copenhagen, Copenhagen, Denmark and ²Roskilde University, Roskilde, Denmark

DNA replication is an extremely accurate process and cells have evolved intricate control mechanisms to ensure that each region of their genome is replicated only once during S phase. Here, we compare what is known about the processes that prevent re-replication in prokaryotic and eukaryotic cells by using the model organisms *Escherichia coli* and *Schizosaccharomyces pombe* as examples. Although the underlying molecular details are different, the logic behind the control mechanisms is similar. For example, after initiation, crucial molecules required for the loading of replicative helicases in both prokaryotes and eukaryotes are inactivated until the next cell cycle. Furthermore, in both systems the β -clamp of the replicative polymerase associates with enzymatic activities that contribute to the inactivation of the helicase loaders. Finally, recent studies suggest that the control mechanism that prevents re-replication in both systems also increases the synthesis of DNA building blocks.

Keywords: initiation of DNA replication; helicase loading; *Escherichia coli*; *Schizosaccharomyces pombe*; DNA building blocks
EMBO reports (2008) 9, 151–156. doi:10.1038/embor.2008.2

Introduction

Genomic DNA is organized differently in prokaryotic and eukaryotic cells. The bacterium *Escherichia coli* contains a single circular chromosome and replication is initiated bi-directionally from a fixed origin (*oriC*; Fig 1). Consequently, a single initiation event will ensure replication of the entire 4.6 Mbp genome in a process that is completed when the two divergent replication forks collide at the opposite side of the circular chromosome.

In eukaryotic cells, the genomic DNA is distributed between multiple chromosomes that are contained within the nucleus. At S phase, replication is simultaneously initiated from many different origins that are scattered throughout the genome (Fig 1), and replication is completed when all replication forks have either met a convergent

fork from an adjacent origin or reached the telomeres at the end of the chromosomes.

In the fission yeast *Schizosaccharomyces pombe*, many potential replication origins are found in intragenic regions of the 12.5 Mbp genome distributed across three chromosomes. However, only a limited subset of these is used in a given S phase. Furthermore, the specific origins that actually fire vary from one S phase to another, suggesting that origin selection occurs by a stochastic mechanism.

Replicating once, and only once

Despite these differences in organization, both cell types are faced with the challenge of ensuring that the entire genome is replicated once, and only once, in any given S phase. At face value, this problem seems to be different depending on whether cells have a single origin or many scattered ones. However, under optimal nutritional conditions, *E. coli* cells are able to grow with a doubling time that is much shorter than the time required for replication and segregation of the chromosome (S + G₂ phases). Consequently, initiation of replication occurs one, two or even three generations before cell birth, depending on the growth rate (Cooper & Helmstetter, 1968). Fast-growing cells are therefore born with chromosomes containing several active origins of replication, and such cells are also able to coordinate initiation at multiple—but identical—origins (Fig 1).

In both *E. coli* and fission yeast, initiation of replication is coupled to cell growth and is triggered by a specific signal that is generated when the cell has obtained a critical mass. Once activated, each replication origin is inhibited from re-firing until the next S phase (see below). Furthermore, in eukaryotic cells, passive replication by an incoming fork also prevents an origin that has not yet fired from firing until the next S phase. Together, these mechanisms ensure that the entire genome is replicated only once in each cell cycle.

Mechanisms of initiation

Initiation of replication in both *E. coli* and *S. pombe* occurs by a series of discrete steps. First, the origin recognition complex (ORC) is formed by the recruitment of replication factors to origin sequences. Subsequently, loading of the replicative helicase converts the ORC into a pre-replicative complex (pre-RC)—a process often referred to as ‘licensing’. This paves the way for loading of the polymerase itself; in both organisms, the crucial step seems to be loading of the helicase.

¹Cell Cycle and Genome Stability Group, Department of Molecular Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 Copenhagen K, Denmark

²Department of Science, Systems and Models, Roskilde University, Universitetsvej 1, PO Box 260, DK-4000 Roskilde, Denmark

*Corresponding author. Tel: +45 3532 2102; Fax: +45 3532 2128; E-mail: onigen@my.molbio.ku.dk

**Corresponding author. Tel: +45 4674 2615; Fax: +45 4674 3011; E-mail: lobner@ruc.dk

Submitted 21 November 2007; accepted 3 January 2008




Fig 1 | Initiation of replication at multiple origins in *Escherichia coli* and *Schizosaccharomyces pombe*. Origins that have not yet initiated are shown in green, whereas those initiated or passively replicated are shown in red. The *E. coli* cell is fast-growing with S + G₂ phases spanning more than two generation times. Consequently, initiations occur in synchrony at four cellular origins. For simplicity, only one chromosome with six autonomously replicating sequences is shown in the *S. pombe* cell. Four of these are firing, whereas two are being passively replicated.

The *S. pombe* ORC is a six-subunit complex that consists of the three proteins Orc1, Orc2 and Orc4, each of which contain an AAA⁺ ATP-binding domain (Kong & DePamphilis, 2002). Orc proteins specifically associate with origins of replication (autonomously replicating sequences (ARSs); Fig 2). One of the ORC subunits, Orc4, contains several AT-hook motifs in its amino-terminal domain, which interact with the AT-rich origin sequences. The ORC is bound to chromatin throughout the cell cycle and therefore its binding is unlikely to regulate initiation (Lygerou & Nurse, 1999). In *E. coli*, the ORC is formed by DnaA, which is the only protein that is specific to replication initiation at *oriC*. The DnaA protein associated with either ATP or ADP binds to three 9-bp binding sites within the origin of replication called R1, R2 and R4 (reviewed by Kaguni, 2006; Mott & Berger, 2007). Similar to the situation in yeast, the ORC remains bound to the origin of replication throughout most of the cell cycle (Samitt *et al*, 1989).

The transition from the ORC-like stage to a pre-RC stage represents the next step in the initiation process. In fission yeast, the pre-RC is formed in late M and G₁ phases, when the two initiation factors Cdt1 and Cdc18 facilitate loading of the replicative helicase. Both proteins are cell-cycle regulated, being absent from late S phase until cells exit mitosis (see below). Cdc18 is yet another AAA⁺ ATPase, whereas the biochemical function of Cdt1 is unknown. Our understanding of pre-RC formation is largely based on studies of Cdc6, the budding yeast orthologue of Cdc18, but we anticipate that the mechanism in fission yeast is similar. The presence of Cdc18 and Cdt1 at the ORC enables the recruitment of several complexes of minichromosome maintenance (MCM) 2–7 hexamers, in a cyclic process that requires hydrolysis of ATP on both Cdc18 and on ORC subunits (Randell *et al*, 2006). The MCM2–7 complex is believed to act as the replicative helicase, although biochemical evidence for this is still circumstantial.

Assembly of the replisome at the pre-RC and activation of the replication process involve several additional factors, including Cdc45 and the GINS complex (reviewed by Legouras *et al*, 2006). In addition, the activity of two protein kinases is required at this stage: cyclin-dependent protein kinase (Cdk)—the main driver of the cell cycle in fission yeast—and the conserved Hsk1–Dfb1 kinase (also known as Ddk). The phosphorylation targets for Cdk were recently identified as Sdl2 and Sdl3 in budding yeast (Zegeerman & Diffley, 2007; Tanaka *et al*, 2007).

In *E. coli*, pre-RC formation is initiated by further binding of DnaA to the weaker recognition sites within the origin—that is, R3 and R5, which are indifferent to the nucleotide-bound status of DnaA—and to three I-boxes (McGarry *et al*, 2004; Kawakami *et al*, 2005) that are specific for DnaA-ATP. With the help of the accessory proteins IHF, HU and DiaA (Ryan *et al*, 2002; Keyamura *et al*, 2007), this induces formation of a DnaA–DNA nucleoprotein complex on *oriC*, where the DNA is remodelled to a right-handed DNA wrapped around a right-handed DnaA-ATP filament (Erzberger *et al*, 2006). The DnaA–DNA complex promotes duplex opening in an adjacent AT-rich region. This open complex is stabilized by the binding of DnaA-ATP to specific 6-bp sequences found in the single-stranded region (Speck & Messer, 2001). The requirement for DnaA-ATP in origin remodelling explains why this configuration of the protein is limiting for initiation *in vivo* (Nishida *et al*, 2002; Riber *et al*, 2006). Subsequently, the DnaA protein recruits the hexameric DnaB helicase associated with ATP-bound DnaC as a B₆C₆ complex to the single-stranded region of the open complex. DnaC loads the DnaB helicase on the open complex to promote further duplex opening to form the pre-RC stage. During this process, ATP is hydrolysed and DnaC is released. In *E. coli*, there does not seem to be any control on the pre-RC stage and the transition to replication proceeds immediately by the loading of two or three DNA polymerase III holoenzymes on the origin (McInerney *et al*, 2007). When dNTPs are present, replication can then commence (Herrick & Sclavi, 2007).

The spatial arrangement of DnaA protein domains involved in nucleotide binding, DNA binding and oligomerization is similar to the fission yeast initiation factor Cdc18, and it has been suggested that the helical DNA-binding domain could direct similar functioning AAA⁺ domains to their respective origins (Erzberger *et al*, 2002). It is worth noting that both DnaA and Cdc18 can switch between active and inactive configurations depending on the nature of the bound nucleotide, and that this molecular switch is one of the determinants for initiation control.

Cascades of initiation

As discussed above, rapidly growing *E. coli* cells contain many origins of replication that all fire simultaneously (Fig 1), and synchronous initiation presumably results from the release of the DnaA protein from the first initiated origin in a cell. This will momentarily increase the DnaA:*oriC* ratio for remaining ‘old’ origins and their initiation will follow in a cascade-like manner—known as the initiation cascade (Løbner-Olesen *et al*, 1994). Eukaryotic cells are faced with a similar problem: origins are selected by a stochastic mechanism, and therefore there is a risk that large chromosomal regions will occasionally remain unreplicated during any given S phase. Analogous to the initiation cascade model, it has been proposed that a crucial replication factor is rate limiting for initiation. As replication proceeds, this factor is released and can be redistributed to other origins, thereby increasing their probability of firing (Lucas




Fig 2 | Mechanisms of replication initiation, and the prevention of re-replication in *Escherichia coli* and *Schizosaccharomyces pombe*. The symbols for the different replication proteins are indicated in the figure with the exception of the replicative DNA polymerase (green) and the DNA-loaded β -clamp–proliferating cell nuclear antigen (PCNA; yellow rings). The replicative DNA polymerase is indicated as a dimer, although recent evidence suggests, at least in *E. coli*, that the replicase might be a trimer (McInerney *et al*, 2007). For simplicity, DnaA and Cdc18 have been given the same symbol as these proteins share homology in important functional domains. In the steps after pre-replicative complex formation, the *E. coli* process is shown on the upper strand/leftward fork, whereas the fission yeast system is illustrated by the lower strand/rightward fork. In *E. coli*, the newly synthesized DNA is unmethylated at GATC sites (blue, middle panel)—that is, DNA is hemi-methylated immediately after passage of the replication fork. As the fork progresses further away, both strands become methylated (lower panel). In fission yeast, Cdk phosphorylation of ORC and Cdc18 (indicated by 'P') prevent binding of the MCM2–7 complex to the ORC and causes SCF-mediated destruction of Cdc18. ARS, autonomously replicating sequence; CDK, cyclin-dependent kinase; MCM, minichromosome maintenance; ORC, origin recognition complex; Pre-RC, pre-replicative complex; RIDA, regulatory inactivation of DnaA; SCF, an E3 ubiquitin ligase.

et al, 2000). The nature of this factor has not been established, but it would have to be a protein that is not degraded in the initiation process—that is, not Cdt1 or Cdc18 (see below).

Mechanisms to prevent re-initiation

In both *E. coli* and *S. pombe*, inactivation of the helicase loader proteins has a crucial role in preventing immediate re-firing of a recently

activated origin. In *E. coli*, the DnaA protein is the target for this regulation, whereas fission yeast cells regulate both Cdt1 and Cdc18. In both systems, inhibition of re-replication is accomplished both by physically preventing pre-RC assembly and by a reduction in the activity of AAA⁺ ATPase proteins. This is mediated by post-translational inactivation of the proteins and by the modulation of gene expression.

Prevention of pre-RC assembly

The *E. coli* origin of replication is rich in GATC sites, which is the substrate for the Dam methyltransferase. As methylation is a post-replicative process, newly replicated origins are methylated on only one strand (Fig 2). These hemi-methylated origins are bound (sequestered) by SeqA, a protein with high affinity for hemi-methylated GATC sites (Lu *et al*, 1994). Sequestration renders the origin inaccessible to DnaA for approximately one-third of the generation time to prevent immediate re-initiation (Campbell & Kleckner, 1990; von Freiesleben *et al*, 2000).

Origin sequestration is instrumental not only in preventing the immediate re-initiation at an origin, but also in preparing the origin for the next round of initiation. During sequestration, the DnaA protein is only able to bind the high affinity sites R1, R2 and R4, to re-set the origin to the ORC stage (Nievera *et al*, 2006). Although sequestration lasts less than one generation, it ensures that successive initiations at the same origin are separated by a doubling time, because it provides a time window during which the origin cannot be initiated and the amount of DnaA-ATP is reduced to a level that cannot sustain initiation (see below). Consequently, a period of growth is necessary before origins of replication are released from sequestration and can re-initiate.

In eukaryotic cells, the increase in Cdk activity that initiates S phase has an additional function in preventing re-replication during S, G2 and M phases. The importance of this mechanism follows from the observation that G2 cells can be manipulated to erroneously enter another round of S phase by temporarily inactivating a temperature-sensitive Cdk allele (Hayles *et al*, 1994). Cdk seems to inhibit re-initiation of DNA replication partly by phosphorylating subunits in the ORC (Fig 2), thereby preventing *de novo* assembly of pre-RCs until Cdk activity becomes low again as cells exit mitosis (Nguyen *et al*, 2001; Vas *et al*, 2001). Phosphorylation of the ORC has not been reported to involve a sequestration mechanism as in *E. coli*; presumably it simply prevents recruitment of Cdt1 and Cdc18.

Post-translational inactivation of AAA⁺ proteins

The activity of the DnaA protein is reduced during S phase by a process known as the 'regulatory inactivation of DnaA' (RIDA; Fig 2), in which the active ATP-bound DnaA protein is converted to the inactive ADP-bound form by ATP hydrolysis (Katayama *et al*, 1998). RIDA activity involves two proteins: the DnaA-related protein Hda (Kato & Katayama, 2001) and the β -clamp of the DNA polymerase (Pol) III holoenzyme (encoded by the *dnaN* gene; Katayama *et al*, 1998). These proteins form a complex even before the clamp is loaded onto the DNA (Kawakami *et al*, 2006). However, only the DNA-loaded β -subunit of Pol III in complex with the Hda protein stimulates the ATPase activity of DnaA to promote conversion of DnaA-ATP to the inactive DnaA-ADP (Su'etsugu *et al*, 2004). At the end of the initiation process, hydrolysis of DnaA-ATP by RIDA is accelerated because new replication forks are formed, and more β -clamps are loaded onto the DNA.

In fission yeast, the two helicase-loader proteins Cdc18 and Cdt1 also become inactivated after initiation of replication, but here this is accomplished by physical degradation rather than biochemical inactivation. The increase in Cdk activity that brings about S phase also causes phosphorylation of Cdc18, which targets the protein for SCF-mediated ubiquitination and subsequent degradation by proteolysis (Fig 2; Jallepalli *et al*, 1997). The importance of this regulation is clear from the fact that ectopic over-production of Cdc18 causes massive re-initiation of DNA replication (Nishitani & Nurse, 1995).

The Cdt1 protein also becomes degraded after successful initiation of DNA replication, but this process does not require Cdk. Instead, Cdt1 is targeted for degradation by a different E3 ubiquitin ligase, the Cullin4–Ddb1–Roc1 complex (Ralph *et al*, 2006). Interestingly, Cdt1 ubiquitination is tightly coupled to its function in initiation by means of two different mechanisms. First, substrate recognition requires a specific adaptor protein, the WD40-repeat protein Cdt2, which becomes transcriptionally induced when cells enter S phase (Liu *et al*, 2005). Second, Cdt1 only becomes ubiquitinated when it is associated with the proliferating cell nuclear antigen (PCNA) processivity clamp of the polymerase (Fig 2; Arias & Walter, 2006; Jin *et al*, 2006; Nishitani *et al*, 2006; Senga *et al*, 2006). Presumably, Cdt1 molecules are consumed when they have been actively engaged in initiation. Therefore, enzymatic activities that negatively regulate helicase-loader proteins seem to associate with the processivity clamp in both *E. coli* and *S. pombe*.

Modulation of gene expression

E. coli does not seem to regulate DnaA activity by degrading the protein; however, in addition to RIDA, a second mechanism for reducing DnaA activity in the post-initiation period exists. This method uses the sequestration mechanism to reduce expression of the *dnaA* gene. On replication, the *dnaA* gene promoter region, which is rich in GATC sequences, is hemi-methylated and sequestered for the same time period as the origin of replication. Sequestration of the *dnaA* promoter completely blocks transcription of the *dnaA* gene (Campbell & Kleckner, 1990). As the *dnaA* gene is close to the origin, sequestration of *dnaA* is virtually coincident with sequestration of *oriC*, and *de novo* DnaA synthesis is prevented during the origin sequestration period (Fig 2). In cells in which origin and *dnaA* gene sequestration no longer coincide, DnaA synthesis continues during origin sequestration. In such cells, re-initiations occasionally occur at some origins within the same cell cycle (Riber & Løbner-Olesen, 2005).

Transcription of the genes encoding the helicase-loader proteins Cdc18 and Cdt1 also oscillates in fission yeast and is high in late mitosis and G1 (Hofmann & Beach, 1994; Kelly *et al*, 1993). However, this is actively controlled by the cell-cycle-regulated MBF transcription factor complex rather than by an intricate system that monitors ongoing replication.

Titration of DnaA to reservoir sites

During origin sequestration, replication generates new DnaA protein-binding sites outside *oriC*. These titrate DnaA protein away from the origin and, in the absence of *de novo* DnaA synthesis (Campbell & Kleckner, 1990), efficiently reduce the intracellular concentration of DnaA protein available for initiation (Fig 2). The *E. coli* chromosome contains a hierarchy of 308 evenly distributed R-type DnaA boxes with different affinities for the DnaA protein. The *datA* locus, which contains five R-type DnaA boxes, seems to have the highest DnaA-binding capacity, and might bind to several hundred molecules of DnaA protein associated with either ATP or ADP. The *datA* locus is

located approximately 470Kbp away from *oriC* and is replicated within the period of origin sequestration during which no new DnaA protein is synthesized. This generates a sink for free DnaA protein (Kitagawa *et al*, 1998).

Coupling nucleotide synthesis to chromosome replication

In most cells, the intracellular concentration of DNA precursors (dNTPs) is low and can only sustain limited chromosome replication unless they are continuously synthesized to match the demand of ongoing replication forks. Upregulation of dNTP synthesis in S phase is carefully controlled because imbalances between the four individual nucleotide pools, as well as balanced deviation from the normal level, are mutagenic (reviewed by Mathews, 2006). In both *E. coli* and fission yeast, dNTPs are synthesized from their corresponding NTPs exclusively by the ribonucleotide reductase (RNR) complex. RNR is a heterodimeric tetramer consisting of two large and two small subunits. RNR activity is the rate-limiting step in dNTP synthesis.

The RNR subunits of *E. coli* are encoded by the *nrdAB* operon, and *nrdAB* expression is adjusted to DNA synthesis (reviewed by Herrick & Sclavi, 2007). Transcription of *nrdAB* is induced at the time of initiation by a DnaA-independent mechanism (Jacobson & Fuchs, 1998). Superimposed on this cell-cycle regulation is modulation of transcription by the DnaA protein (Augustin *et al*, 1994). DnaA was initially reported to stimulate *nrdAB* transcription (Jacobson & Fuchs, 1998) although a recent study indicates that DnaA-ATP—but not DnaA-ADP—is an efficient repressor of its transcription (Gon *et al*, 2006). The *nrdAB* expression level is therefore determined by the DnaA-ATP:DnaA-ADP ratio.

The RIDA-imposed variation in DnaA-ATP:DnaA-ADP ratio throughout the cell cycle (Kurokawa *et al*, 1999) could therefore couple dNTP synthesis to the elongation step of chromosome replication. Before initiation, when the cellular DnaA-ATP:DnaA-ADP ratio is high (Kurokawa *et al*, 1999), DnaA regulation would favour *nrdAB* repression. After initiation, RIDA is accelerated, resulting in a reduced DnaA-ATP:DnaA-ADP ratio and consequently an increase in *nrdAB* transcription. Therefore, the RNR level is increased in S phase, resulting in an increased synthesis of dNTPs to match the demand from the ongoing replication forks.

Precursor synthesis in eukaryotic cells is also adjusted to ongoing DNA replication by the regulation of RNR activity; however, the molecular basis is different. In *S. pombe*, transcription of the gene encoding the large subunit (Cdc22) is cell-cycle regulated (Fernandez Sarabia *et al*, 1993). In addition, assembly of RNR is actively prevented outside S phase by the presence of the RNR inhibitor protein Spd1 (Liu *et al*, 2003). When cells enter S phase, Spd1 is degraded by the same pathway that downregulates the Cdt1 helicase loader—that is, the Cullin4–Ddb1–Roc1 E3 ubiquitin ligase and the adaptor protein Cdt2 (Holmberg *et al*, 2005; Liu *et al*, 2005). It is unclear whether the degradation of Spd1—similar to the degradation of Cdt1—is coupled to PCNA.

Perspectives

The development of the eukaryotic type of genome organization—with multiple chromosomes and many scattered origins of replication—was probably important for the expansion of genome size that allowed the development of complex organisms. Taken at face value, control of replication seems to be organized differently in prokaryotic and eukaryotic cells; however, the control mechanisms found

in the two systems seem to regulate the same steps in the process. First, in both prokaryotes and eukaryotes the crucial step in the establishment of a replication origin is loading of the replicative helicase. This process is mediated when the concentration of the helicase-loading AAA⁺ ATPases builds up to a certain threshold in the cell. In *E. coli*, this seems to be the rate-limiting step; loading of the replicative polymerase and initiation immediately follows. In *S. pombe*, further progress requires the action of S-phase-activating kinases. Second, once an origin of replication has fired, re-firing is prevented for a period of time. In both systems, this is accomplished by a combination of physical modification of the origin and/or associated protein factors (by sequestration or by phosphorylation), such that the helicase loader cannot access it, and by removing the helicase-loader activity. Eukaryotic cells literally get rid of the protein by switching on ubiquitin-mediated degradation. The prokaryotic cell does not have this option and therefore it is dependent on several other methods of reducing the active concentration of the helicase loader, such as through hydrolysis of its bound ATP, binding of the loader to unproductive sites or downregulation of its expression. The development of ubiquitin-mediated protein degradation made these mechanisms redundant.

In this review, we have attempted to draw parallels between the basic mechanisms that prevent re-replication in two simple unicellular model organisms. Failure to restrict replication to once per cell cycle leads to DNA damage through the generation of double-stranded breaks and can result in development of tumours (reviewed by Arias & Walter, 2007). It is therefore not surprising that metazoans have evolved additional mechanisms—such as inactivation of Cdt1 by Geminin binding—to minimize the likelihood of untimely replication initiations.

ACKNOWLEDGEMENTS

We thank R. Egel, S. MacNeill and O. Fleck for their comments on the manuscript. Work in our laboratories is supported by The Danish Cancer Society, the Lundbeck Foundation, The Novo-Nordisk Foundation, the Danish Medical Research Council and the Danish Natural Science Research Council.

REFERENCES

- Arias EE, Walter JC (2006) PCNA functions as a molecular platform to trigger Cdt1 destruction and prevent re-replication. *Nat Cell Biol* **8**: 84–90
- Arias EE, Walter JC (2007) Strength in numbers: preventing rereplication via multiple mechanisms in eukaryotic cells. *Genes Dev* **21**: 497–518
- Augustin LB, Jacobson BA, Fuchs JA (1994) *Escherichia coli* Fis and DnaA proteins bind specifically to the *nrd* promoter region and affect expression of an *nrd*–lac fusion. *J Bacteriol* **176**: 378–387
- Campbell JL, Kleckner N (1990) *E. coli* *oriC* and the *dnaA* gene promoter are sequestered from dam methyltransferase following the passage of the chromosomal replication fork. *Cell* **62**: 967–979
- Cooper S, Helmstetter CE (1968) Chromosome replication and the division cycle of *Escherichia coli* B/r. *J Mol Biol* **31**: 519–540
- Erzberger JP, Pirruccello MM, Berger JM (2002) The structure of bacterial DnaA: implications for general mechanisms underlying DNA replication initiation. *EMBO J* **21**: 4763–4773
- Erzberger JP, Mott ML, Berger JM (2006) Structural basis for ATP-dependent DnaA assembly and replication-origin remodeling. *Nat Struct Mol Biol* **13**: 676–683
- Fernandez Sarabia MJ, McNerny C, Harris P, Gordon C, Fantes P (1993) The cell cycle genes *cdc22+* and *suc22+* of the fission yeast *Schizosaccharomyces pombe* encode the large and small subunits of ribonucleotide reductase. *Mol Gen Evol* **238**: 241–251
- Gon S, Camara JE, Klungsoyr HK, Crooke E, Skarstad K, Beckwith J (2006) A novel regulatory mechanism couples deoxyribonucleotide synthesis and DNA replication in *Escherichia coli*. *EMBO J* **25**: 1137–1147
- Hayles J, Fisher D, Woollard A, Nurse P (1994) Temporal order of S phase and mitosis in fission yeast is determined by the state of the p34cdc2–mitotic B cyclin complex. *Cell* **78**: 813–822

- Herrick J, Sclavi B (2007) Ribonucleotide reductase and the regulation of DNA replication: an old story and an ancient heritage. *Mol Microbiol* **63**: 22–34
- Hofmann JF, Beach D (1994) *cdt1* is an essential target of the Cdc10/Sct1 transcription factor: requirement for DNA replication and inhibition of mitosis. *EMBO J* **13**: 425–434
- Holmberg C, Fleck O, Hansen HA, Liu C, Slaaby R, Carr AM, Nielsen O (2005) Ddb1 controls genome stability and meiosis in fission yeast. *Genes Dev* **19**: 853–862
- Jacobson BA, Fuchs JA (1998) A 45 bp inverted repeat is required for cell cycle regulation of the *Escherichia coli* *nrd* operon. *Mol Microbiol* **28**: 1307–1314
- Jallepalli PV, Brown GW, Muzi-Falconi M, Tien D, Kelly TJ (1997) Regulation of the replication initiator protein p65cdc18 by CDK phosphorylation. *Genes Dev* **11**: 2767–2779
- Jin J, Arias EE, Chen J, Harper JW, Walter JC (2006) A family of diverse Cul4–Ddb1-interacting proteins includes Cdt2, which is required for S phase destruction of the replication factor Cdt1. *Mol Cell* **23**: 709–721
- Kaguni JM (2006) DnaA: controlling the initiation of bacterial DNA replication and more. *Annu Rev Microbiol* **60**: 351–375
- Katayama T, Kubota T, Kurokawa K, Crooke E, Sekimizu K (1998) The initiator function of DnaA protein is negatively regulated by the sliding clamp of the *E. coli* chromosomal replicase. *Cell* **94**: 61–71
- Kato J, Katayama T (2001) Hda, a novel DnaA-related protein, regulates the replication cycle in *Escherichia coli*. *EMBO J* **20**: 4253–4262
- Kawakami H, Keyamura K, Katayama T (2005) Formation of an ATP-DnaA-specific initiation complex requires DnaA Arginine 285, a conserved motif in the AAA+ protein family. *J Biol Chem* **280**: 27420–27430
- Kawakami H, Su’etsugu M, Katayama T (2006) An isolated Hda–clamp complex is functional in the regulatory inactivation of DnaA and DNA replication. *J Struct Biol* **156**: 220–229
- Kelly TJ, Martin GS, Forsburg SL, Stephen RJ, Russo A, Nurse P (1993) The fission yeast *cdc18+* gene product couples S phase to START and mitosis. *Cell* **74**: 371–382
- Keyamura K, Fujikawa N, Ishida T, Ozaki S, Su’etsugu M, Fujimitsu K, Kagawa W, Yokoyama S, Kurumizaka H, Katayama T (2007) The interaction of DnaA and DnaA regulates the replication cycle in *E. coli* by directly promoting ATP DnaA-specific initiation complexes. *Genes Dev* **21**: 2083–2099
- Kitagawa R, Ozaki T, Moriya S, Ogawa T (1998) Negative control of replication initiation by a novel chromosomal locus exhibiting exceptional affinity for *Escherichia coli* DnaA protein. *Genes Dev* **12**: 3032–3043
- Kong D, DePamphilis ML (2002) Site-specific ORC binding, pre-replication complex assembly and DNA synthesis at *Schizosaccharomyces pombe* replication origins. *EMBO J* **21**: 5567–5576
- Kurokawa K, Nishida S, Emoto A, Sekimizu K, Katayama T (1999) Replication cycle-coordinated change of the adenine nucleotide-bound forms of DnaA protein in *Escherichia coli*. *EMBO J* **18**: 6642–6652
- Legouras I, Xouri G, Dimopoulos S, Lygeros J, Lygerou Z (2006) DNA replication in the fission yeast: robustness in the face of uncertainty. *Yeast* **23**: 951–962
- Liu C, Powell KA, Mundt K, Wu L, Carr AM, Caspari T (2003) Cop9/ signalosome subunits and Pcu4 regulate ribonucleotide reductase by both checkpoint-dependent and -independent mechanisms. *Genes Dev* **17**: 1130–1140
- Liu C, Poitelea M, Watson A, Yoshida SH, Shimoda C, Holmberg C, Nielsen O, Carr AM (2005) Transactivation of *Schizosaccharomyces pombe* *cdt2+* stimulates a Pcu4–Ddb1–CSN ubiquitin ligase. *EMBO J* **24**: 3940–3951
- Løbner-Olesen A, Hansen FG, Rasmussen KV, Martin B, Kuempel PL (1994) The initiation cascade for chromosome replication in wild-type and Dam methyltransferase deficient *Escherichia coli* cells. *EMBO J* **13**: 1856–1862
- Lu M, Campbell JL, Boye E, Kleckner N (1994) SeqA: a negative modulator of replication initiation in *E. coli*. *Cell* **77**: 413–426
- Lucas I, Chevrier-Miller M, Sogo JM, Hyrien O (2000) Mechanisms ensuring rapid and complete DNA replication despite random initiation in *Xenopus* early embryos. *J Mol Biol* **296**: 769–786
- Lygerou Z, Nurse P (1999) The fission yeast origin recognition complex is constitutively associated with chromatin and is differentially modified through the cell cycle. *J Cell Sci* **112**: 3703–3712
- Mathews CK (2006) DNA precursor metabolism and genomic stability. *FASEB J* **20**: 1300–1314
- McGarry KC, Ryan VT, Grimwade JE, Leonard AC (2004) Two discriminatory binding sites in the *Escherichia coli* replication origin are required for DNA strand opening by initiator DnaA-ATP. *Proc Natl Acad Sci USA* **101**: 2811–2816
- McInerney P, Johnson A, Katz F, O’Donnell M (2007) Characterization of a triple DNA polymerase replisome. *Mol Cell* **27**: 527–538
- Mott ML, Berger JM (2007) DNA replication initiation: mechanisms and regulation in bacteria. *Nat Rev Microbiol* **5**: 343–354
- Nguyen VQ, Co C, Li JJ (2001) Cyclin-dependent kinases prevent DNA re-replication through multiple mechanisms. *Nature* **411**: 1068–1073
- Nievera C, Torgue JJ, Grimwade JE, Leonard AC (2006) SeqA blocking of DnaA–oriC interactions ensures staged assembly of the *E. coli* pre-RC. *Mol Cell* **24**: 581–592
- Nishida S, Fujimitsu K, Sekimizu K, Ohmura T, Ueda T, Katayama T (2002) A nucleotide switch in the *Escherichia coli* DnaA protein initiates chromosomal replication: evidence from a mutant DnaA protein defective in regulatory ATP hydrolysis *in vitro* and *in vivo*. *J Biol Chem* **277**: 14986–14995
- Nishitani H, Nurse P (1995) p65cdc18 plays a major role controlling the initiation of DNA replication in fission yeast. *Cell* **83**: 397–405
- Nishitani H *et al* (2006) Two E3 ubiquitin ligases, SCF–Skp2 and DDB1–Cul4, target human Cdt1 for proteolysis. *EMBO J* **25**: 1126–1136
- Ralph E, Boye E, Kearsy SE (2006) DNA damage induces Cdt1 proteolysis in fission yeast through a pathway dependent on Cdt2 and Ddb1. *EMBO Rep* **7**: 1134–1139
- Randell JC, Bowers JL, Rodriguez HK, Bell SP (2006) Sequential ATP hydrolysis by Cdc6 and ORC directs loading of the Mcm2–7 helicase. *Mol Cell* **21**: 29–39
- Riber L, Løbner-Olesen A (2005) Coordinated replication and sequestration of oriC and dnaA are required for maintaining controlled once-per-cell-cycle initiation in *Escherichia coli*. *J Bacteriol* **187**: 5605–5613
- Riber L, Olsson JA, Jensen RB, Skovgaard O, Dasgupta S, Marinus MG, Løbner-Olesen A (2006) Hda-mediated inactivation of the DnaA protein and *dnaA* gene autoregulation act in concert to ensure homeostatic maintenance of the *Escherichia coli* chromosome. *Genes Dev* **20**: 2121–2134
- Ryan VT, Grimwade JE, Nievera CJ, Leonard AC (2002) IHF and HU stimulate assembly of pre-replication complexes at *Escherichia coli* oriC by two different mechanisms. *Mol Microbiol* **46**: 113–124
- Samitt CE, Hansen FG, Miller JF, Schaechter M (1989) *In vivo* studies of DnaA binding to the origin of replication of *Escherichia coli*. *EMBO J* **8**: 989–993
- Senga T, Sivaprasad U, Zhu W, Park JH, Arias EE, Walter JC, Dutta A (2006) PCNA is a cofactor for Cdt1 degradation by CUL4/DDB1-mediated N-terminal ubiquitination. *J Biol Chem* **281**: 6246–6252
- Speck C, Messer W (2001) Mechanism of origin unwinding: sequential binding of DnaA to double- and single-stranded DNA. *EMBO J* **20**: 1469–1476
- Su’etsugu M, Takata M, Kubota T, Matsuda Y, Katayama T (2004) Molecular mechanism of DNA replication-coupled inactivation of the initiator protein in *Escherichia coli*: interaction of DnaA with the sliding clamp-loaded DNA and the sliding clamp–Hda complex. *Genes Cells* **9**: 509–522
- Tanaka S, Umemori T, Hirai K, Muramatsu S, Kamimura Y, Araki H (2007) CDK-dependent phosphorylation of Sld2 and Sld3 initiates DNA replication in budding yeast. *Nature* **445**: 328–332
- Vas A, Mok W, Leatherwood J (2001) Control of DNA rereplication via Cdc2 phosphorylation sites in the origin recognition complex. *Mol Cell Biol* **21**: 5767–5777
- von Freiesleben U, Krekling MA, Hansen FG, Løbner-Olesen A (2000) The eclipse period of *Escherichia coli*. *EMBO J* **19**: 6240–6248
- Zegerman P, Diffley JF (2007) Phosphorylation of Sld2 and Sld3 by cyclin-dependent kinases promotes DNA replication in budding yeast. *Nature* **445**: 281–285



Olaf Nielsen



Anders Løbner-Olesen