



Isolation and characterization of antifungal dairy propionibacteria

Min, Min; Aunbjerg, Stina Dissing; Vogensen, Finn Kvist; Knøchel, Susanne

Publication date:
2012

Document version
Early version, also known as pre-print

Citation for published version (APA):
Min, M., Aunbjerg, S. D., Vogensen, F. K., & Knøchel, S. (2012). *Isolation and characterization of antifungal dairy propionibacteria*. Abstract from International ICFMH Symposium FoodMicro 2012, Istanbul, Turkey.





23rd INTERNATIONAL ICFMH SYMPOSIUM
FoodMicro 2012



Global Issues In Food Microbiology

Abstract Book



FoodMicro

3-7 SEPTEMBER 2012 ISTANBUL

re and bacteriocins on
Cytophaga, *Salmonella*
H7 in beef carpaccio

a MEDINA

be contaminated with pathogens
in beef products include *L.*
colli O157:H7. Combined
individual applications have been
food safety. In this work, high
bacteriocins present in supernatants of
bacteriocin-based preparations were
and *E. coli* O157:H7 and in beef
carpaccio was inoculated
450 MPa for 5 min was applied
independent trials, one for each
microbial counts were determined
The levels of the three pathogens
during refrigerated storage. HP
bacteriocins by 0.2 to 1.4 log units.
antimicrobial effect. Reductions of *S.*
bacteriocins were not active
activities were observed against *S.*
and *E. coli* O157:H7 by 2.7 log
units of bacteriocins. A synergistic
effect with the combinations of HP and
nisin, with inactivation rates
with the sum of the individual
450 MPa for 5 min would sensitize
carpaccio, exhibiting a synergistic
effect against *Cytophaga* or *S. Enteritidis*.

Bacteria for bioprotection

P-567

Isolation and characterization of antifungal dairy
propionibacteria

Min MIN, Stina Dissing AUNSBJERG, Finn Kvist VOGENSEN,
Susanne KNØCHEL

University of Copenhagen, Food Science, Copenhagen, DENMARK
Corresponding author: sofa@life.ku.dk

Spoilage of stored food caused by food-borne fungi is a huge problem in the food industry and can result in significant economic losses. In addition, fungi can cause severe health problems due to the production of mycotoxins by some moulds. The use of biopreservation to control food-borne fungi has gained increased interest due to consumers' requirements for reduced use of chemical preservatives. Propionibacteria are important organisms in specific food fermentations such as certain cheeses and other dairy products and some strains can in addition be used as biopreservatives due to their antimicrobial nature. The mechanism behind the antifungal activity of these organisms is not yet fully elucidated but a synergistic effect between produced antifungal metabolites is indicated as a likely explanation. Strains of dairy propionibacteria were isolated from Swiss raw milk Emmental cheeses and identified by the use of Rep-PCR, PFGE and 16S rDNA sequencing. Three restriction endonucleases were studied for their digestion of the isolated propionibacteria. The antifungal activity of the propionibacteria was tested against indicator yeasts and moulds isolated from foods with an overlay assay. Some of the propionibacteria displayed a marked inhibitory effect against moulds, whereas no or limited inhibition was seen against the indicator yeasts. Propionibacteria showing high anti-mould activity were selected for further analysis. The antifungal activity was highly influenced by the carbon source in the growth media, with the highest activity observed in media with the lowest end pH. An increase in incubation time of propionibacteria from 3 to 8 days prior to inhibition tests increased antifungal activity. However, although activity could still be observed after 17 days of incubation, some of the antifungal effect was lost after prolonged incubation time of propionibacteria and moulds. The nature of the antifungal metabolites was studied in order to clarify the mechanism behind the antifungal activity of the isolated propionibacteria.